



Self-Extensionality of Finitely-Valued Logics: Advances

Alexej Pynko

EasyChair preprints are intended for rapid dissemination of research results and are integrated with the rest of EasyChair.

September 29, 2021

SELF-EXTENSIONALITY OF FINITELY-VALUED LOGICS: ADVANCES

ALEXEJ P. PYNKO

ABSTRACT. We start from proving a general characterization of the self-extensionality of sentential logics implying the decidability of this problem for (not necessarily uniform) finitely-valued logics. And what is more, in case of logics defined by finitely many either implicative or both disjunctive and conjunctive finite *hereditarily* simple (viz., having no non-simple submatrix) matrices, we then derive a characterization yielding a quite effective algebraic criterion of checking their self-extensionality via analyzing homomorphisms between (viz., in the uniform case, endomorphisms of) the underlying algebras of their defining matrices and equally being a quite useful heuristic tool, manual applications of which are demonstrated within the framework of Łukasiewicz' finitely-valued logics, uniform three-valued logics with subclassical negation (U3VLSN), uniform four-valued expansions of Belnap's "useful" four-valued logic as well as their (not necessarily uniform) no-more-than-four-valued extensions, [uniform inferentially consistent proper {in particular, no-more-than-three-valued} non-]classical ones proving to be [non-]self-extensional.

1. INTRODUCTION

Perhaps, the principal value of universal logical investigations consists in discovering uniform points behind particular results originally proved *ad hoc*. This thesis is the main paradigm of the present universal logical study.

Recall that a sentential logic (cf., e.g., [5]) is said to be *self-extensional*, whenever its inter-derivability relation is a congruence of the formula algebra (i.e. is preserved under subformula replacement). This feature is typical of both two-valued (in particular, classical)¹ and super-intuitionistic logics as well as some interesting many-valued ones (like Belnap's "useful" four-valued one [1]). Here, we explore self-extensionality laying a special emphasis onto the general framework of finitely-valued logics and the decidability issue with reducing the complexity of effective procedures of verifying self-extensionality, when restricting our consideration to finitely-valued logics of special kind — namely, those defined by finitely many either implicative or both conjunctive and disjunctive (and so having either classical implication or both classical conjunction and classical disjunction in Tarski's conventional sense) *hereditarily simple* (viz., having no non-simple submatrix; i.e., having an *equality determinant* in a sense extending [14]) finite matrices. We then exemplify our universal elaboration by discussing four (perhaps, most representative) generic classes of logics of the kind involved: Łukasiewicz' finitely-valued logics [6]; uniform three-valued logics with subclassical negation (U3VLSN); uniform four-valued expansions of Belnap's "useful" four-valued logic [1] as well as

2020 *Mathematics Subject Classification*. 03B20, 03B22, 03B50, 03B53, 03G10.

Key words and phrases. logic; matrix; model; congruence; [semi-]lattice.

¹Properly speaking, within the context of General Logic, the notorious classical logic arises just as the clone of miscellaneous functionally complete two-valued logics with classical negation. Here, we follow this natural paradigm, equally adopted in [18] even without the stipulation of functional completeness, calling functionally complete classical logics *genuinely* so, that naturally gives rise to the conception of *subclassical* logic/negation.

their (not necessarily uniform) no-more-than-four-valued extensions, [uniform inferentially consistent proper {in particular, no-more-than-three-valued} non-]classical ones proving to be [non-]self-extensional.

The rest of the paper is as follows. The exposition of the material of the paper is entirely self-contained (of course, modulo very basic issues concerning Set and Lattice Theory, Universal Algebra and Logic to be found, if necessary, in standard mathematical handbooks like [8]). Section 2 is a concise summary of particular basic issues underlying the paper, most of which, though having become a part of algebraic and logical folklore, are still recalled just for the exposition to be properly self-contained. Likewise, in Section 3, we then summarize certain advanced generic issues concerning simple matrices, equality determinants, intrinsic varieties as well as both disjunctivity and implicativity. Section 4 is a collection of main *general* results of the paper that are then exemplified in Section 5 (aside from Łukasiewicz' finitely-valued logics, whose non-self-extensionality has actually been due [15], as we briefly discuss within Example 4.16 — this equally concerns certain particular instances discussed in Section 5 and summarized in Example 4.17). Finally, Section 6 is a brief summary of principal contributions of the paper.

2. BASIC ISSUES

2.1. Set-theoretical background. We follow the standard set-theoretical convention, according to which natural numbers (including 0) are treated as finite ordinals (viz., sets of lesser natural numbers), the ordinal of all them being denoted by ω . In this way, when dealing with n -tuples to be viewed as either [comma separated] sequences of length n or functions with domain n , where $n \in \omega$, π_i , where $i \in n$, denotes the i -th projection operator under enumeration started from rather 0 than 1. (In particular, when $n = 2$, $\pi_{0/1}$ denotes the left/right projection operator, respectively.) The proper class of all ordinals is denoted by ∞ . Also, functions are viewed as binary relations, while singletons are identified with their unique elements, unless any confusion is possible. A function f is said to be *singular*, provided $|\text{img } f| \in 2$, that is, $(\ker f) = (\text{dom } f)^2$.

Given a set S , let $\Delta_S \triangleq \{(a, a) \mid a \in S\}$, functions of such a kind being referred to as *diagonal*, and $\wp_{[K]}(S)$ the set of all subsets of S [of cardinality $\in K \subseteq \infty$], respectively. Then, given any *equivalence relation* θ on S , viz., a *transitive* (in the sense that $(\theta \circ \theta) \subseteq \theta$) *symmetric* (in the sense that $\theta^{-1} \subseteq \theta$) *reflexive binary relation on S* (in the sense that $\Delta_S \subseteq \theta \subseteq S^2$), ν_θ denotes the function with domain S defined by $\nu_\theta(a) \triangleq \theta[\{a\}]$, for all $a \in S$, while $(T/\theta) \triangleq \nu_\theta[T]$, for every $T \subseteq S$. Next, any S -*tuple* (viz., a function with domain S) is often written in the sequence form \bar{t} , its s -th component (viz., the value under argument s), where $s \in S$, being written as t_s , in that case. Given two more sets A and B , any relation $R \subseteq (A \times B)$ (in particular, a mapping $R : A \rightarrow B$) determines the equally-denoted relation $R \subseteq (A^S \times B^S)$ (resp., mapping $R : A^S \rightarrow B^S$) point-wise. Furthermore, any $f : S^n \rightarrow S$, where $n \in \omega$, is said to be *R -monotonic*, where $R \subseteq S^2$, provided, for all $\bar{a} \in R^n$, it holds that $\langle f(\bar{a} \circ \pi_0), f(\bar{a} \circ \pi_1) \rangle \in R$. Then, $\text{Tr}(R) \triangleq \{(\pi_0(a_0), \pi_1(a_{m-1})) \mid m \in (\omega \setminus 1), \bar{a} \in R^m, \forall i \in (m-1) : \pi_1(a_i) = \pi_0(a_{i+1})\}$ is the least transitive binary relation on S including R , called the *transitive closure of R* . Finally, given any $T \subseteq S$, we have the *characteristic function/mapping* $\chi_S^T \triangleq ((T \times \{1\}) \cup ((S \setminus T) \times \{0\})) \in 2^S$ of T in S .

Let A be a set. Then, a $U \subseteq \wp(A)$ is said to be *upward-directed*, provided, for every $S \in \wp_\omega(U)$, there is some $T \in U$ such that $(\bigcup S) \subseteq T$, in which case $U \neq \emptyset$, when taking $S = \emptyset$. Further, a subset of $\wp(A)$ is said to be *inductive*, whenever it is closed under unions of upward-directed subsets. Further, a *closure system over A* is any $\mathcal{C} \subseteq \wp(A)$ such that, for every $S \subseteq \mathcal{C}$, it holds that $(A \cap \bigcap S) \in \mathcal{C}$. In that

case, any $\mathcal{B} \subseteq \mathcal{C}$ is called a (*closure*) *basis* of \mathcal{C} , provided $\mathcal{C} = \{A \cap \bigcap S \mid S \subseteq \mathcal{B}\}$. Furthermore, an *operator over* A is any unary operation O on $\wp(A)$. This is said to be *monotonic*, whenever it is $(\subseteq \cap \wp(A)^2)$ -monotonic. Likewise, it is said to be *idempotent|transitive*, provided, for all $X \subseteq A$, it holds that $(X \mid O(O(X))) \subseteq O(X)$, respectively. Finally, it is said to be *inductive|finitary*, provided, for any upward-directed $U \subseteq \wp(A)$, it holds that $O(\bigcup U) \subseteq (\bigcup O[U])$. Then, a *closure operator over* A is any monotonic idempotent transitive operator over A , in which case $\text{img } C$ is a [n inductive] closure system over A [iff C is inductive], determining C uniquely, as, for every basis \mathcal{B} of $\text{img } C$ (in particular, $\text{img } C$ itself) and each $X \subseteq A$, $C(X) = (A \cap \bigcap \{Y \in \mathcal{B} \mid X \subseteq Y\})$, C and $\text{img } C$ being said to be *dual* to one another.

2.2. Algebraic background. Unless otherwise specified, abstract algebras are denoted by Fraktur letters [possibly, with indices], their carriers (viz., underlying sets) being denoted by corresponding Italic letters [with same indices, if any].

A (*propositional|sentential*) *language|signature* is any algebraic (viz., functional) signature Σ (to be dealt with throughout the paper by default) constituted by function (viz., operation) symbols of finite arity to be treated as (*propositional|sentential*) [*primary*] *connectives*, the set of all nullary ones being denoted by $\Sigma \mid 0$.

Given a Σ -algebra \mathfrak{A} , the set $\text{Con}(\mathfrak{A})$ of all *congruences of* \mathfrak{A} (viz., equivalence relations θ on A such that *primary* operations of \mathfrak{A} — i.e., those of the form $\varsigma^{\mathfrak{A}}$, where $\varsigma \in \Sigma$ — are θ -monotonic) is an inductive closure system over A^2 , the dual closure operator (of congruence generation) being denoted by $\text{Cg}^{\mathfrak{A}}$. Then, given any function f with $(\text{dom } f) = A$ and $(\text{ker } f) \in \text{Con}(\mathfrak{A})$, we have the Σ -algebra $f[\mathfrak{A}]$ with carrier $f[A]$ and primary operations $\varsigma^{f[\mathfrak{A}]} \triangleq f[\varsigma^{\mathfrak{A}}]$, where $\varsigma \in \Sigma$. In particular, given any $\theta \in \text{Con}(\mathfrak{A})$, $(\mathfrak{A}/\theta) \triangleq \nu_{\theta}[\mathfrak{A}]$ is known as the *quotient of* \mathfrak{A} *by* θ . Finally, given a class \mathbf{K} of Σ -algebras, set $\text{hom}(\mathfrak{A}, \mathbf{K}) \triangleq (\bigcup \{\text{hom}(\mathfrak{A}, \mathfrak{B}) \mid \mathfrak{B} \in \mathbf{K}\})$, in which case $\text{ker}[\text{hom}(\mathfrak{A}, \mathbf{K})] \subseteq \text{Con}(\mathfrak{A})$, so $(A^2 \cap \bigcap \text{ker}[\text{hom}(\mathfrak{A}, \mathbf{K})]) \in \text{Con}(\mathfrak{A})$.

Given any *rank*, viz., $\alpha \subseteq \omega$, put $\bar{x}_{\alpha} \triangleq \langle x_i \rangle_{i \in \alpha}$ and $\text{Var}_{\alpha} \triangleq (\text{img } \bar{x}_{\alpha})$, elements of which being viewed as (*propositional|sentential*) *variables of rank* α . (In general, any mention of rank α within any context is normally omitted, whenever $\alpha = \omega$.) Then, providing either $\alpha \neq \emptyset$ or Σ has a nullary connective, in which case α is called a Σ -*rank*, we have the absolutely-free Σ -algebra $\mathfrak{Fm}_{\Sigma}^{\alpha}$ freely-generated by the set Var_{α} , “its endomorphisms”/“elements of its carrier $\text{Fm}_{\Sigma}^{\alpha}$ (viz., Σ -terms of rank α)” being called (*propositional|sentential*) Σ -*substitutions/-formulas of* $\{\Sigma\}$ -*rank* α . Any homomorphism h from $\mathfrak{Fm}_{\Sigma}^{\alpha}$ to a Σ -algebra $\mathfrak{A}(= \mathfrak{Fm}_{\Sigma}^{\alpha})$ is uniquely determined by {and so identified with} $h' = (h \upharpoonright (\text{Var}_{\alpha} \setminus V))$ (where $V \subseteq \text{Var}_{\alpha}$ such that $h \upharpoonright V$ is diagonal) as well as often written in the standard assignment (resp., substitution) form $[v/h(v)]_{v \in (\text{dom } h')}$, $\varphi^{\mathfrak{A}} \langle \upharpoonright h \upharpoonright \rangle$, where $\varphi \in \text{Fm}_{\Sigma}^{\alpha}$, standing for $h(\varphi)$ (the algebra superscript being normally omitted just like in denoting primary operations of \mathfrak{A}). Then, given any $n \in \omega$, a *secondary n -ary connective of* Σ is any Σ -formula φ of rank $m_{\Sigma}(n) \triangleq (n + (1 - \min(1, \max(n, |\Sigma \setminus 0|))))$, in which case, given any Σ -algebra \mathfrak{A} , an $f : A^n \rightarrow A$ is said to be *secondary/“(term-wise) definable {by } φ ” of/in* \mathfrak{A} , provided, for all $\bar{a} \in A^{m_{\Sigma}(n)}$, it holds that $f(\bar{a} \upharpoonright n) = \varphi^{\mathfrak{A}}[x_i/a_i]_{i \in m_{\Sigma}(n)}$. For the sake of formal unification, any primary n -ary connective $\varsigma \in \Sigma$ is identified with the secondary one $\varsigma(\bar{x}_n)$. A $\theta \in \text{Con}(\mathfrak{Fm}_{\Sigma}^{\alpha})$ is said to be *fully-invariant*, if, for every Σ -substitution σ of rank α , it holds that $\sigma[\theta] \subseteq \theta$. Recall that, for any [surjective] $h \in \text{hom}(\mathfrak{A}, \mathfrak{B})$, where \mathfrak{A} and \mathfrak{B} are Σ -algebras, it holds that:

$$(2.1) \quad [\text{hom}(\mathfrak{Fm}_{\Sigma}^{\alpha}, \mathfrak{B}) \subseteq] \{h \circ g \mid g \in \text{hom}(\mathfrak{Fm}_{\Sigma}^{\alpha}, \mathfrak{A})\} \subseteq \text{hom}(\mathfrak{Fm}_{\Sigma}^{\alpha}, \mathfrak{B}).$$

Any $\langle \phi, \psi \rangle \in \text{Eq}_{\Sigma}^{\alpha} \triangleq (\text{Fm}_{\Sigma}^{\alpha})^2$ is referred to as a Σ -*equation/-identity of* $\{\Sigma\}$ -*rank* α and normally written in the standard equational form $\phi \approx \psi$. In this way, given any $h \in \text{hom}(\mathfrak{Fm}_{\Sigma}^{\alpha}, \mathfrak{A})$, $\text{ker } h$ is the set of all Σ -identities of rank α *true/satisfied in* \mathfrak{A} *under* h . Likewise, given a class \mathbf{K} of Σ -algebras, $\theta_{\mathbf{K}}^{\alpha} \triangleq (\text{Eq}_{\Sigma}^{\alpha} \cap \bigcap \text{ker}[\text{hom}(\mathfrak{Fm}_{\Sigma}^{\alpha}, \mathbf{K})]) \in$

$\text{Con}(\mathfrak{Fm}_\Sigma^\alpha)$, being fully invariant, in view of (2.1), is the set of all all Σ -identities of rank α *true/satisfied in* \mathbf{K} , in which case we set $\mathfrak{F}_\mathbf{K}^\alpha \triangleq (\mathfrak{Fm}_\Sigma^\alpha/\theta_\mathbf{K}^\alpha)$. (In case α as well as both \mathbf{K} and all elements of it are finite, the class $I \triangleq \{\langle \mathfrak{A}, h \rangle \mid \mathfrak{A} \in \mathbf{K}, h \in \text{hom}(\mathfrak{Fm}_\Sigma^\alpha, \mathfrak{A})\}$ is a finite set — more precisely, $|I| = \sum_{\mathfrak{A} \in \mathbf{K}} |\mathfrak{A}|^\alpha$, in which case, putting, for each $i \in I$, $\mathfrak{A}_i \triangleq \pi_0(i) \in \mathbf{K}$, $h_i \triangleq \pi_1(i) \in \text{hom}(\mathfrak{Fm}_\Sigma^\alpha, \mathfrak{A}_i)$ and $\mathfrak{B}_i \triangleq (\mathfrak{A}_i \upharpoonright (\text{img } h_i))$, we have $\text{hom}(\mathfrak{Fm}_\Sigma^\alpha, \prod_{i \in I} \mathfrak{B}_i) \ni g : \text{Fm}_\Sigma^\alpha \rightarrow (\prod_{i \in I} \mathfrak{B}_i), \varphi \mapsto \langle h_i(\varphi) \rangle_{i \in I}$ with $(\ker g) = \theta \triangleq \theta_\mathbf{K}^\alpha$, and so, by the Homomorphism Theorem, $e \triangleq (\nu_\theta^{-1} \circ g)$ is an isomorphism from $\mathfrak{F}_\mathbf{K}^\alpha$ onto the subdirect product $(\prod_{i \in I} \mathfrak{B}_i) \upharpoonright (\text{img } g)$ of $\langle \mathfrak{B}_i \rangle_{i \in I}$. In this way, the former is finite, for the latter is so — more precisely, $|F_\mathbf{K}^\alpha| \leq (\max\{|\mathfrak{A}| \mid \mathfrak{A} \in \mathbf{K}\})^{|I|}$.)

Given a Σ -algebra \mathfrak{A} and any $a, b \in A$, according to Mal'cev [Principal Congruence] Lemma [7]:

$$(2.2) \quad \text{Cg}^\mathfrak{A}(\langle a, b \rangle) = \text{Tr}(\nabla^\mathfrak{A}(a, b) \cup \nabla^\mathfrak{A}(a, b)^{-1}), \text{ where } \nabla^\mathfrak{A}(a, b) \triangleq \{ \langle \varphi^\mathfrak{A}[x_i/c_i; x_n/a]_{i \in n}, \varphi^\mathfrak{A}[x_i/c_i; x_n/b]_{i \in n} \mid n \in \omega, \varphi \in \text{Fm}_\Sigma^{n+1}, \bar{c} \in A^n \}.$$

The class of all Σ -algebras satisfying every element of an $\mathcal{E} \subseteq \text{Eq}_\Sigma^\omega$ is called the *variety axiomatized by* \mathcal{E} . Then, the variety $\mathbf{V}(\mathbf{K})$ axiomatized by $\theta_\mathbf{K}^\omega$ is the least variety including \mathbf{K} and is said to be *generated by* \mathbf{K} , in which case $\theta_{\mathbf{V}(\mathbf{K})}^\alpha = \theta_\mathbf{K}^\alpha$, and so $\mathfrak{F}_{\mathbf{V}(\mathbf{K})}^\alpha = \mathfrak{F}_\mathbf{K}^\alpha$.

Given a fully invariant $\theta \in \text{Con}(\mathfrak{Fm}_\Sigma^\omega)$, by (2.1), $\mathfrak{Fm}_\Sigma^\omega/\theta$ belongs to the variety \mathbf{V} axiomatized by θ , in which case any Σ -identity satisfied in \mathbf{V} belongs to θ , and so $\theta_\mathbf{V}^\omega = \theta$. In particular, given a variety \mathbf{V} of Σ -algebras, we have $\mathfrak{F}_\mathbf{V}^\alpha \in \mathbf{V}$.

Finally, let $\text{Var} : \text{Fm}_\Sigma^\omega \rightarrow \wp_\omega(\text{Var}_\omega)$ be the mapping assigning the set of all *actually* occurring variables.

2.2.1. Lattice-theoretic background.

2.2.1.1. Semi-lattices. Let \diamond be a (possibly, secondary) binary connective of Σ .

A Σ -algebra \mathfrak{A} is called a \diamond -*semi-lattice*, provided it satisfies *semi-lattice* identities for \diamond (viz., *idempotence* $(x_0 \diamond x_0) \approx x_0$, *commutativity* $(x_0 \diamond x_1) \approx (x_1 \diamond x_0)$ and *associativity* $(x_0 \diamond (x_1 \diamond x_2)) \approx ((x_0 \diamond x_1) \diamond x_2)$ ones), in which case we have the partial ordering $\leq_\diamond^\mathfrak{A}$ on A , given by $(a \leq_\diamond^\mathfrak{A} b) \stackrel{\text{def}}{\iff} (a = (a \diamond b))$, for all $a, b \in A$. Then, in case the [dual] poset $\langle A, (\leq_\diamond^\mathfrak{A})^{-1} \rangle$ has the least element (viz., *lower bound*), this is called the [dual] $\langle \diamond \rangle$ -*bound of* \mathfrak{A} and denoted by $[\delta] \beta_\diamond^\mathfrak{A}$, while \mathfrak{A} is referred to as a \diamond -*semi-lattice with [dual] bound* $\{a, \text{ whenever } a = [\delta] \beta_\diamond^\mathfrak{A}\}$.

Lemma 2.1. *Let \mathfrak{A} and \mathfrak{B} be \diamond -semi-lattices with bound and $h \in \text{hom}(\mathfrak{A}, \mathfrak{B})$. Suppose $h[A] = B$. Then, $h(\beta_\diamond^\mathfrak{A}) = \beta_\diamond^\mathfrak{B}$.*

Proof. There is some $a \in A$ such that $h(a) = \beta_\diamond^\mathfrak{B}$, in which case $(a \diamond^\mathfrak{A} \beta_\diamond^\mathfrak{A}) = \beta_\diamond^\mathfrak{A}$, so $h(\beta_\diamond^\mathfrak{A}) = (h(a) \diamond^\mathfrak{B} h(\beta_\diamond^\mathfrak{A})) = (\beta_\diamond^\mathfrak{B} \diamond^\mathfrak{B} h(\beta_\diamond^\mathfrak{A})) = \beta_\diamond^\mathfrak{B}$, as required. \square

2.2.1.2. Lattices. Let $\bar{\vee}$ and $\underline{\vee}$ be (possibly, secondary) binary connectives of Σ .

A Σ -algebra \mathfrak{A} is called a [distributive] $(\bar{\vee}, \underline{\vee})$ -*lattice*, provided it satisfies [distributive] *lattice* identities for $\bar{\vee}$ and $\underline{\vee}$ (viz., *semi-lattice* identities for both $\bar{\vee}$ and $\underline{\vee}$ as well as *mutual [both] absorption* $(x_0 \diamond_1 (x_0 \diamond_2 x_1)) \approx x_0$ [and *distributivity* $(x_0 \diamond_1 (x_1 \diamond_2 x_2)) \approx ((x_0 \diamond_1 x_1) \diamond_2 (x_0 \diamond_2 x_2))$] identities for $\bar{\vee}$ and $\underline{\vee}$, where $\diamond : 2 \rightarrow \{\bar{\vee}, \underline{\vee}\}$ is either bijection), in which case $\leq_{\bar{\vee}}^\mathfrak{A}$ and $\leq_{\underline{\vee}}^\mathfrak{A}$ are inverse/dual to one another, and so, in case \mathfrak{A} is a $\underline{\vee}$ -semi-lattice with bound (in particular, when A is finite), $\beta_{\underline{\vee}}^\mathfrak{A}$ is the dual $\bar{\vee}$ -bound of \mathfrak{A} (viz., the greatest element of the poset $\langle A, \leq_{\bar{\vee}}^\mathfrak{A} \rangle$). Then, in case \mathfrak{A} is a {distributive} $(\bar{\vee}, \underline{\vee})$ -lattice, it is said to be that *with zero/unit* (a) , whenever it is a $(\bar{\vee}|\underline{\vee})$ -semi-lattice with bound (a) .

2.2.1.2.1. Bounded lattices. Let $\Sigma_{\langle \emptyset \rangle \{+\} [01]} \triangleq (\emptyset \{ \cup \{ \wedge, \vee \} \} [\cup \{ \perp, \top \}])$ be the $\{[\text{bounded}] \text{ lattice}\}$ signature $\{\text{with binary } \wedge \text{ (conjunction) and } \vee \text{ (disjunction)}\}$ $[\{\text{as well as}\}]$ with nullary \perp and \top (falsehood/zero and truth/unit constants, respectively)]. Then, a $\Sigma_{+[01]}$ -algebra \mathfrak{A} is called a *[bounded] (distributive) lattice*, whenever it is a (distributive) (\wedge, \vee) -lattice [with zero $\perp^{\mathfrak{A}}$ and unit $\top^{\mathfrak{A}}$]. Given any $\Sigma \supseteq \Sigma_{+[01]}$ and any $\phi, \psi \in \text{Fm}_{\Sigma}^{\omega}$, $\phi \lesssim \psi$ stands for $\phi \approx (\phi \wedge \psi)$. Likewise, given any Σ -algebra \mathfrak{A} with Σ_+ -reduct being a lattice, $\leq^{\mathfrak{A}}$ stands for $\leq_{\wedge}^{\mathfrak{A}}$. Then, given any $n \in (\omega \setminus 2)$, $\mathfrak{D}_{n[01]}$ denotes the [bounded] distributive lattice with carrier $(n \div (n-1)) \triangleq \{ \frac{m}{n-1} \mid m \in n \}$ and $\leq^{\mathfrak{D}_{n[01]}} \triangleq (\leq \cap D_{n[01]}^2)$.

2.3. Logical background.

2.3.1. *Propositional calculi and logics.* A (*propositional*||*sentential*) [*finitary*|*unary*]*axiomatic* Σ -*rule/-calculus* $\{\text{of } \langle \Sigma - \rangle\text{-rank } \alpha\}$ is any element/subset of the set $\wp_{[\omega | (2 \setminus 1) | 1]}(\text{Fm}_{\Sigma}^{\omega \{ \cap \alpha \}} \times \text{Fm}_{\Sigma}^{\omega \{ \cap \alpha \}})$, any Σ -rule $\langle \Gamma, \varphi \rangle$ being normally written in the standard sequent form $\Gamma \vdash \varphi$, “the left”/“any element of the right” component|side of it being referred to as the/a *conclusion/premise* of it. Then, we set $\sigma(\Gamma \vdash \varphi) \triangleq (\sigma[\Gamma] \vdash \sigma(\varphi))$, where σ is a Σ -substitution. Axiomatic Σ -rules are called (*propositional/sentential*) Σ -*axioms* and are identified with their conclusions.

A (*propositional/sentential*) Σ -*logic* is any closure operator C over $\text{Fm}_{\Sigma}^{\omega}$ that is *structural* in the sense that $\sigma[C(X)] \subseteq C(\sigma[X])$, for all $X \subseteq \text{Fm}_{\Sigma}^{\omega}$ and all $\sigma \in \text{hom}(\mathfrak{Fm}_{\Sigma}^{\omega}, \mathfrak{Fm}_{\Sigma}^{\omega})$, that is, $\text{img } C$ is closed under inverse Σ -substitutions. Then, we have the equivalence relation $\equiv_C^{\alpha} \triangleq \{ \langle \phi, \psi \rangle \in \text{Eq}_{\Sigma}^{\alpha} \mid C(\phi) = C(\psi) \}$ on $\text{Fm}_{\Sigma}^{\alpha}$, where $\alpha \in \wp_{\infty[1]}(\omega)$ [unless Σ has a nullary connective], called the *inter-derivability relation* of C , whenever $\alpha = \omega$. A *congruence* of C is any $\theta \in \text{Con}(\mathfrak{Fm}_{\Sigma}^{\omega})$ such that $\theta \subseteq \equiv_C^{\omega}$, the set of all them being denoted by $\text{Con}(C)$. Then, given any $\theta, \vartheta \in \text{Con}(C)$, $\text{Tr}(\theta \cup \vartheta)$, being well-known to be a congruence of $\mathfrak{Fm}_{\Sigma}^{\omega}$, is then that of C , for θ_C^{ω} , being an equivalence relation, is transitive. In particular, any maximal congruence of C (that exists, by Zorn Lemma, because $\text{Con}(C) \ni \Delta_{\text{Fm}_{\Sigma}^{\omega}}$ is both non-empty and inductive, for $\text{Con}(\mathfrak{Fm}_{\Sigma}^{\omega})$ is so) is the greatest one to be denoted by $\wp(C)$. Then, C is said to be *self-extensional*, whenever $\equiv_C^{\omega} \in \text{Con}(\mathfrak{Fm}_{\Sigma}^{\omega})$.

Definition 2.2 (cf. [12]). Given a Σ -logic C , the variety $\text{IV}(C)$ axiomatized by $\wp(C)$ is called the *intrinsic variety* of C . \square

Next, a Σ -logic C is said to be [*inferentially*] (*in*)*consistent*, provided $x_1 \notin (\in)C(\emptyset \cup \{x_0\})$ [(in which case $\equiv_C^{\omega} = \text{Eq}_{\Sigma}^{\omega} \in \text{Con}(\mathfrak{Fm}_{\Sigma}^{\omega})$, and so C is self-extensional)]. Further, a Σ -rule $\Gamma \rightarrow \Phi$ is said to be *satisfied/derivable* in C , provided $\Phi \in C(\Gamma)$, Σ -axioms satisfied in C being referred to as *theorems* of C .

Definition 2.3. A Σ -logic C' is said to be a (*proper*) [*K*-]*extension* of a Σ -logic C [where $K \subseteq \infty$], whenever ($C' \neq C$ and) $C(X) \subseteq C'(X)$, for all $X \in \wp_{[K]}(\text{Fm}_{\Sigma}^{\omega})$, C being said to be a (*proper*) [*K*-]*sublogic* of C' , in which case C' is said to be *axiomatized* by a Σ -calculus \mathcal{C} *relatively* to C , whenever C' is the least (w.r.t. the extension partial ordering) extension of C satisfying every rule in \mathcal{C} . \square

Next, a Σ -logic C is said to be (*strongly*)/*weakly* $\{\text{classically}\}$ $\bar{\wedge}$ -*conjunctive*, provided $C(\{x_0, x_1\}) = / \subseteq C(x_0 \bar{\wedge} x_1)$. Likewise, C is said to be (*strongly*)/*weakly* $\{\text{classically}\}$ $\bar{\vee}$ -*disjunctive*, if $C(X \cup \{\phi \bar{\vee} \psi\}) = / \subseteq (C(X \cup \{\phi\}) \cap C(X \cup \{\psi\}))$, where $(X \cup \{\phi, \psi\}) \subseteq \text{Fm}_{\Sigma}^{\omega}$, “in which case”/“that is, the first two — viz., (2.3) — of” the following four rules:

$$(2.3) \quad x_i \vdash (x_0 \bar{\vee} x_1), \quad \text{where } i \in 2,$$

$$(2.4) \quad (x_0 \bar{\vee} x_1) \vdash (x_1 \bar{\vee} x_0),$$

$$(2.5) \quad (x_0 \bar{\vee} x_0) \vdash x_0$$

are satisfied in C . Further, C is said to *have/satisfy Deduction Theorem (DT) with respect to* a (possibly, secondary) binary connective \sqsupset of Σ (fixed throughout the paper by default), provided, for all $\phi \in X \subseteq \text{Fm}_\Sigma^\omega$ and all $\psi \in C(X)$, it holds that $(\phi \sqsupset \psi) \in C(X \setminus \{\phi\})$, in which case the following axioms:

$$(2.6) \quad x_0 \sqsupset x_0,$$

$$(2.7) \quad x_0 \sqsupset (x_1 \sqsupset x_0)$$

are satisfied in C . Then, C is said to be *weakly {classically} \sqsupset -implicative*, if it has DT w.r.t. \sqsupset as well as satisfies the *Modus Ponens* rule:

$$(2.8) \quad \{x_0, x_0 \sqsupset x_1\} \vdash x_1,$$

in which case the following axiom:

$$(2.9) \quad (x_0 \uplus_{\sqsupset} (x_0 \sqsupset x_1)),$$

where $(x_0 \uplus_{\sqsupset} x_1) \triangleq ((x_0 \sqsupset x_1) \sqsupset x_1)$ is the *intrinsic disjunction of (implication) \sqsupset* , is satisfied in C . Likewise, C is said to be *(strongly) {classically} \sqsupset -implicative*, whenever it is weakly so and satisfies the *Peirce Law* axiom:

$$(2.10) \quad ((x_0 \sqsupset x_1) \uplus_{\sqsupset} x_0).$$

Furthermore, C is said to be *\wr -paraconsistent*, where \wr is a (possibly, secondary) unary connective of Σ (tacitly fixed throughout the paper by default), provided it does not satisfy the *Ex Contradictione Quodlibet* rule:

$$(2.11) \quad \{x_0, \wr x_0\} \vdash x_1.$$

Likewise, C is said to be *\vee, \wr -paracomplete*, whenever it does not satisfy the *Excluded Middle Law* axiom:

$$(2.12) \quad x_0 \vee \wr x_0.$$

Given any $\Sigma' \subseteq \Sigma$, the Σ' -logic C' , defined by $C'(X) \triangleq (\text{Fm}_{\Sigma'}^\omega \cap C(X))$, for all $X \subseteq \text{Fm}_{\Sigma'}^\omega$, is called the *Σ' -fragment of C* , C being referred to as a *$(\Sigma'$ -)expansion of C'* , in which case $\equiv_{C'}^\omega = (\equiv_C^\omega \cap \text{Eq}_{\Sigma'}^\omega)$, and so C' is self-extensional, whenever C is so. Finally, C is said to be *theorem-less/purely-inferential*, whenever it has no theorem, that is, $\emptyset \in (\text{img } C)$. In general, $(\text{img } C) \cup \{\emptyset\}$ is closed under inverse Σ -substitutions, for $\text{img } C$ is so, in which case the dual closure operator C_{+0} is the greatest purely-inferential sublogic of C , called the *purely-inferential version of C* and being an $(\infty \setminus 1)$ -extension of C (cf. Definition 2.3), so

$$(2.13) \quad \equiv_C^\omega = \equiv_{C_{+0}}^\omega$$

(in particular, C_{+0} is self-extensional iff C is so).

2.3.2. Logical matrices. A (logical) Σ -matrix is any pair of the form $\mathcal{A} = (\mathfrak{A}, D^{\mathcal{A}})$, where \mathfrak{A} is a Σ -algebra, called the *underlying algebra of \mathcal{A}* , while A is called the *carrier/“underlying set” of \mathcal{A}* , whereas $D^{\mathcal{A}} \subseteq A$ is called the *truth predicate of \mathcal{A}* , elements of $A[\cap D^{\mathcal{A}}]$ being referred to as *[distinguished] values of \mathcal{A}* . (In general, matrices are denoted by Calligraphic letters [possibly, with indices], their underlying algebras being denoted by corresponding capital Fraktur letters [with same indices, if any].) This is said to be *[no-more/less-than-] n -valued*, where $n \in (\omega \setminus 1)$, provided $|A| = [\leq/\geq]n$. Next, it is said to be *[in]consistent*, whenever $D^{\mathcal{A}} \neq [=]A$, respectively. Likewise, it is said to be *truth[-non]-empty*, whenever $D^{\mathcal{A}} = [\neq]\emptyset$. Further, it is said to be *truth-/false-singular*, provided $|((D^{\mathcal{A}}/(A \setminus D^{\mathcal{A}}))|) \in 2$. Finally, \mathcal{A} is said to be *finite[ly generated]/“generated by $B \subseteq A$ ”*, if \mathfrak{A} is so.

Given any Σ -rank α and any class \mathbf{M} of Σ -matrices, we have the closure operator $\text{Cn}_{\mathbf{M}}^{\alpha}$ over $\text{Fm}_{\Sigma}^{\alpha}$ dual to the closure system with basis $\mathcal{B}_{\mathbf{M}}^{\alpha} \triangleq \{h^{-1}[D^{\mathcal{A}}] \mid \mathcal{A} \in \mathbf{M}, h \in \text{hom}(\mathfrak{Fm}_{\Sigma}^{\alpha}, \mathfrak{A})\}$, in which case:

$$(2.14) \quad \text{Cn}_{\mathbf{M}}^{\alpha}(X) = (\text{Fm}_{\Sigma}^{\alpha} \cap \text{Cn}_{\mathbf{M}}^{\omega}(X)),$$

for all $X \subseteq \text{Fm}_{\Sigma}^{\alpha}$. Then, by (2.1), $\text{Cn}_{\mathbf{M}}^{\omega}$ is a Σ -logic, called the *logic of/“defined by”* \mathbf{M} . A Σ -logic is said to be *{“unitary/uniform[ly]”|double|finitely} (no-more-than-) n -valued*, where $n \in (\omega \setminus 1)$, whenever it is defined by a *{one-element|two-element|finite} class of (no-more-than-) n -valued Σ -matrices* {in which case it is finitary, and so is the logic of any finite class of finite Σ -matrices; cf. [5]}. Then, a [uniformly] n -valued Σ -logic, where $n \in (\omega \setminus 2)$, is said to be *minimally* so, unless it is [uniformly] no-more-than- $(n - 1)$ -valued.

As usual, Σ -matrices are treated as first-order model structures (viz., algebraic systems; cf. [8]) of the first-order signature $\Sigma \cup \{D\}$ with unary predicate D , any [in]finitary Σ -rule $\Gamma \vdash \phi$ being viewed as the [in]finitary equality-free basic strict Horn formula $(\bigwedge \Gamma) \rightarrow \phi$ under the standard identification of any propositional Σ -formula ψ with the first-order atomic formula $D(\psi)$, as well as being *true/satisfied* in a class \mathbf{M} of Σ -matrices (in the conventional model-theoretic sense; cf., e.g., [8]) iff it being satisfied in the logic of \mathbf{M} .

Remark 2.4. Since any [non-]axiomatic rule is [not] non-true in any truth-empty matrix, given any class \mathbf{M} of Σ -matrices, the purely-inferential version of the logic of \mathbf{M} is defined by MUS , where \mathbf{S} is any non-empty class of truth-empty Σ -matrices. \square

Let \mathcal{A} and \mathcal{B} be two Σ -matrices. A *(strict) [surjective] {injective} homomorphism from \mathcal{A} [on]to \mathcal{B}* is any *{injective} $h \in \text{hom}(\mathfrak{A}, \mathfrak{B})$ such that $h[A] = B$ and $D^{\mathcal{A}} \subseteq h^{-1}[D^{\mathcal{B}}](\subseteq D^{\mathcal{A}})$* , the set of all them being denoted by $\text{hom}_{\mathbf{S}}^{[\mathbf{S}]}(\mathcal{A}, \mathcal{B})$, in which case \mathcal{B}/\mathcal{A} is said to be a *(strictly) [surjectively] {injectively} homomorphic image/counter-image* ([{as well as called an *isomorphic copy*})] of \mathcal{A}/\mathcal{B} , respectively. Then, by (2.1), we have:

$$(2.15) \quad (\text{hom}_{\mathbf{S}}^{[\mathbf{S}]}(\mathcal{A}, \mathcal{B}) \neq \emptyset) \Rightarrow (\text{Cn}_{\mathbf{B}}^{\alpha}(X) \subseteq \text{Cn}_{\mathcal{A}}^{\alpha}(X) [\subseteq \text{Cn}_{\mathbf{B}}^{\alpha}(X)]),$$

for all Σ -ranks α and all $X \subseteq \text{Fm}_{\Sigma}^{\alpha}$. Further, $\mathcal{A}[\neq \mathcal{B}]$ is said to be a *[proper] submatrix of \mathcal{B}* , whenever $\Delta_{\mathcal{A}} \in \text{hom}_{\mathbf{S}}(\mathcal{A}, \mathcal{B})$, in which case we set $(\mathcal{B} \upharpoonright \mathcal{A}) \triangleq \mathcal{A}$. Injective/bijective strict homomorphisms from \mathcal{A} to \mathcal{B} are called *embeddings/isomorphisms of/from \mathcal{A} into/onto \mathcal{B}* , in case of existence of which \mathcal{A} is said to be *embeddable/isomorphic into/to \mathcal{B}* .

Given a Σ -matrix \mathcal{A} , $(\chi^{\mathcal{A}}/\theta^{\mathcal{A}}) \triangleq (\chi_{\mathcal{A}}^{D^{\mathcal{A}}})/(\ker \chi^{\mathcal{A}})$ is referred to as the *characteristic function/relation of \mathcal{A}* . Then, any $\theta \in \text{Con}(\mathfrak{A})$ such that $\theta \subseteq \theta^{\mathcal{A}}$, in which case ν_{θ} is a strict surjective homomorphism from \mathcal{A} onto $(\mathcal{A}/\theta) \triangleq \langle \mathfrak{A}/\theta, D^{\mathcal{A}}/\theta \rangle$, is called a *congruence of \mathcal{A}* , the set of all them being denoted by $\text{Con}(\mathcal{A})$. Given any $\theta, \vartheta \in \text{Con}(\mathcal{A})$, $\text{Tr}(\theta \cup \vartheta)$, being well-known to be a congruence of \mathfrak{A} , is then that of \mathcal{A} , for $\theta^{\mathcal{A}}$, being an equivalence relation, is transitive. In particular, any maximal congruence of \mathcal{A} (that exists, by Zorn Lemma, because $\text{Con}(\mathcal{A}) \ni \Delta_{\mathcal{A}}$ is both non-empty and inductive, for $\text{Con}(\mathfrak{A})$ is so) is the greatest one to be denoted by $\mathfrak{D}(\mathcal{A})$, in which case, by (2.2), for all $a, b \in \mathcal{A}$, $(a \mathfrak{D}(\mathcal{A}) b) \Leftrightarrow (\nabla^{\mathfrak{A}}(a, b) \subseteq \theta^{\mathcal{A}})$, and so this is traditionally called the *Leibniz congruence of \mathcal{A}* and denoted, but for quite unclear reasons, by rather $\Omega(\mathcal{A})$ than, e.g., $\Lambda(\mathcal{A})$ (here we though naturally adapt conventions adopted in [18] to use its results *immediately*). Finally, \mathcal{A} is said to be *[(finitely) hereditarily] simple*, whenever it has no non-diagonal congruence [as well as no non-simple (finitely-generated) submatrix].

Remark 2.5. Let \mathcal{A} and \mathcal{B} be two Σ -matrices and $h \in \text{hom}(\mathcal{A}, \mathcal{B})$ strict [and surjective]. Then, $\theta^{\mathcal{A}} = h^{-1}[\theta^{\mathcal{B}}]$ and, for every $\theta \in \text{Con}(\mathfrak{B})$, $(\ker h) \subseteq h^{-1}[\theta] \in \text{Con}(\mathfrak{A})$

[while $h[\theta^A] = \theta^B$ as well as $h[h^{-1}[\theta]] = \theta$, whereas, for every $(\ker h) \subseteq \vartheta \in \text{Con}(\mathfrak{A})$, $h[\vartheta] \in \text{Con}(\mathfrak{B})$ and $h^{-1}[h[\vartheta]] = \vartheta$]. Therefore:

- (i) for every $\theta \in \text{Con}(\mathfrak{B})$, $(\ker h) \subseteq h^{-1}[\theta] \in \text{Con}(\mathfrak{A})$ [while $h[h^{-1}[\theta]] = \theta$, whereas, for every $(\ker h) \subseteq \vartheta \in \text{Con}(\mathfrak{A})$, $h[\vartheta] \in \text{Con}(\mathfrak{B})$ and $h^{-1}[h[\vartheta]] = \vartheta$].

In particular (when $\theta = \Delta_B$), we have $(\ker h) = h^{-1}[\Delta_B] \in \text{Con}(\mathfrak{A})$, in which case we get $(\ker h) \subseteq \varnothing(\mathfrak{A})$, and so:

- (ii) h is injective, whenever \mathfrak{A} is simple.

[Likewise, when $\vartheta = \varnothing(\mathfrak{A}) \supseteq (\ker h)$ and $\theta = \varnothing(\mathfrak{B})$, we have $h[\vartheta] \in \text{Con}(\mathfrak{B})$ and $h^{-1}[\theta] \in \text{Con}(\mathfrak{A})$, in which case we get $h[\vartheta] \subseteq \theta$ and $h^{-1}[\theta] \subseteq \vartheta$, and so:

- (iii) $\varnothing(\mathfrak{A}) = h^{-1}[\varnothing(\mathfrak{B})]$ and $\varnothing(\mathfrak{B}) = h[\varnothing(\mathfrak{A})]$.

In particular, when $\mathfrak{B} = (\mathfrak{A}/\vartheta)$ and $h = \nu_\vartheta$, we have $\theta = h[\vartheta] = \Delta_B$, and so:

- (iv) $\mathfrak{A}/\varnothing(\mathfrak{A})$ is simple.] □

Definition 2.6. A Σ -matrix \mathcal{A} is said to be a $[K-]$ model of a Σ -logic C {over \mathfrak{A} } [where $K \subseteq \infty$], provided C is a $[K-]$ sublogic the logic of \mathcal{A} (cf. Definition 2.3), the class of all (simple of) them being denoted by $\text{Mod}_{[K]}^{(*)}(C\{\mathfrak{A}\})$, respectively. Then, $\text{Fi}_C(\mathfrak{A}) \triangleq \pi_1[\text{Mod}(C, \mathfrak{A})]$, whose elements are called *filters of C over \mathfrak{A}* , is a closure system over A , $\text{Fg}_C^{\mathfrak{A}}$ denoting the dual closure operator (of filter generation). □

A Σ -matrix \mathcal{A} is said to be \wr -paraconsistent/ (\vee, \wr) -paracomplete, whenever its logic is so. Next, \mathcal{A} is said to be (strongly)/weakly {classically} \diamond -conjunctive, provided $(\{a, b\} \subseteq D^{\mathcal{A}}) \Leftrightarrow / \Leftarrow ((a \diamond^{\mathfrak{A}} b) \in D^{\mathcal{A}})$, for all $a, b \in A$, that is, the logic of \mathcal{A} is strongly/weakly \diamond -conjunctive. Then, \mathcal{A} is said to be (strongly)/weakly {classically} \diamond -disjunctive, whenever $\langle \mathfrak{A}, A \setminus D^{\mathcal{A}} \rangle$ is strongly/weakly \diamond -conjunctive, “in which case”/“that is,” the logic of \mathcal{A} is strongly/weakly \diamond -disjunctive, and so is the logic of any class of strongly/weakly \diamond -disjunctive Σ -matrices. Likewise, \mathcal{A} is said to be (weakly/strongly){classically} \sqsupset -implicative, whenever $((a \in D^{\mathcal{A}}) \Rightarrow (b \in D^{\mathcal{A}})) \Leftrightarrow ((a \sqsupset^{\mathfrak{A}} b) \in D^{\mathcal{A}})$, for all $a, b \in A$, in which case it is \sqsupset_{\sqsupset} -disjunctive, while the logic of \mathcal{A} is \sqsupset -implicative, for both (2.8) and (2.10) are true in any \sqsupset -implicative (and so \sqsupset_{\sqsupset} -disjunctive) Σ -matrix, while DT is immediate, and so is the logic of any class of \sqsupset -implicative Σ -matrices. Furthermore, given any $\Sigma' \subseteq \Sigma$, \mathcal{A} is said to be a (Σ') -expansion of its Σ' -reduct $(\mathcal{A} \upharpoonright \Sigma') \triangleq \langle \mathfrak{A} \upharpoonright \Sigma', D^{\mathcal{A}} \rangle$, clearly defining the Σ' -fragment of the logic of \mathcal{A} . Finally, \mathcal{A} is said to be weakly/(strongly) {classically} \wr -negative, provided, for all $a \in A$, $(a \in D^{\mathcal{A}}) \Leftarrow / \Leftrightarrow (\wr^{\mathfrak{A}} a \notin D^{\mathcal{A}})$, in which case it is truth-non-empty / “and so consistent”.

Remark 2.7. For any Σ -matrices \mathcal{A} and \mathfrak{B} , the following hold:

- (i) \mathcal{A} is:

- (a) [weakly] \diamond -disjunctive/-conjunctive iff it is [weakly] \diamond^{\wr} -conjunctive/-disjunctive, respectively, whenever it is \wr -negative, where $(x_0 \diamond^{\wr} x_1) \triangleq \wr(\wr x_0 \diamond x_1)$ is the \wr -dual|De-Morgan counterpart of \diamond ;
- (b) $\sqsupset_{\diamond}^{\wr}$ -implicative, whenever it is both \wr -negative and \diamond -disjunctive, where $(x_0 \sqsupset_{\diamond}^{\wr} x_1) \triangleq (\wr x_0 \diamond x_1)$ is the material implication of/“defined|given by” {negation} \wr and {disjunction} \diamond .
- (c) not \wr -paraconsistent, whenever it is \wr -negative, as well as not (\diamond, \wr) -paracomplete, whenever it is weakly both \wr -negative and \diamond -disjunctive;

- (ii) for any strict [surjective] (injective) $h \in \text{hom}(\mathcal{A}, \mathfrak{B})$, the following hold:

- (a) \mathcal{A} is {weakly} \wr -negative| \diamond -conjunctive/-disjunctive/-implicative iff [f] \mathfrak{B} is so;
- (b) \mathfrak{B} is consistent/truth-non-empty iff [f] \mathcal{A} is so;
- (c) \mathcal{A} is false-/truth-singular (if) ([and]) [only if] \mathfrak{B} is so. □

Given a set I and an I -tuple $\bar{\mathcal{A}}$ of Σ -matrices, [any submatrix \mathcal{B} of] the Σ -matrix $(\prod_{i \in I} \mathcal{A}_i) \triangleq \langle \prod_{i \in I} \mathfrak{A}_i, \prod_{i \in I} D^{\mathcal{A}_i} \rangle$ is called the [a] *[sub]direct product of $\bar{\mathcal{A}}$* [whenever, for each $i \in I$, $\pi_i[B] = A_i$]. As usual, if $(\text{img } \bar{\mathcal{A}}) \subseteq \{\mathcal{A}\}$, where \mathcal{A} is a Σ -matrix, we set $\mathcal{A}^I \triangleq (\prod_{i \in I} \mathcal{A}_i)$.

Given a class \mathbf{M} of Σ -matrices, the class of all “strictly surjectively homomorphic [counter-]images” / “isomorphic copies” / “(consistent) submatrices” of elements of \mathbf{M} is denoted by $(\mathbf{H}^{[-1]} / \mathbf{I} / \mathbf{S}_{(*)})(\mathbf{M})$, respectively. Likewise, the class of all [sub]direct products of tuples (of cardinality $\in K \subseteq \infty$) constituted by elements of \mathbf{M} is denoted by $\mathbf{P}_{(K)}^{[\text{SD}]}(\mathbf{M})$.

2.3.2.1. Classical matrices and logics. Σ -matrices with diagonal characteristic function (and so relation) are said to be *classically-canonical*, isomorphisms between them being diagonal, in which case isomorphic ones being equal. Then, the characteristic function of any Σ -matrix \mathcal{A} with diagonal characteristic relation — viz., injective characteristic function — (and so no-more-than-two-valued) is an isomorphism from it onto the classically-canonical Σ -matrix $\mathfrak{C}(\mathcal{A}) \triangleq \langle \chi^{\mathcal{A}}[\mathfrak{A}], \{1\} \rangle$, called the *[classical] canonization of \mathcal{A}* .

A (classically-canonical) two-valued Σ -matrix \mathcal{A} [with functionally complete underlying algebra] is said to be *[genuinely] (canonical{ly}) \wr -classical*, whenever it is \wr -negative, in which case it is both false- and truth-singular (and so its characteristic relation is diagonal) but is not \wr -paraconsistent, by Remark 2.7(i)(c).

A Σ -logic is said to be *(genuinely) \wr -[sub]classical*, whenever it is [a sublogic of] the logic of a (genuinely) \wr -classical Σ -matrix, in which case it is inferentially consistent. Then, a Σ -matrix is said to be *\wr -classically-defining*, whenever its logic is \wr -classical. Likewise, a unary $\sim \in \Sigma$ is called a *subclassical negation* for a Σ -logic C , whenever the \sim -fragment of C is \sim -subclassical, in which case:

$$(2.16) \quad \sim^m x_0 \notin C(\sim^n x_0),$$

for all $m, n \in \omega$ such that the integer $m - n$ is odd, where the secondary unary connective \wr^l of Σ is defined by induction on $l \in \omega$ via setting $\wr^{0[l+1]} x_0 \triangleq [\wr^l] x_0$.

3. PRELIMINARY KEY ADVANCED GENERIC ISSUES

3.1. **Equality determinants versus matrix hereditary simplicity.** Following the paradigm of the works [14] and [15], an *equality determinant* for a class of Σ -matrices \mathbf{M} is any infinitary quantifier-free equality-free formula Φ of the first-order signature $L \triangleq (\Sigma \cup \{D\})$ (that is, any equality-free formula of the infinitary language $L_{\infty,0}$) with variables in Var_2 such that the infinitary universal sentence $\forall x_0 \forall x_1 (\Phi \leftrightarrow (x_0 \approx x_1))$ with equality is true in \mathbf{M} , in which case Φ is an equality determinant for $\mathbf{I}(\mathbf{S}(\mathbf{M}))$ (cf. Lemma 3.3 of [18] for the “unitary” case discussed in Subsubsection 3.1.1). Then, a *canonical equality determinant* for \mathbf{M} is any Σ -calculus ε of rank 2 such that $\bigwedge \varepsilon$ is an equality determinant for \mathbf{M} . The main distinctive feature of Σ -matrices with equality determinant is as follows:

Lemma 3.1 (cf. Lemma 3.2 of [18] for the “unitary” case). *Any Σ -matrix \mathcal{A} with equality determinant Φ is simple, and so hereditarily so.*

Proof. Then, for any $\bar{a} \in \theta \in \text{Con}(\mathcal{A})$, and all $\varphi \in \text{Fm}_{\Sigma}^2$, we have $\varphi^{\mathfrak{A}}(a_0, a_0) \theta \varphi^{\mathfrak{A}}(a_0, a_1)$, in which case we get $(\varphi^{\mathfrak{A}}(a_0, a_0) \in D^{\mathcal{A}}) \Leftrightarrow (\varphi^{\mathfrak{A}}(a_0, a_1) \in D^{\mathcal{A}})$, and so $\mathcal{A} \models \Phi[x_i/a_i]_{i \in 2}$, for $\mathcal{A} \models \Phi[x_i/a_0]_{i \in 2}$, as $a_0 = a_0$ (in particular, $a_0 = a_1$, in which case $\theta = \Delta_A$, and so \mathcal{A} is simple). \square

Conversely, we have:

Theorem 3.2. *A $\langle n$ implicative Σ -matrix \mathcal{A} is [finitely] hereditarily simple iff it has a ($\{\text{finitary/unary}/\langle\text{axiomatic}\rangle\}$ canonical) equality determinant, in which case this is so for $\mathbf{IS}(\{\langle\mathbf{P}\rangle\})\mathcal{A}$, and so all elements of this class are hereditarily simple.*

Proof. The “if” part is by Lemma 3.1. Conversely, assume \mathcal{A} is finitely hereditarily simple. Let $\varepsilon \triangleq \{\phi_i \vdash \phi_{1-i} \mid i \in 2, \bar{\phi} \in (\text{Fm}_\Sigma^2)^2, (\phi_0[x_1/x_0]) = (\phi_1[x_1/x_0])\}$. Clearly, $\mathcal{A} \models (\bigwedge \varepsilon)[x_i/a]_{i \in 2}$, for all $a \in A$, because every element of $\varepsilon[x_1/x_0]$ is a first-order tautology of the form $\xi \vdash \xi$, where $\xi \in \text{Fm}_\Sigma^1$. Conversely, consider any $\bar{a} \in (A^2 \setminus \Delta_A)$. Let \mathcal{B} be the submatrix of \mathcal{A} generated by the finite set $\text{img } \bar{a}$. Then, it, being finitely-generated is simple, in which case $\theta \triangleq \text{Cg}^{\mathfrak{B}}(\bar{a}) \ni \bar{a} \notin \Delta_B$ of \mathfrak{B} is non-diagonal, and so $\theta \not\subseteq \theta^{\mathfrak{B}}$. Therefore, by (2.2), $\theta^{\mathfrak{B}} \supseteq \Delta_B$, being transitive and symmetric, does not include $\nabla^{\mathfrak{B}}(\bar{a})$, in which case there are some $j \in 2$, some $n \in \omega$, some $\varphi \in \text{Fm}_\Sigma^{n+1}$ and some $\bar{c} \in B^n$ such that $\langle \varphi^{\mathfrak{B}}[x_n/a_j; x_k/c_k]_{k \in n}, \varphi^{\mathfrak{B}}[x_n/a_{1-j}; x_k/c_k]_{k \in n} \rangle \notin \theta^{\mathfrak{B}}$, and so there is some $i \in 2$ such that $\varphi^{\mathfrak{B}}[x_n/a_i; x_k/c_k]_{k \in n} \in D^{\mathfrak{B}} \not\equiv \varphi^{\mathfrak{B}}[x_n/a_{1-i}; x_k/c_k]_{k \in n}$, while, as \mathfrak{B} is generated by $\text{img } \bar{a}$, for each $k \in n$, there is some $\psi_k \in \text{Fm}_\Sigma^2$ such that $c_k = \psi_k^{\mathfrak{B}}[x_l/a_l]_{l \in 2}$. Then, $\phi_i^{\mathfrak{B}}[x_l/a_l]_{l \in 2} \in D^{\mathfrak{B}} \not\equiv \phi_{1-i}^{\mathfrak{B}}[x_l/a_l]_{l \in 2}$, where, for all $m \in 2$, $\phi_m \triangleq (\varphi[x_n/x_m; x_k/\psi_k]_{k \in n}) \in \text{Fm}_\Sigma^2$. And what is more, $(\phi_0[x_1/x_0]) = (\varphi[x_k/(\psi_k[x_1/x_0])]_{k \in n}) = (\phi_1[x_1/x_0])$, in which case $(\phi_i \vdash \phi_{1-i}) \in \varepsilon$, and so $\mathcal{B} \not\models (\bigwedge \varepsilon)[x_l/a_l]_{l \in 2}$. Hence, $\mathcal{A} \not\models (\bigwedge \varepsilon)[x_l/a_l]_{l \in 2}$, for $\bigwedge \varepsilon$ is quantifier-free, and so ε is a unary (in particular, finitary) canonical equality determinant for \mathcal{A} . (Then, $\epsilon \triangleq \{\phi \sqsupset \psi \mid (\phi \vdash \psi) \in \varepsilon\}$ is an axiomatic canonical equality determinant for \mathcal{A} .) On the other hand, any $\Xi \subseteq \text{Fm}_\Sigma^2$ is an axiomatic canonical equality determinant for a Σ -matrix \mathcal{D} iff the universal infinitary strict Horn sentences with equality $\forall x_0 \forall x_1 ((\bigwedge \Xi) \rightarrow (x_0 \approx x_1))$ and $\forall x_0 (\xi[x_1/x_0])$, where $\xi \in \Xi$, of the first-order signature $\Sigma \cup \{D\}$ are true in \mathcal{D} . In this way, the well-known fact that model classes of universal infinitary (strict Horn) theories with equality are closed under \mathbf{I} and \mathbf{S} (as well as \mathbf{P}) — cf., e.g., [8] — completes the argument. \square

3.1.1. Unitary equality determinants versus matrix non-diagonal partial automorphisms. A [partial] (strict) endomorphism of a Σ -matrix \mathcal{A} is any (strict) homomorphism from [a submatrix of] \mathcal{A} to \mathcal{A} ([injective ones being referred to as *partial automorphisms of \mathcal{A}*]).

A *unitary equality determinant* for a class \mathbf{M} of Σ -matrices is any $\Upsilon \subseteq \text{Fm}_\Sigma^1$ such that $\varepsilon_\Upsilon \triangleq \{(v[x_0/x_i]) \vdash (v[x_0/x_{1-i}]) \mid i \in 2, v \in \Upsilon\}$ is a (unary) canonical equality determinant for \mathbf{M} . It is unitary equality determinants that are equality determinants in the sense of [14].

Theorem 3.3. *A Σ -matrix \mathcal{A} has a unitary equality determinant iff it is (finitely) hereditarily simple and has no non-diagonal [injective] partial strict endomorphism.*

Proof. First, let Υ be a unitary equality determinant for \mathcal{A} , \mathcal{B} a submatrix of \mathcal{A} and $h \in \text{hom}(\mathcal{B}, \mathcal{A})$ strict. Then, for every $b \in B$ and each $v \in \Upsilon$, we have $(v^{\mathfrak{A}}(b) = v^{\mathfrak{B}}(b) \in D^{\mathcal{A}}) \Leftrightarrow (v^{\mathfrak{B}}(b) \in D^{\mathcal{B}}) \Leftrightarrow (v^{\mathfrak{A}}(h(b)) = h(v^{\mathfrak{B}}(b)) \in D^{\mathcal{A}})$, in which case we get $h(b) = b$, and so h is diagonal. Thus, the “only if” part is by Lemma 3.1. Conversely, assume \mathcal{A} is finitely hereditarily simple and has no non-diagonal partial automorphism. Consider any $\bar{a} \in A^2$ such that

$$(3.1) \quad (\varphi^{\mathfrak{A}}(a_0) \in D^{\mathcal{A}}) \Leftrightarrow (\varphi^{\mathfrak{A}}(a_1) \in D^{\mathcal{A}}),$$

for all $\varphi \in \text{Fm}_\Sigma^1$. Let f be the carrier of the subalgebra of \mathfrak{A}^2 generated by $\{\bar{a}\}$, and, for each $i \in 2$, \mathcal{B}_i the submatrix of \mathcal{A} generated by $\{a_i\}$, in which case \mathcal{B}_i , being finitely-generated, is simple, while $B_i = \pi_i[f]$, for $\pi_i(\bar{a}) = a_i$ and $\pi_i \in \text{hom}(\mathfrak{A}^2, \mathfrak{A})$. Consider any $i \in 2$ and any $\bar{b}, \bar{c} \in f$ such that $b_i \neq c_i$, in which case there are some $\phi, \psi \in \text{Fm}_\Sigma^1$ such that $\bar{b} = \phi^{\mathfrak{A}^2}(\bar{a})$ and $\bar{c} = \psi^{\mathfrak{A}^2}(\bar{a})$. And what is more,

$\theta \triangleq \text{Cg}^{\mathfrak{B}_i}(\langle b_i, c_i \rangle) \ni \langle b_i, c_i \rangle \notin \Delta_{B_i}$, being a non-diagonal congruence of \mathfrak{B}_i , is not a congruence of \mathcal{B}_i , for this is simple, in which case $B_i^2 \supseteq \theta \not\subseteq \theta^{\mathcal{B}_i} = (\theta^{\mathcal{A}} \cap B_i^2)$, and so, by (2.2), $\nabla^{\mathfrak{B}_i}(b_i, c_i) \not\subseteq \theta^{\mathcal{A}}$, for $\theta^{\mathcal{B}_i}$ is both symmetric and transitive. Hence, there are some $n \in \omega$, some $\xi \in \text{Fm}_{\Sigma}^{n+1}$ and some $\bar{d} \in B_i^n$ such that $(\xi^{\mathfrak{A}}[x_j/d_j; x_n/b_i]_{j \in n} \in D^{\mathcal{A}}) \Leftrightarrow (\xi^{\mathfrak{A}}[x_j/d_j; x_n/c_i]_{j \in n} \notin D^{\mathcal{A}})$. Then, for each $j \in n$, there is some $v_j \in \text{Fm}_{\Sigma}^1$ such that $d_j = v_j^{\mathfrak{A}}(a_i)$. Let $(\eta|\zeta) = (\xi[x_j/v_j; x_n/(\phi|\psi)]_{j \in n} \in \text{Fm}_{\Sigma}^1)$, in which case, for each $k \in 2$, $(\eta|\zeta)^{\mathfrak{A}}(a_k) = \xi^{\mathfrak{A}}[x_j/v_j^{\mathfrak{A}}(a_k); x_n/(b|c)_k]_{j \in n}$, and so, by (3.1), $(\xi^{\mathfrak{A}}[x_j/v_j^{\mathfrak{A}}(a_{1-i}); x_n/b_{1-i}]_{j \in n} \in D^{\mathcal{A}}) \Leftrightarrow (\xi^{\mathfrak{A}}[x_j/v_j^{\mathfrak{A}}(a_{1-i}); x_n/c_{1-i}]_{j \in n} \notin D^{\mathcal{A}})$ (in particular, $b_{1-i} \neq c_{1-i}$). In this way, f is a bijection from B_0 onto B_1 , in which case it, being a subalgebra of \mathfrak{A}^2 , is an embedding of \mathfrak{B}_0 into \mathfrak{A} , and so, by (3.1), is that of \mathcal{B}_0 into \mathcal{A} (in particular, a partial automorphism of \mathcal{A}). Thus, f is diagonal, in which case $a_1 = f(a_0) = a_0$, so Fm_{Σ}^1 is a unitary equality determinant for \mathcal{A} . \square

Clearly, any consistent truth-non-empty two-valued (in particular, classical) Σ -matrix \mathcal{A} is both false- and truth-singular, in which case its characteristic relation is diagonal, and so $\{x_0\}$ is an equality determinant for \mathcal{A} .

3.2. Disjunctivity.

3.2.1. Disjunctivity versus multiplicativity. A Σ -logic C is said to be \vee -(singularly-)multiplicative, provided, for all $X \subseteq \text{Fm}_{\Sigma}^{\omega}$ and all $\phi, \psi \in \text{Fm}_{\Sigma}^{\omega}$, it holds that $(\vee[C(X \cup \{\phi\}) \times \{\psi\}]) \subseteq C(X \cup \{\phi \vee \psi\})$.

Lemma 3.4. *Any Σ -logic C is \vee -disjunctive iff it is both weakly \vee -disjunctive and \vee -multiplicative as well as satisfies both (2.4) and (2.5).*

Proof. The ‘‘only if’’ part is immediate. Conversely, assume C is both weakly \vee -disjunctive and \vee -multiplicative as well as satisfies both (2.4) and (2.5). Consider any $X \subseteq \text{Fm}_{\Sigma}^{\omega}$, any $\phi, \psi \in \text{Fm}_{\Sigma}^{\omega}$ and any $\varphi \in (C(X \cup \{\phi\}) \cap C(X \cup \{\psi\}))$. Then, by the \vee -multiplicativity of C and (2.4), we have $(\psi \vee \varphi) \in C(\varphi \vee \psi) \subseteq C(X \cup \{\phi \vee \psi\})$. Likewise, by the \vee -multiplicativity of C and (2.5), we have $\varphi \in C(\varphi \vee \psi) \subseteq C(X \cup \{\psi \vee \varphi\})$. In this way, we eventually get $\varphi \in C(X \cup \{\phi \vee \psi\})$. \square

3.2.1.1. Implicativity versus intrinsic disjunctivity.

Theorem 3.5. *Let C be a weakly \sqsupset -implicative Σ -logic and $\vee \triangleq \wp \sqsupset$. Then, the following hold:*

- (i) C is both weakly \vee -disjunctive and \vee -multiplicative;
- (ii) C is \sqsupset -implicative iff it is \vee -disjunctive iff it satisfies (2.4).

Proof. (i) First, (2.3) with $i = 0$ is by DT and (2.8). Likewise, (2.3) with $i = 1$ is by (2.7) and (2.8). Now, consider any $X \subseteq \text{Fm}_{\Sigma}^{\omega}$ and any $\phi, \psi, \varphi \in \text{Fm}_{\Sigma}^{\omega}$. Then, by DT and (2.8), we have $((\psi \in C(X \cup \{\phi\}) \Rightarrow ((\phi \sqsupset \varphi) \in C(X \cup \{\psi \sqsupset \varphi\})))$, applying which twice, the second time being with $(\psi \sqsupset \varphi) | (\phi \sqsupset \varphi)$ instead of $\phi | \psi$, respectively, we conclude that C is \vee -multiplicative.

- (ii) Assume C is \sqsupset -implicative. Then, $((x_0 \vee x_0) \sqsupset x_0) = ((2.10)[x_1/x_0])$ is satisfied in C , for this is structural, and so is (2.5), in view of (2.8). Furthermore, by (2.8), we have $x_0 \in C(\{x_0 \vee x_1, x_0 \sqsupset x_1, x_1 \sqsupset x_0\})$, in which case, by DT, we get $((x_0 \sqsupset x_1) \sqsupset x_0) \in C(\{x_0 \vee x_1, x_1 \sqsupset x_0\})$, and so, by (2.8) and (2.10), we eventually get $x_0 \in C(\{x_0 \vee x_1, x_1 \sqsupset x_0\})$ (in particular, by DT, (2.4) is satisfied in C). Then, Lemma 3.4, (i) and (2.9) complete the argument. \square

3.2.2. Disjunctive consistent finitely-generated models of finitely-valued weakly disjunctive logics.

Lemma 3.6. $\mathbf{H}(\mathbf{H}^{-1}(\mathbf{M})) \subseteq \mathbf{H}^{-1}(\mathbf{H}(\mathbf{M}))$, for any class of Σ -matrices \mathbf{M} .

Proof. Let \mathcal{A} and \mathcal{B} be Σ -matrices, $\mathcal{C} \in \mathbf{M}$ and $(h|g) \in \text{hom}_{\mathbb{S}}^{\mathbb{S}}(\mathcal{B}, \mathcal{C}|\mathcal{A})$. Then, by Remark 2.5(i), $(\ker(h|g)) \in \text{Con}(\mathcal{B})$, in which case $(\ker(h|g)) \subseteq \theta \triangleq \vartheta(\mathcal{B}) \in \text{Con}(\mathcal{B})$, and so, by the Homomorphism Theorem, $(\nu_{\theta} \circ (h|g)^{-1}) \in \text{hom}_{\mathbb{S}}^{\mathbb{S}}(\mathcal{C}|\mathcal{A}, \mathcal{B}/\theta)$. \square

Lemma 3.7 (Finite Subdirect Product Lemma; cf. Lemma 2.7 of [18]). *Let \mathbf{M} be a finite class of finite Σ -matrices and \mathcal{A} a [non-]simple finite(ly-generated) model of the logic of \mathbf{M} . Then, $(\mathcal{A}[\vartheta(\mathcal{A})]) \in \mathbf{HP}_{\omega}^{\mathbb{S}\mathbb{D}} \mathbf{S}_* \mathbf{M}$.*

Lemma 3.8. *Let \mathbf{M} be a class of weakly \vee -disjunctive Σ -matrices, I a finite set, $\overline{\mathcal{C}} \in \mathbf{M}^I$, and \mathcal{D} a consistent \vee -disjunctive submatrix of $\prod \overline{\mathcal{C}}$. Then, there is some $i \in I$ such that $(\pi_i \upharpoonright \mathcal{D}) \in \text{hom}_{\mathbb{S}}^{\mathbb{S}}(\mathcal{D}, \mathcal{C}_i)$.*

Proof. By contradiction. For suppose that, for every $i \in I$, $(\pi_i \upharpoonright \mathcal{D}) \notin \text{hom}_{\mathbb{S}}^{\mathbb{S}}(\mathcal{D}, \mathcal{C}_i)$, in which case $D^{\mathcal{D}} \subsetneq (\pi_i \upharpoonright \mathcal{D})^{-1}[D^{\mathcal{C}_i}] = (D \cap \pi_i^{-1}[D^{\mathcal{C}_i}])$, for $(\pi_i \upharpoonright \mathcal{D}) \in \text{hom}(\mathcal{D}, \mathcal{C}_i)$ is surjective, and so there is some $a_i \in (D \setminus D^{\mathcal{D}})$ such that $\pi_i(a_i) \in D^{\mathcal{C}_i}$. By induction on the cardinality of any $J \subseteq I$, let us prove that there is some $b \in (D \setminus D^{\mathcal{D}})$ such that $\pi_j(b) \in D^{\mathcal{C}_j}$, for all $j \in J$, as follows. In case $J = \emptyset$, take any $b \in (D \setminus D^{\mathcal{D}}) \neq \emptyset$, for \mathcal{D} is consistent. Otherwise, take any $j \in J$, in which case $K \triangleq (J \setminus \{j\}) \subseteq I$, while $|K| < |J|$, so, by the induction hypothesis, there is some $c \in (D \setminus D^{\mathcal{D}})$ such that $\pi_k(c) \in D^{\mathcal{C}_k}$, for all $k \in K$. Then, by the \vee -disjunctivity of \mathcal{D} , $b \triangleq (c \vee^{\mathcal{D}} a_j) \in (D \setminus D^{\mathcal{D}})$, while $\pi_i(b) \in D^{\mathcal{C}_i}$, for all $i \in J = (K \cup \{j\})$, because $(\pi_i \upharpoonright \mathcal{D}) \in \text{hom}(\mathcal{D}, \mathcal{C}_i)$, while \mathcal{C}_i is weakly \vee -disjunctive. In particular, when $J = I$, there is some $b \in (D \setminus D^{\mathcal{D}})$ such that $\pi_i(b) \in D^{\mathcal{C}_i}$, for all $i \in I$. This contradicts to the fact that $D^{\mathcal{D}} = (D \cap \bigcap_{i \in I} \pi_i^{-1}[D^{\mathcal{C}_i}])$, as required. \square

By Lemmas 3.6, 3.7, 3.8 and Remark 2.7(ii), we immediately have:

Theorem 3.9. *Let \mathbf{M} be a finite class of finite weakly \vee -disjunctive Σ -matrices, C the logic of \mathbf{M} and \mathcal{A} a finite[ly-generated] consistent \vee -disjunctive model of C . Then, $\mathcal{A} \in \mathbf{H}^{-1}(\mathbf{H}(\mathbf{S}_*(\mathbf{M})))$.*

3.2.3. *Non-paraconsistency versus Resolution.* Given any Σ -logic C , by $C^{\mathbf{R}}$ we denote the extension of C relatively axiomatized by the *Resolution* rule (cf. [20]):

$$(3.2) \quad \{x_0 \vee x_1, \lambda x_0 \vee x_1\} \vdash x_1.$$

Applying Lemma 3.4 and (2.5) to (2.11) twice, we have:

Lemma 3.10. (3.2) *is satisfied in any \vee -disjunctive non- λ -paraconsistent Σ -logic.*

Theorem 3.11. *Let \mathbf{M} be a finite class of finite \vee -disjunctive Σ -matrices and C the logic of \mathbf{M} . Then, $C^{\mathbf{R}}$ is defined by the class \mathbf{S} of all non- λ -paraconsistent elements of $\mathbf{S}_*(\mathbf{M})$, and so is \vee -disjunctive.*

Proof. Then, C is \vee -disjunctive, while the logic of \mathbf{S} is a both finitary, \vee -disjunctive (in view of Remark 2.7(ii)(a)) and non- λ -paraconsistent extension of C , and so an extension of $C^{\mathbf{R}}$, in view of Lemma 3.10. Conversely, consider any $n \in (\omega \setminus 1)$, any $\Gamma \subseteq \text{Fm}_{\Sigma}^n$ and any $\varphi \in (\text{Fm}_{\Sigma}^n \setminus C^{\mathbf{R}}(\Gamma))$, in which case, by (2.14) with $\alpha = n$, $\varphi \notin C(\Gamma) = \text{Cn}_{\mathbf{M}}^{\omega}(\Gamma) \supseteq \text{Cn}_{\mathbf{M}}^n(\Gamma)$, and so $\mathcal{T} \triangleq \{T \in \mathcal{B}_{\mathbf{M}}^n \mid \Gamma \subseteq T \not\vdash \varphi\} \neq \emptyset$. Then, since n as well as both \mathbf{M} and all elements of it are finite, the class $\{(\mathcal{A}, h) \mid \mathcal{A} \in \mathbf{M}, h \in \text{hom}(\mathfrak{Fm}_{\Sigma}^n, \mathfrak{A})\}$ is finite, in which case the set $\mathcal{B}_{\mathbf{M}}^n$ is finite, and so is $\mathcal{T} \subseteq \mathcal{B}_{\mathbf{M}}^n$. Let $m \triangleq |\mathcal{T}| \in (\omega \setminus 1)$ and \overline{T} any bijection from m onto \mathcal{T} , in which case, for each $i \in m$, there is some $\mathcal{A}_i \in \mathbf{M}$ and some $h_i \in \text{hom}(\mathfrak{Fm}_{\Sigma}^n, \mathfrak{A}_i)$ such that $\Gamma \subseteq T_i = h_i^{-1}[D^{\mathcal{A}_i}] \not\vdash \varphi$, and so $B_i \triangleq (\text{img } h_i)$ forms a subalgebra of \mathfrak{A}_i , while $\mathcal{B}_i \triangleq (\mathcal{A}_i \upharpoonright B_i) \in \mathbf{S}(\mathbf{M})$, whereas $h_i^{-1}[D^{\mathcal{B}_i}] = T_i$ (in particular, \mathcal{B}_i is consistent, for $h_i(\varphi) \in (B_i \setminus D^{\mathcal{B}_i})$), as well as $h_i \in \text{hom}(\mathfrak{Fm}_{\Sigma}^n, \mathfrak{B}_i)$ (In particular, $T_i \in \mathcal{B}_{\mathbf{S}_*(\mathbf{M})}^n$). We prove, by contradiction, that, for some $i \in m$, \mathcal{B}_i is not λ -paraconsistent. For suppose each \mathcal{B}_i , where $i \in m$, is λ -paraconsistent. By induction on any $j \in (m+1)$,

we set $\Xi_j \triangleq (\{\varphi\} \mid \{\lambda^k \psi \vee \phi \mid k \in 2, \psi \in T_{j-1} \ni \lambda \psi, \phi \in \Xi_{j-1}\}) \subseteq \text{Fm}_\Sigma^n$, whenever $j = | \neq 0$, respectively, and prove that

$$(3.3) \quad \varphi \in C^{\text{R}}(\Xi_j),$$

$$(3.4) \quad \Xi_j \subseteq (C(T_i) \cap C(\Xi_i)),$$

for all $i \in j$. The case, when $j = 0 = \emptyset$, is evident. Otherwise, $(j-1) \in (m \cap j)$, in which case \mathcal{B}_{j-1} is \sim -paraconsistent, and so there is some $\psi \in T_{j-1}$ such that $\lambda \psi \in T_{j-1}$. In particular, for each $\phi \in \Xi_{j-1}$ and every $k \in 2$, $(\lambda^k \psi \vee \phi) \in \Xi_j$, in which case, by (3.2)[$x_0/\psi, x_1/\phi$] and the structurality of C^{R} , $\phi \in C^{\text{R}}(\Xi_j)$, and so, by the induction hypothesis, $\varphi \in C^{\text{R}}(\Xi_{j-1}) \subseteq C^{\text{R}}(\Xi_j)$. Thus, (3.3) holds. Likewise, by the \vee -disjunctivity of C , for each $\phi \in \Xi_{j-1}$, every $k \in 2$ and all $\psi \in T_{j-1}$ such that $\lambda \psi \in T_{j-1}$, we have $(\lambda^k \psi \vee \phi) \in (C(\Xi_{j-1}) \cap C(T_{j-1}))$ (in particular, (3.4) with $i = (j-1)$ holds), and so, by the induction hypothesis as well as (3.4) with $i = (j-1)$, we get (3.4), for all $i \in (j-1)$. Thus, (3.4) holds, for all $i \in (\{j-1\} \cup (j-1)) = j$. In this way, by (3.3) with $j = m$, we have $\Xi_m \not\subseteq C^{\text{R}}(\Gamma) \supseteq C(\Gamma) = \text{Cn}_M^\omega(\Gamma)$, in which case, by (2.14) with $\alpha = n$, we get $\Xi_m \not\subseteq \text{Cn}_M^n(\Gamma)$, and so there is some $T \in \mathcal{B}_M^n$ such that $\Gamma \subseteq T \not\subseteq \Xi_m$. In that case, if T contained φ , that is, included Ξ_0 , then, by (3.4) with $j = m$ and $i = 0 \in m$, for $m \neq 0$, we would have $\Xi_m \subseteq C(T)$, and so, by (2.14) with $\alpha = n$, would get $\Xi_m \subseteq \text{Cn}_M^n(T) = T$. Therefore, $\varphi \notin T$, in which case $T \in \mathcal{T}$, and so $T = T_l$, for some $l \in m$. Hence, by (3.4) with $j = m$ and $i = l$, we have $\Xi_m \subseteq C(T)$, in which case, by (2.14) with $\alpha = n$, we get $\Xi_m \subseteq \text{Cn}_M^n(T) = T$, and so this contradiction shows that there is some $i \in m$, such that \mathcal{B}_i is not λ -paraconsistent. In this way, $\mathcal{B}_i \in \mathcal{S}$, in which case $\varphi \notin \text{Cn}_{\mathcal{B}_i}^n(\Gamma) \supseteq \text{Cn}_\Sigma^n(\Gamma)$, and so, by (2.14) with $\alpha = n$, $\varphi \notin \text{Cn}_\Sigma^\omega(\Gamma)$, as required, for $\wp_\omega(\text{Fm}_\Sigma^\omega) \subseteq \bigcup_{n \in (\omega \setminus 1)} \wp(\text{Fm}_\Sigma^n)$. \square

3.3. Some peculiarities of false-singular matrices.

3.3.1. Subdirect products of consistent submatrices of weakly conjunctive matrices.

Lemma 3.12. *Let \mathcal{A} be a false-singular weakly \diamond -conjunctive Σ -matrix, $f \in (A \setminus D^{\mathcal{A}})$, I a finite set, $\bar{\mathcal{B}} \in \mathbf{S}_*(\mathcal{A})^I$ and \mathcal{D} a subdirect product of it. Then, $(I \times \{f\}) \in \mathcal{D}$.*

Proof. By induction on the cardinality of any $J \subseteq I$, let us prove that there is some $a \in D$ including $(J \times \{f\})$. First, when $J = \emptyset$, take any $a \in D \neq \emptyset$, in which case $(J \times \{f\}) = \emptyset \subseteq a$. Now, assume $J \neq \emptyset$. Take any $j \in J \subseteq I$, in which case $K \triangleq (J \setminus \{j\}) \subseteq I$, while $|K| < |J|$, and so, as \mathcal{B}_j is a consistent submatrix of the false-singular Σ -matrix \mathcal{A} , we have $f \in B_j = \pi_j[D]$. Hence, there is some $b \in D$ such that $\pi_j(b) = f$, while, by induction hypothesis, there is some $c \in D$ including $(K \times \{f\})$. Therefore, since $J = (K \cup \{j\})$, while \mathcal{A} is both weakly \diamond -conjunctive and false-singular, we have $D \ni a \triangleq (c \diamond^{\mathcal{D}} b) \supseteq (J \times \{f\})$. Thus, when $J = I$, we eventually get $D \ni (I \times \{f\})$, as required. \square

3.3.2. Models of weakly implicative logics.

Lemma 3.13. *Let \mathcal{A} be a false-singular Σ -matrix. Suppose (2.6), (2.7) and (2.8) are true in \mathcal{A} . Then, \mathcal{A} is \sqsupset -implicative. In particular, any false-singular Σ -matrix is \sqsupset -implicative iff its logic is [weakly] so.*

Proof. Then, for all $a, b \in (A \setminus D^{\mathcal{A}})$, we have $a = b$, in which case, by (2.6), we get $(a \sqsupset^{\mathcal{A}} b) = (a \sqsupset^{\mathcal{A}} a) \in D^{\mathcal{A}}$, and so (2.7) and (2.8) complete the argument. \square

3.4. Logic versus model congruences.

Lemma 3.14. *Let C be a Σ -logic, $\theta \in \text{Con}(C)$, $\mathcal{A} \in \text{Mod}(C)$ and $h \in \text{hom}(\mathfrak{Fm}_\Sigma^\omega, \mathfrak{A})$. Then, $h[\theta] \subseteq \mathfrak{D}(\mathcal{A})$.*

Proof. Then, $\vartheta \triangleq (\bigcup\{g[\theta] \mid g \in \text{hom}(\mathfrak{Fm}_\Sigma^\omega, \mathfrak{A})\})$ is symmetric, for θ is so. And what is more, since $\theta \subseteq \equiv_C^\omega$, while $\mathcal{A} \in \text{Mod}(C)$, $\vartheta \subseteq \theta^{\mathcal{A}}$. Next, consider any $a \in A$. Let $g \triangleq [x_k/a]_{k \in \omega} \in \text{hom}(\mathfrak{Fm}_\Sigma^\omega, \mathfrak{A})$. Then, since $\langle x_0, x_0 \rangle \in \theta$, $\langle a, a \rangle = g(\langle x_0, x_0 \rangle) \in g[\theta] \subseteq \vartheta$, and so $\Delta_{\mathcal{A}} \subseteq \vartheta$. Now, consider any $\varsigma \in \Sigma$ of arity $n \in \omega$, any $i \in n$, any $\langle a, b \rangle \in \vartheta$ and any $\bar{c} \in A^{n-1}$. Then, there are some $\langle \phi, \psi \rangle \in \theta$ and some $f \in \text{hom}(\mathfrak{Fm}_\Sigma^\omega, \mathfrak{A})$ such that $a = f(\phi)$ and $b = f(\psi)$. Let $V \triangleq (\text{Var}(\phi) \cup \text{Var}(\psi) \cup \{x_i\}) \in \wp_\omega(\text{Var}_\omega)$, in which case $|\text{Var}_\omega \setminus V| = \omega \geq (n-1)$, for $|\text{Var}_\omega| = \omega$ is infinite, and so there is some injective $\bar{v} \in (\text{Var}_\omega \setminus V)^{n-1}$. Let $\varphi \triangleq (\varsigma(\bar{x}_n)[x_j/v_j; x_k/v_{k-1}]_{j \in i; k \in (n \setminus (i+1))}) \in \text{Fm}_\Sigma^\omega$ and $g \in \text{hom}(\mathfrak{Fm}_\Sigma^\omega, \mathfrak{A})$ extend $(f \upharpoonright (\text{Var}_\omega \setminus (\text{img } \bar{v}))) \cup (\bar{c} \circ \bar{v}^{-1})$, in which case $\langle \varphi[x_i/\phi], \varphi[x_i/\psi] \rangle \in \theta$, so $\langle \varphi^{\mathfrak{A}}[x_i/a; v_l/c_l]_{l \in (n-1)}, \varphi^{\mathfrak{A}}[x_i/b; v_l/c_l]_{l \in (n-1)} \rangle = g(\langle \varphi[x_i/\phi], \varphi[x_i/\psi] \rangle) \in g[\theta] \subseteq \vartheta$. Thus, unary algebraic operations of \mathfrak{A} are ϑ -monotonic. Therefore, $\eta \triangleq \text{Tr}(\vartheta)$ is a congruence of \mathfrak{A} . And what is more, $\theta^{\mathcal{A}} \supseteq \vartheta$, being transitive, includes η , in which case $\eta \in \text{Con}(\mathcal{A})$, and so $h[\theta] \subseteq \vartheta \subseteq \eta \subseteq \mathfrak{D}(\mathcal{A})$. \square

3.4.1. *Simple models versus intrinsic varieties.* As a particular case of Lemma 3.14, we first have (from now on, we follow Definition 2.2 tacitly):

Corollary 3.15. *Let C be a Σ -logic. Then, $\pi_0[\text{Mod}^*(C)] \subseteq \text{IV}(C)$.*

Corollary 3.16. *Let C be a Σ -logic. Then, $\mathfrak{D}(C)$ is fully-invariant. In particular, $\mathfrak{D}(C) = \theta_{\text{IV}(C)}^\omega$.*

Proof. Consider any $\sigma \in \text{hom}(\mathfrak{Fm}_\Sigma^\omega, \mathfrak{Fm}_\Sigma^\omega)$ and any $T \in (\text{img } C)$, in which case, by the structurality of C , $\mathcal{A}_T \triangleq \langle \mathfrak{Fm}_\Sigma^\omega, T \rangle \in \text{Mod}(C)$, so, by Lemma 3.14, $\sigma[\mathfrak{D}(C)] \subseteq \mathfrak{D}(\mathcal{A}_T)$. Then, $\sigma[\mathfrak{D}(C)] \subseteq \theta \triangleq (\text{Eq}_\Sigma^\omega \cap \bigcap\{\mathfrak{D}(\mathcal{A}_T) \mid T \in (\text{img } C)\}) \subseteq (\text{Eq}_\Sigma^\omega \cap \bigcap\{\theta^{\mathcal{A}_T} \mid T \in (\text{img } C)\}) = \equiv_C^\omega$. Moreover, for each $T \in (\text{img } C)$, $\mathfrak{D}(\mathcal{A}_T) \in \text{Con}(\mathfrak{Fm}_\Sigma^\omega)$, in which case $\theta \in \text{Con}(\mathfrak{Fm}_\Sigma^\omega)$, and so $\sigma[\mathfrak{D}(C)] \subseteq \theta \subseteq \mathfrak{D}(C)$. \square

Lemma 3.17. *Let \mathbf{M} be a class of Σ -matrices, $\mathbf{K} \triangleq \pi_0[\mathbf{M}]$ and C the logic of \mathbf{M} . Then, $\theta_{\mathbf{K}}^\omega \subseteq \equiv_C^\omega$, in which case $\theta_{\mathbf{K}}^\omega \subseteq \mathfrak{D}(C)$, and so $\text{IV}(C) \subseteq \mathbf{V}(\mathbf{K})$.*

Proof. Then, for any $\langle \phi, \psi \rangle \in \theta_{\mathbf{K}}^\omega$, $\mathcal{A} \in \mathbf{M}$ and $h \in \text{hom}(\mathfrak{Fm}_\Sigma^\omega, \mathfrak{A})$, $\mathfrak{A} \in \mathbf{K}$, in which case $\langle h(\phi), h(\psi) \rangle \in \Delta_{\mathcal{A}} \subseteq \theta^{\mathcal{A}}$, and so $\phi \equiv_C^\omega \psi$. \square

By Corollary 3.15 and Lemma 3.17, we then have:

Corollary 3.18. *Let \mathbf{M} be a class of Σ -matrices, $\mathbf{K} \triangleq \pi_0[\mathbf{M}]$ and C the logic of \mathbf{M} . Then, $\pi_0[\text{Mod}^*(C)] \subseteq \mathbf{V}(\mathbf{K})$.*

Theorem 3.19. *Let \mathbf{M} be a class of simple Σ -matrices, $\mathbf{K} \triangleq \pi_0[\mathbf{M}]$ and C the logic of \mathbf{M} . Then, $\text{IV}(C) = \mathbf{V}(\mathbf{K})$.*

4. SELF-EXTENSIONAL LOGICS VERSUS SIMPLE MATRICES

Theorem 4.1. *Let C be a Σ -logic and $\mathbf{V} \triangleq \text{IV}(C)$ (as well as \mathbf{M} a class of simple Σ -matrices, $\mathbf{K} \triangleq \pi_0[\mathbf{M}]$ and $\alpha \triangleq ([1 \cup] (\omega \cap \bigcup\{|A| \mid \mathcal{A} \in \mathbf{M}\}))$). (Suppose C is defined by \mathbf{M} .) Then, (i) \Leftrightarrow (ii) \Leftrightarrow (iii) \Leftrightarrow (iv) \Rightarrow (v) \Rightarrow (vi) \Rightarrow (i), where:*

- (i) C is self-extensional;
- (ii) $\equiv_C^\omega \subseteq \theta_{\mathbf{V}}^\omega$;
- (iii) $\equiv_C^\omega = \theta_{\mathbf{V}}^\omega$;
- (iv) for all distinct $a, b \in F_{\mathbf{V}}^\alpha$, there are some $\mathcal{A} \in \mathbf{M}$ and some $h \in \text{hom}(\mathfrak{Fm}_\Sigma^\omega, \mathfrak{A})$ such that $\chi^{\mathcal{A}}(h(a)) \neq \chi^{\mathcal{A}}(h(b))$;

- (v) *there is some class C of Σ -algebras such that $K \subseteq \mathbf{V}(C)$ and, for each $\mathfrak{A} \in C$ and all distinct $a, b \in A$, there are some $\mathfrak{B} \in M$ and some $h \in \text{hom}(\mathfrak{A}, \mathfrak{B})$ such that $\chi^{\mathfrak{B}}(h(a)) \neq \chi^{\mathfrak{B}}(h(b))$;*
- (vi) *there is some $S \subseteq \text{Mod}(C)$ such that $V \subseteq \mathbf{V}(\pi_0[S])$ and, for each $\mathfrak{A} \in S$, it holds that $(A^2 \cap \bigcap \{\theta^{\mathfrak{B}} \mid \mathfrak{B} \in S, \mathfrak{B} = \mathfrak{A}\}) \subseteq \Delta_A$.*

Proof. In that case, by Corollary 3.16 (and Theorem 3.19), $\mathfrak{D}(C) = \theta_V^\omega$ (as well as $V = \mathbf{V}(K)$, and so $\theta_V^\omega = \theta_K^\omega$). Then, (i) \Leftrightarrow (iii) is immediate, while (ii) is a particular case of (iii), whereas the converse is by the inclusion $\mathfrak{D}(C) \subseteq \equiv_C^\omega$.

(Next, assume (iii) holds. Then, $\theta^{\alpha'} \triangleq \equiv_C^{\alpha'} = \theta_K^{\alpha'} = \theta_V^{\alpha'} \in \text{Con}(\mathfrak{Fm}_\Sigma^{\alpha'})$, for all Σ -ranks α' . Furthermore, consider any distinct $a, b \in F_V^\alpha$. Then, there are some $\phi, \psi \in \text{Fm}_\Sigma^\alpha$ such that $\nu_{\theta^\alpha}(\phi) = a \neq b = \nu_{\theta^\alpha}(\psi)$, in which case, by (2.14), $\text{Cn}_M^\alpha(\phi) \neq \text{Cn}_M^\alpha(\psi)$, and so there are some $\mathfrak{A} \in M$ and some $g \in \text{hom}(\mathfrak{Fm}_\Sigma^\alpha, \mathfrak{A})$ such that $\chi^{\mathfrak{A}}(g(\phi)) \neq \chi^{\mathfrak{A}}(g(\psi))$. In that case, $\theta^\alpha \subseteq (\ker g)$, and so, by the Homomorphism Theorem, $h \triangleq (g \circ \nu_{\theta^\alpha}^{-1}) \in \text{hom}(\mathfrak{F}_V^\alpha, \mathfrak{A})$. Then, $h(a/b) = g(\phi/\psi)$, in which case $\chi^{\mathfrak{A}}(h(a)) \neq \chi^{\mathfrak{A}}(h(b))$, and so (iv) holds.

Now, assume (iv) holds. Consider any $\mathfrak{A} \in K$ and the following cases:

- $|A| \leq \alpha$. Let $h \in \text{hom}(\mathfrak{Fm}_\Sigma^\alpha, \mathfrak{A})$ extend any surjection from Var_α onto A , in which case it is surjective, while $\theta \triangleq \theta_V^\alpha = \theta_K^\alpha \subseteq (\ker h)$, and so, by the Homomorphism Theorem, $g \triangleq (h \circ \nu_\theta^{-1}) \in \text{hom}(\mathfrak{F}_V^\alpha, \mathfrak{A})$ is surjective. Thus, $\mathfrak{A} \in \mathbf{V}(\mathfrak{F}_V^\alpha)$.
- $|A| \not\leq \alpha$. Then, $\alpha = \omega$. Consider any Σ -identity $\phi \approx \psi$ true in \mathfrak{F}_V^ω and any $h \in \text{hom}(\mathfrak{Fm}_\Sigma^\omega, \mathfrak{A})$, in which case, we have $\theta \triangleq \theta_V^\omega = \theta_K^\omega \subseteq (\ker h)$, and so, since $\nu_\theta \in \text{hom}(\mathfrak{Fm}_\Sigma^\omega, \mathfrak{F}_V^\omega)$, we get $\langle \phi, \psi \rangle \in (\ker \nu_\theta) \subseteq (\ker h)$. Thus, $\mathfrak{A} \in \mathbf{V}(\mathfrak{F}_V^\omega)$.

In this way, (v) with $C \triangleq \{\mathfrak{F}_V^\alpha\}$ holds.

Further, assume (v) holds. Let $C' \triangleq \{\mathfrak{A} \in C \mid |A| > 1\}$ and $S \triangleq \{\langle \mathfrak{A}, h^{-1}[D^{\mathfrak{B}}] \mid \mathfrak{A} \in C', \mathfrak{B} \in M, h \in \text{hom}(\mathfrak{A}, \mathfrak{B}) \rangle\}$. Then, for all $\mathfrak{A} \in C'$, each $\mathfrak{B} \in M$ and every $h \in \text{hom}(\mathfrak{A}, \mathfrak{B})$, h is a strict homomorphism from $\mathcal{C} \triangleq \langle \mathfrak{A}, h^{-1}[D^{\mathfrak{B}}] \rangle$ to \mathfrak{B} , in which case, by (2.15), $\mathcal{C} \in \text{Mod}(C)$, and so $S \subseteq \text{Mod}(C)$, while $\chi^{\mathcal{C}} = (h \circ \chi^{\mathfrak{B}})$, whereas $\pi_0[S] = C'$ generates the variety $\mathbf{V}(C)$. In this way, (vi) holds.)

Finally, assume (vi) holds. Consider any $\phi, \psi \in \text{Fm}_\Sigma^\omega$ such that $\phi \equiv_C^\omega \psi$, any $\mathfrak{A} \in S$ and any $h \in \text{hom}(\mathfrak{Fm}_\Sigma^\omega, \mathfrak{A})$. Then, for each $\mathfrak{B} \in S$ with $\mathfrak{B} = \mathfrak{A}$, $h(\phi) \theta^{\mathfrak{B}} h(\psi)$, in which case $h(\phi) = h(\psi)$, so $\mathfrak{A} \models (\phi \approx \psi)$. Thus, $V \subseteq \mathbf{V}(\pi_0[S]) \models (\phi \approx \psi)$, so (ii) holds. \square

When both M and all elements of it are finite, α is finite, in which case \mathfrak{F}_V^α is finite and can be found effectively, and so, taking (2.15) and Remark 2.5(iv) into account, the item (iv) of Theorem 4.1 yields an effective procedure of checking the self-extensionality of any logic defined by a finite class of finite matrices. However, its computational complexity may be too large to count it *practically* applicable. For instance, in the unitary n -valued case, where $n \in (\omega \setminus 1)$, the upper limit n^{n^n} of $|F_V^\alpha|$ as well as the predetermined computational complexity $n^{n^{n^n}}$ of the procedure involved become too large even in the three-/four-valued case. And, though, in the two-valued case, this limit — 16 — as well as the respective complexity — $2^{16} = 65536$ — are reasonably acceptable, this is no longer matter in view of:

Example 4.2. Let \mathcal{A} be a Σ -matrix. Suppose it is both false- and truth-singular (in particular, two-valued as well as both consistent and truth-non-empty [in particular, classical]), in which case $\theta^{\mathcal{A}} = \Delta_A$, for $\chi^{\mathcal{A}}$ is injective, and so \mathcal{A} is simple. Then, by Theorems 3.19 and 4.1(vi) \Rightarrow (i) with $S = \{\mathcal{A}\}$, the logic of \mathcal{A} is self-extensional, its intrinsic variety being generated by \mathfrak{A} . Thus, by the self-extensionality of inferentially inconsistent logics, *any two-valued logic is self-extensional*. \square

Nevertheless, the procedure involved is simplified much under hereditary simplicity as well as either implicativity or both conjunctivity and disjunctivity of finitely many finite defining matrices upon the basis of the item (v) of Theorem 4.1.

4.1. Self-extensionality of conjunctive disjunctive logics versus distributive lattices.

Remark 4.3. Let C be a $\bar{\wedge}$ -conjunctive or/and $\bar{\vee}$ -disjunctive Σ -logic and $\phi \approx \psi$ a semi-lattice/“distributive lattice” identity for $\bar{\wedge}$ or/and $\bar{\vee}$. Then, $\phi \equiv_C^\omega \psi$. \square

Theorem 4.4. *Let C be a \diamond -conjunctive/-disjunctive Σ -logic (defined by a class \mathbf{M} of simple Σ -matrices) and $i = (0/1)$ (as well as $\mathbf{K} \triangleq \pi_0[\mathbf{M}]$). Then, C is self-extensional iff the following hold:*

- (i) *each element of $\mathbf{IV}(C)(= \mathbf{V}(\mathbf{K}))$ is a \diamond -semi-lattice;*
- (ii) *for all $\bar{\varphi} \in (\mathbf{Fm}_\Sigma^\omega)^2$, $(\varphi_1 \in C(\varphi_0)) \Leftrightarrow |\Rightarrow (\mathbf{IV}(C) \models (\varphi_i \approx (\varphi_0 \diamond \varphi_1)))$.*

Proof. The “if” part is by Theorem 4.1(ii) \Rightarrow (i) and semi-lattice identities (more specifically, the commutativity one) for \diamond . Conversely, if C is self-extensional, then, by Theorem 4.1(i) \Rightarrow (iii), we have $\equiv_C^\omega = \theta_{\mathbf{IV}(C)}^\omega$, in which case, since C is \diamond -conjunctive/-disjunctive, (i) is by Remark 4.3 (and Theorem 3.19), while, for all $\bar{\varphi} \in (\mathbf{Fm}_\Sigma^\omega)^2$, $(\varphi_1 \in C(\varphi_0)) \Leftrightarrow (\varphi_i \equiv_C^\omega (\varphi_0 \diamond \varphi_1))$, so (ii) holds. \square

Lemma 4.5. *A [truth-non-empty $\bar{\wedge}$ -conjunctive] Σ -matrix \mathcal{A} is a $(2 \setminus 1)$ -model of a [finitary $\bar{\wedge}$ -conjunctive] Σ -logic C iff [f] $\mathcal{A} \in \text{Mod}(C)$ (cf. Definition 2.6).*

Proof. The “if” part is trivial. [Conversely, assume $\mathcal{A} \in \text{Mod}_{2 \setminus 1}(C)$. Consider any $\varphi \in C(\emptyset)$ and any $h \in \text{hom}(\mathfrak{Fm}_\Sigma^\omega, \mathfrak{A})$, in which case $V \triangleq \text{Var}(\varphi) \in \wp_\omega(\text{Var}_\omega)$, and so $(\text{Var}_\omega \setminus V) \neq \emptyset$, for, otherwise, we would have $V = \text{Var}_\omega$, and so would get $\omega = |\text{Var}_\omega| = |V| \in \omega$. Take any $v \in (\text{Var}_\omega \setminus V)$ and any $a \in D^{\mathcal{A}} \neq \emptyset$. Let $g \in \text{hom}(\mathfrak{Fm}_\Sigma^\omega, \mathfrak{A})$ extend $(h \upharpoonright (V \setminus \{v\})) \cup \{v/a\}$. Then, $\varphi \in C(v)$, $\{v\} \in \wp_{2 \setminus 1}(\mathbf{Fm}_\Sigma^\omega)$ and $g(v) = a \in D^{\mathcal{A}}$, in which case $h(\varphi) = g(\varphi) \in D^{\mathcal{A}}$, for $\mathcal{A} \in \text{Mod}_{2 \setminus 1}(C)$, and so $\mathcal{A} \in \text{Mod}_2(C)$. By induction on any $n \in \omega$, let us prove that $\mathcal{A} \in \text{Mod}_n(C)$. For consider any $X \in \wp_n(\mathbf{Fm}_\Sigma^\omega)$, in which case $n \neq 0$. In case $|X| \leq 2$, $X \in \wp_2(\mathbf{Fm}_\Sigma^\omega)$, and so $C(X) \subseteq \text{Cn}_\mathcal{A}^\omega(X)$, for $\mathcal{A} \in \text{Mod}_2(C)$. Otherwise, $|X| \geq 2$, in which case there are some distinct $\phi, \psi \in X$, and so $Y \triangleq ((X \setminus \{\phi, \psi\}) \cup \{\phi \bar{\wedge} \psi\}) \in \wp_{n-1}(\mathbf{Fm}_\Sigma^\omega)$. Then, by the induction hypothesis and the $\bar{\wedge}$ -conjunctivity of both C and \mathcal{A} , $C(X) = C(Y) \subseteq \text{Cn}_\mathcal{A}^\omega(Y) = \text{Cn}_\mathcal{A}^\omega(X)$. So, $\mathcal{A} \in \text{Mod}(C)$, as $\omega = (\bigcup \omega)$, and C is finitary.] \square

Theorem 4.6. *Let C be a $\bar{\wedge}$ -conjunctive [$\bar{\vee}$ -disjunctive] Σ -logic and $\mathbf{V} \triangleq \mathbf{IV}(C)$ (as well as \mathbf{M} a class of simple Σ -matrices defining C , and $\mathbf{K} \triangleq \pi_0[\mathbf{M}]$). {Suppose C is finitary (in particular, both \mathbf{M} and all elements of it are finite).} Then, (i) \Leftrightarrow (ii){ \Rightarrow }(iii) \Rightarrow (iv) \Rightarrow (i), where:*

- (i) *C is self-extensional;*
- (ii) *for all $\phi, \psi \in \mathbf{Fm}_\Sigma^\omega$, it holds that $(\psi \in C(\phi)) \Leftrightarrow |\Rightarrow (\mathbf{V} \models (\phi \approx (\phi \bar{\wedge} \psi)))$, while every element of \mathbf{V} is a $\bar{\wedge}$ -semi-lattice [resp., distributive $(\bar{\wedge}, \bar{\vee})$ -lattice];*
- (iii) *every truth-non-empty $\bar{\wedge}$ -conjunctive [consistent $\bar{\vee}$ -disjunctive] Σ -matrix with underlying algebra in \mathbf{V} is a model of C , while every element of \mathbf{V} is a $\bar{\wedge}$ -semi-lattice [resp., distributive $(\bar{\wedge}, \bar{\vee})$ -lattice];*
- (iv) *any truth-non-empty $\bar{\wedge}$ -conjunctive [consistent $\bar{\vee}$ -disjunctive] Σ -matrix with underlying algebra in \mathbf{K} is a model of C , while every element of \mathbf{K} is a $\bar{\wedge}$ -semi-lattice [resp., distributive $(\bar{\wedge}, \bar{\vee})$ -lattice].*

{(In particular, (i-iv) are equivalent.)}

Proof. First, (i) \Leftrightarrow (ii) is by Remark 4.3 and Theorem 4.4 with $i = 0$ and $\diamond = \bar{\wedge}$. {Next, (ii) \Rightarrow (iii) is by Lemma 4.5.} (Further, (iv) is a particular case of (iii), in

view of Theorem 3.19.) Finally, assume (iii) (resp., (iv)) holds. Let \mathbf{S} be the class of all truth-non-empty $\bar{\wedge}$ -conjunctive [consistent \vee -disjunctive] Σ -matrices with underlying algebra in \mathbf{V} (resp., in \mathbf{K}). Consider any $\mathcal{A} \in \mathbf{S}$ and any $\bar{a} \in (A^2 \setminus \Delta_A)$, in which case, by the semi-lattice identities (more specifically, the commutativity one) for $\bar{\wedge}$, $a_i \neq (a_i \bar{\wedge}^{\mathcal{A}} a_{1-i})$, for some $i \in 2$, and so $\mathcal{B} \triangleq (\mathfrak{A}, \{b \in A \mid a_i = (a_i \bar{\wedge}^{\mathcal{A}} b)\}) \in \mathbf{S}$ [resp., by the Prime Ideal Theorem, there is some $\mathcal{B} \in \mathbf{S}$] such that $\mathfrak{B} = \mathfrak{A}$ and $a_i \in D^{\mathcal{B}} \not\equiv a_{1-i}$. In this way, (i) is by Theorem(s) 4.1(vi) \Rightarrow (i) (and 3.19). \square

Theorem 4.7. *Let \mathbf{M} be a [finite] class of [finite hereditarily] simple [$\bar{\wedge}$ -conjunctive \vee -disjunctive] Σ -matrices, $\mathbf{K} \triangleq \pi_0[\mathbf{M}]$ and C the logic of \mathbf{M} . Then, C is self-extensional iff, for each $\mathfrak{A} \in \mathbf{K}$ and all distinct $a, b \in A$, there are some $\mathcal{B} \in \mathbf{M}$ and some $h \in \text{hom}(\mathfrak{A}, \mathfrak{B})$ such that $\chi^{\mathcal{B}}(h(a)) \neq \chi^{\mathcal{B}}(h(b))$.*

Proof. The “if” part is by Theorem 4.1(v) \Rightarrow (i) with $\mathbf{C} = \mathbf{K}$. [Conversely, assume C is self-extensional. Consider any $\mathfrak{A} \in \mathbf{K}$ and any $\bar{a} \in (A^2 \setminus \Delta_A)$. Then, by Theorem 4.6(i) \Rightarrow (iv), \mathfrak{A} is a distributive $(\bar{\wedge}, \vee)$ -lattice, in which case, by the commutativity identity for $\bar{\wedge}$, $a_i \neq (a_i \bar{\wedge}^{\mathfrak{A}} a_{1-i})$, for some $i \in 2$, and so, by the Prime Ideal Theorem, there is some $\bar{\wedge}$ -conjunctive \vee -disjunctive Σ -matrix \mathcal{D} with $\mathfrak{D} = \mathfrak{A}$ such that $a_i \in D^{\mathcal{D}} \not\equiv a_{1-i}$, in which case \mathcal{D} is both consistent and truth-non-empty, and so is a model of C . Hence, by Theorem 3.9 and Remark 2.5(ii), there are some $\mathcal{B} \in \mathbf{M}$ and some strict $h \in \text{hom}(\mathcal{D}, \mathcal{B}) \subseteq \text{hom}(\mathfrak{A}, \mathfrak{B})$, in which case $h(a_i) \in D^{\mathcal{B}} \not\equiv h(a_{1-i})$, so $\chi^{\mathcal{B}}(h(a_i)) = 1 \neq 0 = \chi^{\mathcal{B}}(h(a_{1-i}))$.] \square

4.2. Self-extensionality of implicative logics versus implicative intrinsic semi-lattices. A Σ -algebra \mathfrak{A} is called an \sqsupset -implicative intrinsic semi-lattice [with bound (a)], provided it is a \sqsupset -semi-lattice [with bound (a)] and satisfies:

$$(4.1) \quad (x_0 \sqsupset x_0) \approx (x_1 \sqsupset x_1),$$

$$(4.2) \quad ((x_0 \sqsupset x_0) \sqsupset x_1) \approx x_1,$$

in which case it is that with bound $a \sqsupset^{\mathfrak{A}} a$, for any $a \in A$.

Remark 4.8. Let C be a [self-extensional] Σ -logic and $\phi, \psi \in C(\emptyset)$, in which case $\phi \equiv_C^{\omega} \psi$ [and so $\text{IV}(C) \models (\phi \approx \psi)$]. \square

Theorem 4.9. *Let \mathbf{M} be an \sqsupset -implicative Σ -logic C (defined by a class \mathbf{M} of simple Σ -matrices and $\mathbf{K} \triangleq \pi_0[\mathbf{M}]$). Then, C is self-extensional iff, for all $\phi, \psi \in \text{Fm}_{\Sigma}^{\omega}$, it holds that $(\psi \in C(\phi)) \Leftrightarrow (\text{IV}(C) \models (\psi \approx (\phi \sqsupset \psi)))$, while each element of $\text{IV}(C) (= \mathbf{V}(\mathbf{K}))$ is an \sqsupset -implicative intrinsic semi-lattice.*

Proof. First, by (2.6), Remark 4.8 and the structurality of C , (4.1) $\in \equiv_C^{\omega}$. Likewise, by (2.6), (2.7) and (2.8), (4.2) $\in \equiv_C^{\omega}$. Then, Theorems 3.5(ii) and 4.4 with $i = 1$ and $\diamond = \sqsupset$ complete the argument. \square

Lemma 4.10. *Let C' be a finitary Σ -logic and C'' a 1-extension of C' (cf. Definition 2.3). Suppose C' has DT with respect to \sqsupset , while (2.8) is satisfied in C'' . Then, C'' is an extension of C' .*

Proof. By induction on any $n \in \omega$, we prove that C'' is an n -extension of C' . For consider any $X \in \wp_n(\text{Fm}_{\Sigma}^{\omega})$, in which case $n \neq 0$, and any $\psi \in C'(X)$. Then, in case $X = \emptyset$, we have $X \in \wp_1(\text{Fm}_{\Sigma}^{\omega})$, and so $\psi \in C'(X) \subseteq C''(X)$, for C'' is a 1-extension of C' . Otherwise, take any $\phi \in X$, in which case $Y \triangleq (X \setminus \{\phi\}) \in \wp_{n-1}(\text{Fm}_{\Sigma}^{\omega})$, and so, by DT with respect to \sqsupset , that C' has, and the induction hypothesis, we have $(\phi \sqsupset \psi) \in C'(Y) \subseteq C''(Y)$. Therefore, by (2.8)[$x_0/\phi, x_1/\psi$] satisfied in C'' , in view of its structurality, we eventually get $\psi \in C''(Y \cup \{\phi\}) = C''(X)$. Hence, as $\omega = (\bigcup \omega)$, we conclude that C'' is an extension of C' , for this is finitary. \square

Theorem 4.11. *Let \mathbf{M} be a [finite] class of [finite hereditarily] simple [\sqsupset -implicative] Σ -matrices, $\mathbf{K} \triangleq \pi_0[\mathbf{M}]$ and C the logic of \mathbf{M} . Then, C is self-extensional iff, for each $\mathfrak{A} \in \mathbf{K}$ and all distinct $a, b \in A$, there are some $\mathfrak{B} \in \mathbf{M}$ and some $h \in \text{hom}(\mathfrak{A}, \mathfrak{B})$ such that $\chi^{\mathfrak{B}}(h(a)) \neq \chi^{\mathfrak{B}}(h(b))$.*

Proof. The “if” part is by Theorem 4.1(v) \Rightarrow (i) with $\mathbf{C} = \mathbf{K}$. [Conversely, assume C is self-extensional. Consider any $\mathfrak{A} \in \mathbf{K}$ and any $\bar{a} \in (A^2 \setminus \Delta_A)$. Then, by Theorem 4.9, $\mathfrak{A} \in \text{IV}(C)$ is an \sqsupset -implicative intrinsic semi-lattice, in which case, by the commutativity identity for \sqsupset , $a_{1-i} \neq (a_i \sqsupset^{\mathfrak{A}} a_{1-i})$, for some $i \in 2$. Let $n \triangleq |A| \in (\omega \setminus 1)$. Take any bijection $\bar{c} : n \rightarrow A$. Let $g \in \text{hom}(\mathfrak{Fm}_{\Sigma}^{\omega}, \mathfrak{A})$ extend $[x_j/c_j; x_k/c_0]_{j \in n; k \in (\omega \setminus n)}$, in which case $A = (\text{img } \bar{c}) \subseteq (\text{img } g) \subseteq A$, and so there is some $\bar{\varphi} \in (\text{Fm}_{\Sigma}^{\omega})^2$ such that $g(\bar{\varphi}) = \bar{a}$. Then, by (2.15), $S \triangleq g^{-1}[\text{Fg}_C^{\mathfrak{A}}(\emptyset)] \in \text{Fi}_C(\mathfrak{Fm}_{\Sigma}^{\omega})$. Let us prove, by contradiction, that $\varphi_{1-i} \notin T \triangleq C(S \cup \{\varphi_i\})$. For suppose $\varphi_{1-i} \in T$, in which case, by DT, $(\varphi_i \sqsupset \varphi_{1-i}) \in C(S)$, and so $(\varphi_i \sqsupset \varphi_{1-i}) = \sigma(\varphi_i \sqsupset \varphi_{1-i}) \in S$, for $\sigma[S] = S \subseteq S$, where σ is the diagonal Σ -substitution. Then, $(a_i \sqsupset^{\mathfrak{A}} a_{1-i}) \in \text{Fg}_C^{\mathfrak{A}}(\emptyset)$. Clearly, by (2.6), $F \triangleq \{a_i \sqsupset^{\mathfrak{A}} a_i\} \subseteq \text{Fg}_C^{\mathfrak{A}}(\emptyset)$. Conversely, consider any $\phi \in C(\emptyset)$ and any $e \in \text{hom}(\mathfrak{Fm}_{\Sigma}^{\omega}, \mathfrak{A})$, in which case, by the structurality of C , $\sigma'(\phi) \in C(\emptyset)$, where σ' is the Σ -substitution extending $[x_l/x_{l+1}]_{l \in \omega}$, and so, by (2.6) and Remark 4.8, $e(\phi) = e'(\sigma'(\phi)) = e'(x_0 \sqsupset x_0) = (a_i \sqsupset^{\mathfrak{A}} a_i) \in F$, where $e' \in \text{hom}(\mathfrak{Fm}_{\Sigma}^{\omega}, \mathfrak{A})$ extends $[x_0/a_i; x_{m+1}/e(x_m)]_{m \in \omega}$ (in particular, $\mathcal{D} \triangleq \langle \mathfrak{A}, F \rangle \in \text{Mod}_1(C)$; cf. Definition 2.6). And what is more, by (4.2), (2.8) is true in \mathcal{D} , in which case, by Lemma 4.10, $F \in \text{Fi}_C(\mathfrak{A})$, and so $\text{Fg}_C^{\mathfrak{A}}(\emptyset) \subseteq F$ (in particular, $\text{Fg}_C^{\mathfrak{A}}(\emptyset) = F$). In this way, $(a_i \sqsupset^{\mathfrak{A}} a_{1-i}) = (a_i \sqsupset^{\mathfrak{A}} a_i)$, in which case, by (4.2), $(a_i \sqsupset^{\mathfrak{A}} a_{1-i}) = ((a_i \sqsupset^{\mathfrak{A}} a_i) \sqsupset^{\mathfrak{A}} a_{1-i}) = a_{1-i}$, and so this contradiction shows that $\varphi_{1-i} \notin T$. Hence, there are some $\mathfrak{B} \in \mathbf{M}$ and some $f \in \text{hom}(\mathfrak{Fm}_{\Sigma}^{\omega}, \mathfrak{B})$ such that $(S \cup \{\varphi_i\}) \subseteq f^{-1}[D^{\mathfrak{B}}] \not\cong \varphi_{1-i}$, in which case $\mathcal{E} \triangleq \langle \mathfrak{B}, (\text{img } f) \rangle$, being a submatrix of \mathfrak{B} , is simple, for \mathfrak{B} is hereditarily so, as well as, by Remark 2.7(ii), is \sqsupset -implicative, for \mathfrak{B} is so, while $U \triangleq f^{-1}[D^{\mathfrak{E}}] = f^{-1}[D^{\mathfrak{B}}] \in (\text{img } C)$, whereas f is a surjective strict homomorphism from $\mathcal{F} \triangleq \langle \mathfrak{Fm}_{\Sigma}^{\omega}, U \rangle$ onto \mathcal{E} , and so, by Remark 2.5(iii), $\mathcal{D}(\mathcal{F}) = f^{-1}[\mathcal{D}(\mathcal{E})] = f^{-1}[\Delta_{\mathcal{E}}] = (\ker f)$. Consider any $\bar{\psi} \in \theta \triangleq (\ker g)$, in which case, by (2.6), for all $\ell \in 2$, $g(\psi_{\ell} \sqsupset \psi_{1-\ell}) = (g(\psi_{\ell}) \sqsupset^{\mathfrak{A}} g(\psi_{1-\ell})) = (g(\psi_{\ell}) \sqsupset^{\mathfrak{A}} g(\psi_{\ell})) \in \text{Fg}_C^{\mathfrak{A}}(\emptyset)$, and so $(\psi_{\ell} \sqsupset \psi_{1-\ell}) \in S \subseteq U$. Hence, by (2.8), $(\psi_{\ell} \in U) \Rightarrow (\psi_{1-\ell} \in U)$, in which case $\bar{\psi} \in \theta^{\mathcal{F}}$, and so $\theta^{\mathcal{F}} \supseteq \theta \in \text{Con}(\mathfrak{Fm}_{\Sigma}^{\omega})$. Therefore, $\theta \in \text{Con}(\mathcal{F})$, in which case $\theta \subseteq \mathcal{D}(\mathcal{F}) = (\ker f)$, and so, by the Homomorphism Theorem, $h \triangleq (g^{-1} \circ f) \in \text{hom}(\mathfrak{A}, \mathfrak{B})$, while $h(a_i) = f(\varphi_i) \in D^{\mathfrak{B}} \not\cong f(\varphi_{1-i}) = h(a_{1-i})$. \square

4.3. Self-extensionality of uniform finitely-valued logics versus truth discriminators. A truth discriminator for/of a Σ -matrix \mathcal{A} is any $\bar{h} : \text{img}[\theta^{\mathcal{A}} \setminus \Delta_{\mathcal{A}}] \rightarrow \text{hom}(\mathfrak{A}, \mathfrak{A})$ such that, for every $\{a, b\} \in (\text{dom } \bar{h})$, $\langle a, b \rangle \notin \ker(h_{\{a, b\}} \circ \chi^{\mathcal{A}})$. Then, since $\Delta_{\mathcal{A}} \in \text{hom}(\mathfrak{A}, \mathfrak{A})$, by Theorems 4.7 and 4.11, we have:

Corollary 4.12. *Let \mathcal{A} be a [finite hereditarily] simple [either implicative or both conjunctive and disjunctive] Σ -matrix and C the logic of \mathcal{A} . Then, C is self-extensional iff \mathcal{A} has a truth discriminator.*

The effective procedure of verifying the self-extensionality of the logic of an n -valued, where $n \in (\omega \setminus 1)$, hereditarily simple either implicative or both conjunctive and disjunctive Σ -matrix resulted from Corollary 4.12 has the computational complexity n^{n+2} that is quite acceptable for (3|4)-valued logics. And what is more, it provides a quite useful heuristic tool of doing it, manual applications of which (suppressing the factor n^{n+2} at all) are presented below.

Corollary 4.13. *The logic of any no-less-than-three-valued hereditarily simple either implicative or both conjunctive and disjunctive Σ -matrix \mathcal{A} without non-diagonal non-singular (cf. p. 2) endomorphism of \mathfrak{A} is not self-extensional.*

Proof. By contradiction. For suppose the logic of \mathcal{A} is self-extensional, in which case, as $|A| \geq 3 \not\leq 2$, $\chi^{\mathcal{A}}$ is not injective, and so there are some distinct $a, b \in A$ such that $\chi^{\mathcal{A}}(a) = \chi^{\mathcal{A}}(b)$. Then, by Corollary 4.12, there is some $h \in \text{hom}(\mathfrak{A}, \mathfrak{A})$ such that $\chi^{\mathcal{A}}(h(a)) \neq \chi^{\mathcal{A}}(h(b))$, in which case $h(a) \neq h(b)$, and so h is not singular (in particular, diagonal). Hence, $\chi^{\mathcal{A}}(a) = \chi^{\mathcal{A}}(h(a)) \neq \chi^{\mathcal{A}}(h(b)) = \chi^{\mathcal{A}}(b) = \chi^{\mathcal{A}}(a)$. \square

4.3.1. Self-extensionality versus equational implications and unitary equality determinants. According to [15], given any $m, n \in \omega$, a [finitary] (Σ) -equational \vdash_n^m -sequent definition for/of a Σ -matrix \mathcal{A} is any $\mathfrak{U} \in \wp_{[\omega]}(\text{Eq}_{\Sigma}^{m+n})$ such that, for all $\bar{a} \in A^m$ and all $\bar{b} \in A^n$, it holds that $((\text{img } a) \subseteq D^{\mathcal{A}} \Rightarrow ((\text{img } b) \cap D^{\mathcal{A}}) \neq \emptyset) \Leftrightarrow (\mathfrak{A} \models (\bigwedge \mathfrak{U})[x_i/a_i; x_{m+j}/b_j]_{i \in m; j \in n})$. Equational $\vdash_1^{0/1}$ -definitions are also referred to as equational “truth [predicate] definitions”/implications, / (cf. [16]), respectively. Some kinds of equational sequent definitions are equivalent, by:

Remark 4.14. Given a(n \sqsupset -implicative) Σ -matrix \mathcal{A} , each of (i,(ii-iv)) holds, where:

- (i) given a [finitary] equational \vdash_2^2 -definition \mathfrak{U} for \mathcal{A} , $\mathfrak{U}[x_{(2 \cdot i)+j}/x_i]_{i,j \in 2}$ is a [finitary] equational implication for \mathcal{A} (cf. Theorems 10 and 12(ii) \Rightarrow (iii) of [15]);
- (ii) given any [finitary] equational implication \mathfrak{U} for \mathcal{A} , $\mathfrak{U}[x_0/(x_0 \sqsupset x_0), x_1/x_0]$ is a [finitary] equational truth definition for \mathcal{A} ;
- (iii) given any [finitary] equational truth definition \mathfrak{U} for \mathcal{A} , $\mathfrak{U}[x_0/(x_0 \sqsupset (x_1 \sqsupset (x_2 \sqcup x_3)))]$ is a [finitary] equational \vdash_2^2 -definition for \mathcal{A} ;
- (iv) in case \mathcal{A} is truth-singular, $\{x_0 \approx (x_0 \sqsupset x_0)\}$ is a finitary equational truth definition for it. \square

In this way, taking Theorems 10, 12(i) \Leftrightarrow (ii) and 13 of the work [15] as well as Remark 4.14 into account, an either implicative or both conjunctive and disjunctive consistent truth-non-empty finite Σ -matrix \mathcal{M} with unitary equality determinant has a finitary equational implication iff the multi-conclusion two-side sequent calculus $\tilde{\mathcal{S}}_{\mathcal{M}, \mathcal{T}}^{(k,l)}$ (cf. [14] as well as the paragraph -2 on p. 294 of [15] for more detail) is algebraizable (in the sense of [13, 12]). In this connection, by Lemma 9 and Theorem 10 of [15] as well as Corollary 4.13, we immediately get:

Corollary 4.15. *The logic of any no-less-than-tree-valued either implicative or both conjunctive and disjunctive Σ -matrix with unitary equality determinant and equational implication is not self-extensional.*

In view of Theorem 10 and Lemma 8 of [15], Example 4.2 [with $\Sigma = \{\perp, \top\}$, $\chi^{\mathcal{A}} = \Delta_2$ and $\mathfrak{A} = (\mathfrak{D}_{2,01} \upharpoonright \Sigma)$] and the self-extensionality of inferentially inconsistent (in particular, one-valued) logics, the stipulation “no-less-than-tree-valued” cannot be omitted in the formulation[s] of Corollary 4.15 [as well as Corollary 4.13].

Example 4.16 (Łukasiewicz’ finitely-valued logics; cf. [6]). Let $n \in (\omega \setminus 3)$, $\Sigma \triangleq (\Sigma_+ \cup \{\sim, \supset\})$ with binary \supset (implication) and unary \sim (negation) and \mathcal{A} the Σ -matrix with $(\mathfrak{A} \upharpoonright \Sigma_+) \triangleq \mathfrak{D}_n$ (cf. Subparagraph 2.2.1.2.1), $D^{\mathcal{A}} \triangleq \{1\}$, $\sim^{\mathfrak{A}} \triangleq (1 - a)$ and $(a \supset^{\mathfrak{A}} b) \triangleq \min(1, 1 - a + b)$, for all $a, b \in A$, in which case \mathcal{A} is both consistent, truth-non-empty, \wedge -conjunctive and \vee -disjunctive as well as has both an equational implication, by Example 7 of [15], and a unitary equality determinant, by Example 3 of [14]. Hence, by Corollary 4.15, the logic of \mathcal{A} is not self-extensional. \square

Example 4.17. By Example 2 of [14], Remark 1 as well as Theorem 10 and Lemma 9 of [15] and Corollaries 4.13 and 4.15, arbitrary three-valued expansions of both the *logic of paradox LP* [9] and Kleene’s three-valued logic K_3 [4] are not self-extensional, for the matrix defining the former has the equational implication

$(x_0 \wedge (x_1 \vee \sim x_1)) \approx (x_0 \wedge x_1)$, discovered in [11], while the matrix defining the latter has the same underlying algebra as that defining the former. Likewise, by “both Lemma 4.1 of [10] and Remark 4.14(i,iii)”/“Proposition 5.7 of [16]” as well as Corollary 4.15, arbitrary three-valued expansions of P^1/HZ [21]/[3] are not self-extensional, for their being defined by implicative/ matrices with equational “truth definition”/implication. \square

5. APPLICATIONS TO NO-MORE-THAN-FOUR-VALUED LOGICS

All along throughout this section, $(\lrcorner \triangleq \sim)/ \supset$ is supposed to be a primary unary/binary connective of Σ viewed as negation/implication. Let $\Sigma_{\sim(+)[01]}^{\langle \supset \rangle \{ \bar{\varsigma} \}} \triangleq (\{ \sim \} \cup \Sigma_+ \cup \Sigma_{01} \cup \{ \supset \} \cup \{ \text{img } \bar{\varsigma} \})$ [(cf. Subparagraph 2.2.1.2.1)] {where $\bar{\varsigma}$ is a finite sequence of primary connectives beyond $\Sigma_{\sim(+)[01]}^{\langle \supset \rangle}$ }.

5.1. No-more-than-four-valued extensions of uniform four-valued expansions of Belnap’s four-valued logic. A [bounded] De Morgan lattice [13] is any $\Sigma_{\sim,+[01]}$ -algebra, with [bounded] distributive lattice $\Sigma_{+[01]}$ -reduct satisfying:

$$(5.1) \quad \sim \sim x_0 \approx x_0,$$

$$(5.2) \quad \sim(x_0 \vee x_1) \approx (\sim x_0 \wedge \sim x_1),$$

By $\mathfrak{DM}_{4[01]}$ we denote the non-Boolean diamond [bounded] De Morgan lattice with $(\mathfrak{DM}_{4[01]} \upharpoonright \Sigma_{+[01]}) \triangleq \mathfrak{D}_{2[01]}^2$ and $\sim^{\mathfrak{DM}_{4[01]}}(i, j) \triangleq \langle 1 - j, 1 - i \rangle$, for all $i, j \in 2$.

Here, it is supposed that $\Sigma \supseteq \Sigma_{\sim,+[01]}$ and $(\bar{\wedge} | \bar{\vee}) = (\wedge | \vee)$. Fix a Σ -matrix \mathcal{A} with $(\mathfrak{A} \upharpoonright \Sigma_{\sim,+[01]}) \triangleq \mathfrak{DM}_{4[01]}$ and $D^{\mathcal{A}} \triangleq (2^2 \cap \pi_0^{-1}[\{1\}])$. Then, \mathcal{A} as well as its submatrices are both \wedge -conjunctive and \vee -disjunctive as well as both consistent and truth-non-empty (cf. Remark 2.7(ii)(a,b)), while $\{x_0, \sim x_0\}$ is a unitary equality determinant for them (cf. Example 2 of [14]), so they are hereditarily simple (cf. Lemma 3.1). Let C be the logic of \mathcal{A} . Then, since $\mathcal{DM}_{4[01]} \triangleq (\mathcal{A} \upharpoonright \Sigma_{\sim,+[01]})$ defines [the bounded version/expansion of] Belnap’s four-valued logic $B_{4[01]}$ [1] (cf. [13, 18, 17, 19]), C is a uniform four-valued expansion of $B_{4[01]}$. Conversely, according to Corollary 4.9 of [18], any uniform four-valued expansion of $B_{4[01]}$ is defined by a unique expansion of $\mathcal{DM}_{4[01]}$, in which case \mathcal{A} is uniquely determined by C , and so is said to be *characteristic for/of* C . Moreover, by (2.15), Remark 2.5(ii) and Theorem 3.9, C is \sim -subclassical iff Δ_2 forms a subalgebra of \mathfrak{A} , in which case $\mathcal{A} \upharpoonright 2$ is isomorphic to any \sim -classical model of C , and so defines a unique \sim -classical extension of C (cf. Theorem 4.20 of [18]) denoted by C^{PC} and relatively axiomatized as follows, in view of Corollary 2.9 of [18] and Theorem 3.11, for $\mathcal{A} \upharpoonright 2$ is then the only non- \sim -paraconsistent non- (\vee, \sim) -paracomplete submatrix of \mathcal{A} :

Lemma 5.1. *If C is \sim -classical, C^{PC} is relatively axiomatized by $\{(2.12), (3.2)\}$.*

Given any $i \in 2$, put $DM_{3,-,i} \triangleq (2^2 \setminus \{\langle i, 1 - i \rangle\})$. Then, we have the submatrix $\mathcal{A}_{3,i}$ generated by $DM_{3,-,i}$ with carrier (not) distinct from the generating set (in particular, when, e.g., $\Sigma = \Sigma_{\sim,+[01]}$), taking (2.15) into account, the logic $C_{3,i}$ of which is a both \vee -disjunctive and \wedge -conjunctive {for its defining matrix is so} as well as inferentially consistent {for its defining matrix is both consistent and truth-non-empty} uniform no-more-than-four-valued extension of C (and a three-valued expansion of [the bounded version/expansion $LP_{01}|K_{3,01}$ of] “the logic of paradox”/“Kleene’s three-valued logic” $LP|K_3$ [9] [4], whenever $i = (0|1)$, for $\mathcal{DM}_{3,i[01]} \triangleq (\mathcal{A}_{3,i} \upharpoonright \Sigma_{\sim,+[01]})$ defines $LP_{[01]}|K_{3[01]}$), in which case it is \vee -disjunctive as well as \sim -paraconsistent/ (\vee, \sim) -paracomplete, and so is not \sim -classical, in view of Remark 2.7(i)(a).

Lemma 5.2 (Key 4-valued Lemma). *Let $\mathcal{B} \in \text{Mod}(C)$. Then, the following hold:*

- (i) \mathcal{B} is \vee -disjunctive, whenever it is either inconsistent or truth-empty or \sim -negative or [non- \sim -classically-defining or] no-more-than- $(4[-1])$ -valued;
- (ii) providing \mathcal{B} is \vee -disjunctive [and (not) truth-empty] “either \sim -negative or \sim -classically-defining” \parallel \sim -paraconsistent/ (\vee, \sim) -paracomplete], it is a strictly surjectively homomorphic counter-image of a submatrix of \mathcal{A} with carrier in $S_{4[+(-)\emptyset|C|P/PC]} \triangleq ((\{\{01\}, \Delta_2, 2^2\} \cup \{D_{3,-,l} \mid l \in 2\})[\cap(\setminus)(\{\{01\}|\Delta_2\}||\{2^2, DM_{3,-,0/1}\})])$.

Proof. (i) By contradiction. For suppose \mathcal{B} is not \vee -disjunctive. Then, taking Remarks 2.5(iv), 2.7(ii)(a,b) and (2.15) into account, without loss of generality, one can assume that \mathcal{B} is simple, in which case, by Corollary 3.15 and Theorem 3.19, \mathfrak{B} belongs to the variety generated by \mathfrak{A} , and so $\mathfrak{B}|\Sigma_{\sim,+}$ is a De Morgan lattice (in particular, $\mathfrak{B}|\Sigma_+$ is a distributive lattice), for $(\mathfrak{A}|\Sigma_{\sim,+}) = \mathfrak{DM}_4$ is so. And what is more, $\mathcal{B} \in \text{Mod}(C)$ is both \wedge -conjunctive and weakly \vee -disjunctive, for C is so. Hence, since \mathcal{B} is not \vee -disjunctive, there are some $a, b \in (D \setminus D^{\mathcal{B}})$, in which case $c \triangleq (a \wedge^{\mathfrak{B}} b) \notin D^{\mathcal{B}}$, such that $d \triangleq (a \vee^{\mathfrak{B}} b) \in D^{\mathcal{B}}$ (in particular, \mathcal{B} is both consistent and truth-non-empty), in which case $d \notin \{a, b, c\}$, and so $|\{a, b, c, d\}| = 4$. Therefore, if \mathcal{B} was \sim -negative, then, by its \wedge -conjunctivity and (5.2), we would have $D^{\mathcal{B}} \not\equiv \sim^{\mathfrak{B}} d = (\sim^{\mathfrak{B}} a \wedge^{\mathfrak{B}} \sim^{\mathfrak{B}} b) \in D^{\mathcal{B}}$. Thus, $|B| \leq 4$, in which case $B = \{a, b, c, d\}$ (in particular, $|B| = 4 \not\leq 3$), and so \mathcal{B} is not \sim -classically-defining. In this way, \mathfrak{B} is a distributive (\wedge, \vee) -lattice with zero c and unit d , in which case, by (5.1) and (5.2), $\sim^{\mathfrak{B}}(c|d) = (d|c)$, and so, by (5.1), $\sim^{\mathfrak{B}}[\{a, b\}] \subseteq \{a, b\}$, for $(\{a, b\} \cap \{c, d\}) = \emptyset$. Consider the following cases:

- $\sim^{\mathfrak{B}} a = a$, in which case, by (5.1), $\sim^{\mathfrak{B}} b = b$, and so $e \triangleq \{\langle a, 10 \rangle, \langle b, 01 \rangle, \langle c, 00 \rangle, \langle d, 11 \rangle\}$ is an isomorphism from $\mathfrak{B}|\Sigma_{\sim,+}$ onto \mathfrak{DM}_4 . Furthermore, by Lemma 3.7, there are some finite set I , some $\bar{C} \in \mathbf{S}_*(\mathcal{A})^I$, some subdirect product \mathcal{D} of it and some $h \in \text{hom}_{\mathfrak{S}}^{\mathfrak{S}}(\mathcal{D}, \mathcal{B})$, in which case, $(\{h \circ e\} \cup \{\pi_i \upharpoonright D \mid i \in I\}) \in \wp_{\omega}(\text{hom}(\mathfrak{D}|\Sigma_{\sim,+}, \mathfrak{DM}_4))$, while, by Remark 2.7(ii)(b), \mathcal{D} is consistent (in particular, $I \neq \emptyset$), for \mathcal{B} is so, whereas $(\bigcap_{i \in I} \ker(\pi_i \upharpoonright D)) = \Delta_D \subseteq \ker(h \circ e) \neq D^2$, for $\text{img}(h \circ e) = DM_4 = 2^2$ is not a singleton, and so, by Theorem 3.8 of [18], there is some $i \in I$ such that $\ker(\pi_i \upharpoonright D) = \ker(h \circ e) = (\ker h)$, for e is injective. Therefore, by the Homomorphism Theorem, as $(\text{img } h) = B$, $h^{-1} \circ \pi_i$ is an embedding of \mathcal{B} into \mathcal{A} , in which case, by Remark 2.7(ii)(a), \mathcal{B} is \vee -disjunctive.
 - $\sim^{\mathfrak{B}} a \neq a$, in which case $\sim^{\mathfrak{B}} a = b$, and so, by (5.1), $\sim^{\mathfrak{B}} b = a$. Then, for each $e' \in B$, $(e'(\wedge|\vee)^{\mathfrak{B}} \sim^{\mathfrak{B}} e') = (c|d) \notin \{e' \mid e' \in D^{\mathcal{B}}\}$, in which case \mathcal{B} , being \wedge -conjunctive, satisfies both (2.12) and (3.2). And what is more, $\{c, d\}$ forms a subalgebra of \mathfrak{B} , in which case, by (2.15), $\mathcal{B}|\{c, d\}$ is a \sim -classical model of C , and so this is \sim -subclassical. Then, by Lemma 5.1, $\mathcal{B} \in \text{Mod}(C^{\text{PC}})$. Conversely, the logic of the consistent truth-non-empty model \mathcal{B} of C is an inferentially consistent extension of C , in which case, by Theorem 4.21 of [18], \mathcal{B} is \sim -classically-defining.
- (ii) Since $S_{4[+(-)\emptyset|C|P/PC]}$ is the set of the carriers of all [those] elements of $\mathbf{S}_*(\mathcal{A})$ [which are (not) truth-empty] “either \sim -negative or \sim -classically-defining” \parallel \sim -paraconsistent/ (\vee, \sim) -paracomplete], (2.15), Remarks 2.5 (ii), 2.7(ii)(a,b) and Theorem 3.9 complete the argument. \square

By Theorem 4.10 of [18], (2.15), Examples 4.2, 4.17, Lemma 5.2 and the self-extensionality of inferentially inconsistent logics, we first have:

Theorem 5.3. *Let C' be a uniform no-more-than-four-valued proper (in particular, no-more-than-three-valued) extension of C . Then, the following are equivalent:*

- (i) C' is self-extensional;

- (ii) C' is either inferentially inconsistent or \sim -classical;
- (iii) for each $i \in 2$, if $DM_{3,-,i}$ forms a subalgebra of \mathfrak{A} , then $C' \neq C_{3,i}$.

Since $\mathcal{DM}_4 \upharpoonright \{01\}$ is the only truth-empty submatrix of \mathcal{DM}_4 , while $\{01\} \subseteq [\not\subseteq] DM_{3,-,1[-1]} \supseteq \Delta_2$, by Theorem 4.10 of [18], (2.15) and Lemma 5.2, we also get:

Theorem 5.4. *Let \mathbf{M} be a class of no-more-than-four-valued models of C , C' the logic of \mathbf{M} , $\mathbf{M}_{\{01\}}^{(*)[\sim/\not\sim]}$ the class of all (truth-non-empty) [\sim -classically-/non- \sim -classically-defining] [\sim -paraconsistent] (\vee, \sim)-paracomplete} consistent elements of \mathbf{M} and $\mathbf{M}_2 = (\mathbf{M}_0 \cap \mathbf{M}_1)$. Then, C' is defined by $\{\mathcal{A} \mid \mathbf{M}_2 \neq \emptyset\} \cup \{\mathcal{A} \upharpoonright \{01\} \mid (\mathbf{M} \setminus \mathbf{M}^*) \neq \emptyset = \mathbf{M}_1^{*,\not\sim} = \mathbf{M}_2\} \cup \{\mathcal{A} \upharpoonright \Delta_2 \mid (\bigcup_{i \in 2} \mathbf{M}_i^{*,\not\sim}) = \mathbf{M}_2 = \emptyset \neq \mathbf{M}^\sim\} \cup \bigcup_{i \in 2} \{\mathcal{A}_{3,i} \mid \mathbf{M}_i^{*,\not\sim} \neq \emptyset = \mathbf{M}_2\}$. In particular, C is defined by any both \sim -paraconsistent and (\vee, \sim) -paracomplete no-more-than-four-valued model, so it has no both \sim -paraconsistent and (\vee, \sim) -paracomplete no-more-than-three-valued model.*

Taking (2.13), Theorems 5.3, 5.4, Remark 2.4 and Example 4.2 into account, it only remains to study the following no-more-than-four-valued extensions of C .

5.1.1. *Double three-valued and non-proper extensions.* By (2.15), (providing, for each $i \in 2$, $DM_{3,i}$ forms a subalgebra of \mathfrak{A}) the logic C_3 of $\{\mathcal{A}_{3,0}, \mathcal{A}_{3,1}\}$ is a both \vee -disjunctive and \wedge -disjunctive {for its defining matrices are so} as well as inferentially-consistent {for its defining matrices are both consistent and truth-non-empty} (proper) extension of C (for this is minimally four-valued; cf. Theorem 4.10 of [18]). Let $\mu : 2^2 \rightarrow 2^2$, $\langle i, j \rangle \mapsto \langle j, i \rangle$ be the *mirror/specular* function.

Theorem 5.5 (cf. [17, 19]). *It holds that (v) \Leftarrow (i) \Leftrightarrow (ii) \Leftrightarrow (iii) \Rightarrow (iv) \Rightarrow (iii) \Rightarrow (ii) \Rightarrow (i), where:*

- (i) $C_{\{3\}}$ is self-extensional;
- (ii) [for each/some $i \in 2$] $(\mu[\upharpoonright A_{3,i}]) \in \text{hom}(\mathfrak{A}_{[3,i]}, \mathfrak{A})$;
- (iii) there is a non-singular {cf. p. 2} non-diagonal (injective) homomorphism from [a subalgebra of] \mathfrak{A} to \mathfrak{A} ;
- (iv) \mathcal{A} has no equational implication;
- (v) $C_{\{3\}}$ is \sim -subclassical.

In particular, C_3 is self-extensional, whenever C is so.

Proof. First, the fact that (iv) is equivalent to the $(\)$ -non-optional \square -optional version of (iii) is due to Theorems 10, 13 and 15 of [15], while the \square -optional version of (iii) is a particular case of the \square -non-optional one, whereas the $(\)$ -non-optional version of (iii) is a particular case of the $(\)$ -optional one, being, in its turn, a particular case of (ii), for μ is injective. Next, the fact that (i) implies the $(\)$ -non-optional version of (iii) is by Theorem 4.7, for $D^{\mathcal{A}}[\cap DM_{3,-,0}]$ has two distinct elements. [Furthermore, by the injectivity of μ and the fact that, for any $i \in 2$, $\mu[DM_{3,-,i}] = DM_{3,-,1-i}$, while $2 = \{i, 1-i\}$, the alternatives in (ii) are equivalent.] Further, assume (ii) holds. Consider [any $i \in 2$ and] any distinct $a, b \in A_{[3,i]}$, in which case there is some $j \in 2$ such that $\pi_j(a) \neq \pi_j(b)$, and so $\chi^{\mathcal{A}_{[3,k_j]}}(h_j(a)) \neq \chi^{\mathcal{A}_{[3,k_j]}}(h_j(b))$, where $[k_{01}] \triangleq (i|(1-i))$ and] $h_{01} \triangleq (\Delta_{A_{[3,i]}} | (\mu[\upharpoonright A_{3,i}])) \in \text{hom}(\mathfrak{A}_{[3,i]}, \mathfrak{A})$. In this way, Theorem 4.7 yields (i). Now, assume the $(\)$ -non-optional version of (iii) holds. Then, there is some non-diagonal homomorphism h from [a subalgebra of] \mathfrak{A} to \mathfrak{A} with $B \triangleq (\text{img } h)$ not being a singleton, in which case B forms a non-one-element subalgebra of \mathfrak{A} , and so does $D \triangleq (\text{dom } h)$. Hence, $\Delta_2 \subseteq (B \cap D)$. Then, both of $(\mathfrak{B}|\mathfrak{D}) \triangleq (\mathfrak{A} \upharpoonright (B|D))$ are (\wedge, \vee) -lattices with zero/unit $\langle 0/1, 0/1 \rangle$, for \mathfrak{A} is so, in which case, as $h \in \text{hom}(\mathfrak{D}, \mathfrak{B})$ is surjective, by Lemma 2.1, $h \upharpoonright \Delta_2$ is diagonal, and so, since h is not so, there is some $i \in 2$ such that $DM_{3,-,i} \subseteq D$ {in particular, $A_{[3,i]} \subseteq D$ }, while $h(\langle 1-i, i \rangle) \neq \langle 1-i, i \rangle$. On the other hand, for all $a \in A$, it holds that $(\sim^{\mathfrak{A}} a = a) \Leftrightarrow (a \notin \Delta_2)$, in which case $\sim^{\mathfrak{A}} h(\langle 1-i, i \rangle) = h(\sim^{\mathfrak{A}} \langle 1-i, i \rangle) = h(\langle 1-i, i \rangle)$, and so $h(\langle 1-i, i \rangle) =$

$\langle i, 1 - i \rangle$. And what is more, [if $A_{3,i} = A$, then] $\langle i, 1 - i \rangle \in D$, in which case we have $\langle \langle i, 1 - i \rangle (\wedge | \vee)^{\mathfrak{D}} (1 - i, i) \rangle = \langle 0|1, 0|1 \rangle$, and so, by the diagonality of $h|_{\Delta_2}$, we get $\langle h(\langle i, 1 - i \rangle) (\wedge | \vee)^{\mathfrak{A}} (i, 1 - i) \rangle = \langle h(\langle i, 1 - i \rangle) (\wedge | \vee)^{\mathfrak{A}} h(\langle 1 - i, i \rangle) \rangle = h(\langle 0|1, 0|1 \rangle) = \langle 0|1, 0|1 \rangle$ (in particular, $h(\langle i, 1 - i \rangle) = \langle 1 - i, i \rangle$). In this way, $\text{hom}(\mathfrak{D}, \mathfrak{A}) \ni h = (\mu|D)$, in which case, as $A_{[3,i]} \subseteq D$, $(\mu|A_{3,i}) \in \text{hom}(\mathfrak{A}_{[3,i]}, \mathfrak{A})$, and so (ii) holds. Finally, if $\Delta_2 \langle = (\bigcap_{i \in 2} DM_{3,-,i}) \subseteq (\bigcap_{i \in 2} A_{3,i}) \rangle$ does not form a subalgebra of \mathfrak{A} , then there are some $\zeta \in \Sigma$ of arity $n \in \omega$ and some $\bar{a} \in \Delta_2^n$ such that $b \triangleq \zeta^{\mathfrak{A}}(\bar{a}) \in (A_{[3,i]} \setminus \Delta_2)$ [where $i \triangleq \pi_1(b) \in 2$], in which case $\mu(b) \neq b = \zeta^{\mathfrak{A}}(\mu \circ \bar{a})$, and so $(\mu|A_{3,i}) \notin \text{hom}(\mathfrak{A}_{[3,i]}, \mathfrak{A})$. Thus, (ii) \Rightarrow (v) is by (2.15), so, as the \square -optional version of (ii) is a particular case of the non- \square -optional one, (i) \Leftrightarrow (ii) ends the proof. \square

As μ is not diagonal, according to Example 11 of [15], the optional and non-optional versions of the item (ii) of Theorem 5.5 are non-equivalent to one another, and so are those of (i/iii) (in particular, the converse of the final assertion of Theorem 5.5 does not hold). Theorem 5.5(ii) \Rightarrow (i) positively covers both $B_{4\{01\}[3]}$ and the *classically-negative* case, when $\Sigma = \Sigma_{\sim,+\{01\}}^-$ with unary \neg (classical — viz., Boolean — negation) and $\neg^{\mathfrak{A}} \langle i, j \rangle \triangleq \langle 1 - i, 1 - j \rangle$, for all $i, j \in 2$, being the complement operation (cf. [13] and Subsection 5.1 of [18]), and so \mathfrak{A} has no three-element subalgebra. In view of Theorem 5.5(i) \Rightarrow (iv), the self-extensionality of these three instances of uniform four-valued expansions of B_4 provides a new insight and a new proof (convergent with those given by [15]) to the non-algebraizability of the sequent calculi associated (according to [14]) with their characteristic matrices, proved originally in [13] by a quite different (though equally generic) method based upon universal tools elaborated in [12]. This well justifies the thesis of the first paragraph of Section 1. Conversely, using Theorem 5.5(i) \Rightarrow (iv) / “and Remark 4.14”, we immediately conclude that arbitrary bilattice/implicative (in the /restricted sense of Subsection 5.2/5.3 of [18], respectively) uniform four-valued expansions of B_4 , when $\Sigma \supseteq \Sigma_{\sim,+\{01\}}^{\langle \square, \sqcup \rangle / \supset}$ “with binary \square and \sqcup ” / and $\langle \langle i, j \rangle (\langle \square | \sqcup \rangle / \supset)^{\mathfrak{A}} \langle k, l \rangle \rangle \triangleq \langle \langle (\min | \max)(i, k), (\max | \min)(j, l) \rangle / \langle \max(1 - i, k), \max(1 - i, l) \rangle \rangle$, for all $i, j, k, l \in 2$, / “as well as their double three-valued extensions in the *purely-implicative* case $\Sigma = \Sigma_{\sim,+\{01\}}^{\supset}$ ” are not self-extensional, for their $/\supset$ -implicative characteristic matrices have equational “implication $\{(((x_0 \sqcup \sim x_0) \sqcup (x_1 \sqcup \sim x_1)) \wedge x_0) \lesssim (((x_0 \sqcup \sim x_0) \sqcup (x_1 \sqcup \sim x_1)) \vee x_1)\}$, in view of the proof of Theorem 4.30 of [13]” / “truth definition $\{x_0 \approx (x_0 \supset x_0)\}$ ”. According to Corollary 5.2/5.3 of [18], this does equally/not ensue from Theorem 5.5(i) \Rightarrow (v) / “so refuting the inverse”.

Finally, since inferentially inconsistent logics are self-extensional, by (2.13), Theorems 5.3, 5.4, 5.5(i) \Leftrightarrow (iii) \Rightarrow (iv)[\Rightarrow (iii)], Remark 2.4 and Example 4.2, we get:

Theorem 5.6. *Let \mathbf{M} be a class of no-more-than-four-valued models of C and C' the logic of \mathbf{M} . Then, C' is self-extensional iff either \mathbf{M} contains no non- \sim -classically-defining truth-non-empty consistent element or there are a non-diagonal non-singular homomorphism from [a subalgebra of] \mathfrak{A} to \mathfrak{A} [i.e., \mathfrak{A} has no equational implication] as well as both \sim -paraconsistent and [truth-non-empty] (\vee, \sim) -paracomplete [distinct] element[s] of \mathbf{M} . In particular, any inferentially consistent non- \sim -classical no-more-than-four-valued extension of C is self-extensional only if it is both \sim -paraconsistent and (\vee, \sim) -paracomplete.*

5.2. Uniform three-valued logics with subclassical negation.

5.2.1. *U3VLSN versus super-classical matrices.* Σ -matrices with \sim -reduct having a (canonical) \sim -classical submatrix {and so being both consistent and truth-non-empty, for latter ones are so; cf. Remark 2.7(ii)(b)} (and carrier $3 \div 2$; cf. Subparagraph 2.2.1.2.1) are said to be (*[3-]canonical*(ly)) \sim -*super-classical*, in which case,

by (2.15), \sim is a subclassical negation for their logics {cf. Paragraph 2.3.2.1}, and so we have the “if” part of the following marking the framework of this subsection:

Theorem 5.7. *Let \mathcal{A} be a [no-more-than-three-valued] Σ -matrix. Then, \sim is a subclassical negation for the logic of \mathcal{A} iff \mathcal{A} is \sim -super-classical.*

Proof. [Assume \sim is a subclassical negation for the logic of \mathcal{A} . First, by (2.16) with $m = 1$ and $n = 0$, there is some $a \in D^{\mathcal{A}}$ such that $\sim^{\mathfrak{A}}a \notin D^{\mathcal{A}}$. Likewise, by (2.16) with $m = 0$ and $n = 1$, there is some $b \in (A \setminus D^{\mathcal{A}})$ such that $\sim^{\mathfrak{A}}b \in D^{\mathcal{A}}$, in which case $a \neq b$, and so $|A| \neq 1$. Then, if $|A| = 2$, we have $A = \{a, b\}$, in which case \mathcal{A} is \sim -classical, and so \sim -super-classical. Now, assume $|A| = 3$.

Claim 5.8. *Let \mathcal{A} be a three-valued Σ -matrix, $\bar{a} \in A^2$ and $i \in 2$. Suppose \sim is a subclassical negation for the logic of \mathcal{A} and, for each $j \in 2$, $(a_j \in D^{\mathcal{A}}) \Leftrightarrow (\sim^{\mathfrak{A}}a_j \notin D^{\mathcal{A}}) \Leftrightarrow (a_{1-j} \notin D^{\mathcal{A}})$. Then, either $\sim^{\mathfrak{A}}a_i = a_{1-i}$ or $\sim^{\mathfrak{A}}\sim^{\mathfrak{A}}a_i = a_i$.*

Proof. By contradiction. For suppose both $\sim^{\mathfrak{A}}a_i \neq a_{1-i}$ and $\sim^{\mathfrak{A}}\sim^{\mathfrak{A}}a_i \neq a_i$. Then, in case $a_i \in / \notin D^{\mathcal{A}}$, as $|A| = 3$, we have both $(D^{\mathcal{A}}/(A \setminus D^{\mathcal{A}})) = \{a_i\}$, in which case $\sim^{\mathfrak{A}}a_{1-i} = a_i$, and $((A \setminus D^{\mathcal{A}})/D^{\mathcal{A}}) = \{a_{1-i}, \sim^{\mathfrak{A}}a_i\}$, respectively. Consider the following exhaustive cases:

- $\sim^{\mathfrak{A}}\sim^{\mathfrak{A}}a_i = a_{1-i}$. Then, $\sim^{\mathfrak{A}}\sim^{\mathfrak{A}}\sim^{\mathfrak{A}}a_i = a_i$. This contradicts to (2.16) with $(n/m) = 0$ and $(m/n) = 3$, respectively.
- $\sim^{\mathfrak{A}}\sim^{\mathfrak{A}}a_i = \sim^{\mathfrak{A}}a_i$. Then, for each $c \in ((A \setminus D^{\mathcal{A}})/D^{\mathcal{A}})$, $\sim^{\mathfrak{A}}\sim^{\mathfrak{A}}\sim^{\mathfrak{A}}c = \sim^{\mathfrak{A}}a_i \notin / \in D^{\mathcal{A}}$. This contradicts to (2.16) with $(n/m) = 3$ and $(m/n) = 0$.

Thus, in any case, we come to a contradiction, as required. \square

Set $d_0 \triangleq a$ and $d_1 \triangleq b$. Consider the following complementary cases:

- for each $k \in 2$, $\sim^{\mathfrak{A}}d_k = d_{1-k}$. Then, $\{a, b\}$ forms a subalgebra of $\mathfrak{A}|\{\sim\}$, $(\mathcal{A}|\{\sim\})|\{a, b\}$ being a \sim -classical submatrix of $\mathcal{A}|\{\sim\}$, as required.
- for some $k \in 2$, $\sim^{\mathfrak{A}}d_k \neq d_{1-k}$, in which case, by Claim 5.8, $\sim^{\mathfrak{A}}\sim^{\mathfrak{A}}d_k = d_k$, so $\{d_k, \sim^{\mathfrak{A}}d_k\}$ forms a subalgebra of $\mathfrak{A}|\{\sim\}$, $(\mathcal{A}|\{\sim\})|\{d_k, \sim^{\mathfrak{A}}d_k\}$ being a \sim -classical submatrix of $\mathcal{A}|\{\sim\}$, as required. \square

The stipulation “no-more-than-three-valued” is essential in Theorem 5.7, by:

Example 5.9. Let $n \in \omega$ and \mathcal{A} any Σ -matrix with $A \triangleq (n \cup (2 \times 2))$, $D^{\mathcal{A}} \triangleq \{(1, 0), \langle 1, 1 \rangle\}$, $\sim^{\mathfrak{A}}\langle i, j \rangle \triangleq \langle 1 - i, (1 - i + j) \bmod 2 \rangle$, for all $i, j \in 2$, and $\sim^{\mathfrak{A}}k \triangleq \langle 1, 0 \rangle$, for all $k \in n$. Then, for any subalgebra \mathfrak{B} of $\mathfrak{A}|\{\sim\}$, we have $(2 \times 2) \subseteq \mathfrak{B}$, in which case $4 \leq |B|$, and so \mathcal{A} is not \sim -super-classical, for $4 \not\leq 2$. On the other hand, 2×2 forms a subalgebra of $\mathfrak{A}|\{\sim\}$, while $\mathfrak{B} \triangleq (\mathcal{A}|\{\sim\})|(2 \times 2)$ is \sim -negative, in which case $\theta^{\mathfrak{B}} \in \text{Con}(\mathfrak{B})$, and so $h \triangleq \chi^{\mathfrak{B}}$ is a surjective strict homomorphism from \mathfrak{B} onto the classically-canonical (in particular, two-valued) $\{\sim\}$ -matrix $\mathcal{C} \triangleq \langle h[\mathfrak{B}], \{1\} \rangle$, (in particular, by Remark 2.7(ii)(a), \mathcal{C} is \sim -classical, so, by (2.15), \sim is a subclassical negation for the logic of \mathcal{A}). \square

In general, given any three-valued \sim -super-classical Σ -matrix \mathcal{A} with \sim -classical submatrix \mathfrak{B} of its \sim -reduct, the bijection $e \triangleq (\chi^{\mathfrak{B}} \cup ((A \setminus B) \times \{\frac{1}{2}\})) : A \rightarrow (3 \div 2)$ is an isomorphism from \mathcal{A} onto the canonical \sim -super-classical Σ -matrix $\mathcal{C}_{[3]}(\mathcal{A}) \triangleq \langle e[\mathfrak{A}], e[D^{\mathcal{A}}] \rangle$, called the [3]-canonization of \mathcal{A} .

Throughout the rest of this subsection, unless otherwise specified, \mathcal{C} is supposed to be the logic of an arbitrary but fixed canonical \sim -super-classical Σ -matrix \mathcal{A} (that exhausts all uniform three-valued Σ -logics with subclassical negation \sim , in view of Theorem 5.7 and (2.15)), in which case this is false-singular iff it is not truth-singular iff $\mathbb{k}^{\mathcal{A}} \triangleq \chi^{\mathcal{A}}(\frac{1}{2}) = 1$, and so is false-/truth-singular, whenever it is \sim -paraconsistent/“both weakly \vee -disjunctive and (\vee, \sim) -paracomplete”, respectively,

in which case C is not \sim -classical, in view of Remark 2.7(i)(c). And what is more, any proper submatrix \mathcal{B} of \mathcal{A} is either \sim -classical or one-valued, in which case \mathcal{B} is simple, and so \mathcal{A} is simple iff it is hereditarily so. Also, \mathcal{A} is [weakly] \diamond -conjunctive/ \neg -disjunctive/ \neg -implicative iff C is so, in view of the following results:

Lemma 5.10. *Let \mathcal{B} be a Σ -matrix and C' the logic of \mathcal{B} . Suppose \mathcal{B} is [not] false-singular [as well as both no-more-than-three-valued and \sim -super-classical]. Then, the following are equivalent:*

- (i) C' is ∇ -disjunctive;
- (ii) \mathcal{B} is ∇ -disjunctive;
- (iii) (2.3) with $i = 0$, (2.4) and (2.5) [as well as (3.2)] are satisfied in C' .

Proof. First, (ii) \Rightarrow (i) is immediate. Next, assume (i) holds. Then, (2.3) with $i = 0$, (2.4) and (2.5) are immediate. [And what is more, once \mathcal{B} is not false-singular, it is both no-more-than-three-valued (and so truth-singular) and \sim -super-classical, in which case it is not \sim -paraconsistent, and so is C' . Then, by (i) and Lemma 3.10, (3.2) is satisfied in C' .] Thus, (iii) holds. Finally, assume (iii) holds. Consider any $a, b \in B$. Then, by (2.3) with $i = 0$ and (2.4), C' is weakly ∇ -disjunctive, and so is \mathcal{B} , in which case $(a \nabla^{\mathcal{B}} b) \in D^{\mathcal{B}}$, whenever either a or b is in $D^{\mathcal{B}}$. Now, assume $(\{a, b\} \cap D^{\mathcal{B}}) = \emptyset$. Then, in case $a = b$ (in particular, \mathcal{B} is false-singular), by (2.5), we get $D^{\mathcal{B}} \not\ni (a \nabla^{\mathcal{B}} a) = (a \nabla^{\mathcal{B}} b)$. [Otherwise, \mathcal{B} is not false-singular, in which case it is no-more-than-three-valued (in particular, truth-singular) and \sim -super-classical, while (3.2) is true in \mathcal{B} , and so, for some $c \in (B \setminus D^{\mathcal{B}}) = \{a, b\}$, it holds that $\sim^{\mathcal{B}} c \in D^{\mathcal{B}}$, while $\sim^{\mathcal{B}} \sim^{\mathcal{B}} c = c$. Let d be the unique element of $\{a, b\} \setminus \{c\}$, in which case $\{a, b\} = \{c, d\}$. Then, since $\sim^{\mathcal{B}} c \in D^{\mathcal{B}}$, we conclude that $(c \nabla^{\mathcal{B}} d) = (\sim^{\mathcal{B}} \sim^{\mathcal{B}} c \nabla^{\mathcal{B}} d) \notin D^{\mathcal{B}}$, for, otherwise, by (2.3) with $i = 0$ and (3.2), we would get $d \in D^{\mathcal{B}}$. Hence, by (2.4), we eventually get $(a \nabla^{\mathcal{B}} b) \notin D^{\mathcal{B}}$.] \square

Corollary 5.11. *[Providing \mathcal{A} is false-singular (in particular, \sim -paraconsistent)] \mathcal{A} is \sqsupset -implicative iff C is [weakly] so.*

Proof. By (2.7), (2.8), (2.10), Theorem 3.5(ii) and Lemma[s 3.13 and] 5.10. \square

Remark 5.12. \mathcal{A} is not \sim -negative iff it has unitary equality determinant $\{x_0, \sim x_0\}$.

Next, \mathcal{A} is said to be (\sim -)involutive, provided $\sim^{\mathfrak{A}} \frac{1}{2} = \frac{1}{2}$, that is, the Σ -identity $\sim \sim x_0 \approx x_0$ is true in \mathfrak{A} , in which case \mathcal{A} is not \sim -negative. Further, \mathcal{A} is said to be [extra-]classically-hereditary, provided $[A \setminus] 2$ forms a subalgebra of \mathfrak{A} [in which case \mathcal{A} is involutive]. Then, \mathcal{A} , being classically-hereditary, is said to be genuinely \diamond -conjunctively/ \neg -disjunctively/ \neg -implicatively so, whenever $\mathcal{A}[2]$ is “genuinely \sim -classical” \diamond -conjunctive/ \neg -disjunctive/ \neg -implicative, respectively. Finally, \mathcal{A} is said to be classically-valued, provided, for all $\varsigma \in \Sigma$, $(\text{img } \varsigma^{\mathfrak{A}}) \subseteq 2$, in which case \mathcal{A} is [not extra-]classically-hereditary as well as not involutive.

5.2.1.1. Examples.

5.2.1.1.1. Kleene-style logics. Let $\Sigma \triangleq \Sigma_{\sim, +[01]}$ and \mathcal{A} both involutive and truth-/false-singular with $(\mathfrak{A}[\Sigma_{+[01]}) \triangleq \mathfrak{D}_{3[01]}$. Then, \mathcal{A} is both \wedge -conjunctive, \vee -disjunctive and non- \sim -negative, in which case it is (\vee, \sim) -paracomplete/ \sim -paraconsistent, and so, by Remark 2.7(i)(c), C is not \sim -classical, as well as both classically-hereditary and [not] extra-classically-hereditary, while \mathfrak{A} is a distributive (\wedge, \vee) -lattice with zero 0 and unit 1, whereas C is [the bounded version|expansion $K_{3,01}/LP_{01}$ of] “Kleene’s three-valued logic”/“the logic of paradox” K_3/LP [4]/[9].

5.2.1.1.2. Gödel-style logics. Let $\Sigma \triangleq \Sigma_{\sim, +, 01}^{\supset}$ and \mathcal{A} [not] truth-singular as well as neither \sim -negative nor involutive with $(\mathfrak{A}[\Sigma_{+, 01}]) \triangleq \mathfrak{D}_{3,01}$ (in which case $\sim^{\mathfrak{A}}$ is the [dual] pseudo-complement operation)/“as well as $\supset^{\mathfrak{A}}$ being the [dual] relative pseudo-complement operation”. Then, \mathcal{A} is both \wedge -conjunctive, \vee -disjunctive and

[not] (\vee, \sim) -paracomplete as well as [not] non- \sim -paraconsistent, and so, by Remark 2.7(i)(c), C is not \sim -classical, while \mathcal{A} is classically-hereditary but not extra-classically-hereditary, whereas C is [the (\sim) -paraconsistent counterpart PG_3^* of] “the implication-less fragment G_3^* of” / Gödel’s three-valued logic G_3 [2].

5.2.1.1.3. Hałkowska-Zajac’ logic. Let $\Sigma \triangleq \Sigma_{\sim,+}$ and \mathcal{A} both false-singular and involutive with \mathfrak{A} being the distributive (\wedge, \vee) -lattice with zero $\frac{1}{2}$ and unit 1. Then, \mathcal{A} is \sim -paraconsistent (in particular, C is not \sim -classical; cf. Remark 2.7(i)(c)) as well as both classically- and extra-classically-hereditary but weakly neither \wedge -conjunctive nor \vee -disjunctive, C being the logic HZ [3]. On the other hand, since the identity $\sim x_0 \approx x_0$ is true in \mathfrak{A} , \mathfrak{A} is a distributive (\vee, \wedge) -lattice (cf. Remark 2.7(i)(a) for definition of these secondary binary connectives) with zero $\sim^{\mathfrak{A}}1 = 0$ and unit $\sim^{\mathfrak{A}}\frac{1}{2} = \frac{1}{2}$. Then, \mathcal{A} is both \vee -conjunctive and \wedge -disjunctive.

5.2.1.1.4. Sette-style logics. Let $\Sigma \triangleq \Sigma_{\supset}$ and \mathcal{A} classically-valued, non- \sim -negative, \supset -implicative (in particular, \uplus_{\supset} -disjunctive) and [not] false-singular. Then, \mathcal{A} is [not] \sim -paraconsistent as well as [not] non- (\uplus_{\supset}, \sim) -paracomplete, and so, by Remark 2.7(i)(c), C , being [the intuitionistic/ (\uplus_{\supset}, \sim) -paracomplete counterpart IP^1 of] P^1 [21], is not \sim -classical.

5.2.2. Non-classical U3VLSN. C need not be non- \sim -classical, by:

Example 5.13. Let $\Sigma \triangleq \Sigma_{\sim,+,\mathfrak{01}}$ and $(\mathcal{B}/\mathcal{D})|\mathcal{E}$ the \wedge -conjunctive \vee -disjunctive canonical “ \sim -negative false-/truth-singular \sim -super-classical” $|\sim$ -classical Σ -matrix, with $((\mathfrak{B}/\mathfrak{D})|\mathfrak{E})|\Sigma_{\sim,+,\mathfrak{01}} \triangleq \mathfrak{D}_{3|2,\mathfrak{01}}$ (cf. Subparagraph 2.2.1.2.1), respectively. Then, $(\mathcal{B}/\mathcal{D})|\mathcal{E}$ is \supset -implicative, in view of Remark 2.7(i)(b). And what is more, $\chi^{\mathcal{B}/\mathcal{D}} \in \text{hom}_{\mathfrak{S}}^{\mathfrak{S}}(\mathcal{B}/\mathcal{D}, \mathcal{E})$. Therefore, by (2.15), \mathcal{B}/\mathcal{D} define the same \sim -classical Σ -logic of \mathcal{E} . On the other hand, \mathfrak{B} and \mathfrak{D} are non-isomorphic (in particular, \mathcal{B} and \mathcal{D} are so), because the Σ -identity $(x_0 \wedge \sim x_0) \approx \perp$, being true in \mathfrak{B} , is not so in \mathfrak{D} under $[x_0/\frac{1}{2}]$. Moreover, $h \triangleq (\chi^{\mathcal{B}/\mathcal{D}} \circ \Delta_2)$ is a non-diagonal (for $h(\frac{1}{2}) = (1/0) \neq \frac{1}{2}$) strict homomorphism from \mathcal{B}/\mathcal{D} to itself, so this does not have a unitary equality determinant, in view of Theorem 3.3. \square

On the other hand, \sim -classical Σ -logics are self-extensional, in view of Example 4.2. This makes the purely algebraic criterion of the classicism of U3VLSN to be obtained here especially acute.

Let $\Delta_2^+ \triangleq \Delta_2 \in 2^2$ and $\Delta_2^- \triangleq (A^2 \setminus \Delta_2) \in 2^2$.

Lemma 5.14 (Key 3-valued Lemma). *Let \mathcal{B} be a canonical \sim -super-classical Σ -matrix, \mathcal{D} a submatrix of \mathcal{A} and $h \in \text{hom}(\mathfrak{D}, \mathfrak{B})$. Then, providing \mathcal{A} is involutive, whenever both \mathcal{B} is so and $\frac{1}{2} \in (\text{img } h)$ (in particular, either $\mathfrak{A} = \mathfrak{B}$ or $\text{hom}(\mathfrak{B}|\text{img } h, \mathfrak{A}) \neq \emptyset$), the following hold:*

- (i) *providing h is not singular {cf. p. 2}, $2 \subseteq D$, while $h[2] = 2$, in which case $h|_2$ is bijective, and so belongs to $\{\Delta_2^+, \Delta_2^-\}$;*
- (ii) *providing $h \not\subseteq \Delta_2^-$ [in particular, $h \in \text{hom}(\mathcal{D}, \mathcal{B})$] is injective, it is diagonal.*

In particular, the following hold:

- (a) *any partial automorphism {cf. Subsubsection 3.1.1} of \mathcal{A} is diagonal;*
- (b) *any isomorphism from \mathcal{A} onto \mathcal{B} is diagonal, in which case $\mathcal{A} = \mathcal{B}$, and so \mathcal{A} and \mathcal{B} are equal, whenever they are isomorphic.*

Proof. First, note that the carrier of any subalgebra of $(\mathfrak{A}|\mathfrak{B})|\{\sim\}$ (in particular, $D|(\text{img } h)$) belongs to $\{A|B, 2, \{\frac{1}{2}\}\}$. And what is more, for each $a \in (A|B)$, we have $(\sim^{\mathfrak{A}|\mathfrak{B}} a = a) \Rightarrow (a = \frac{1}{2})$. In particular, for any $g \in \text{hom}(\mathfrak{D}|(\mathfrak{B}|\text{img } h), \mathfrak{B}|\mathfrak{A})$ with $\frac{1}{2} \in (\text{dom } g)$, providing $\sim^{\mathfrak{A}|\mathfrak{B}}\frac{1}{2} = \frac{1}{2}$, we have $\sim^{\mathfrak{B}|\mathfrak{A}}g(\frac{1}{2}) = g(\frac{1}{2})$, in which case we get $g(\frac{1}{2}) = \frac{1}{2}$, and so $\sim^{\mathfrak{B}|\mathfrak{A}}\frac{1}{2} = \frac{1}{2}$. While proving (i,ii), assume $(\sim^{\mathfrak{B}}\frac{1}{2} = \frac{1}{2}) \Rightarrow (\sim^{\mathfrak{A}}\frac{1}{2} = \frac{1}{2})$, whenever $\frac{1}{2} \in (\text{img } h)$.

- (i) Assume h is not singular, in which case $1 < |\text{img } h| \leq |D|$, and so $D \supseteq 2 \subseteq (\text{img } h)$. Then, as 2 forms a subalgebra of $\mathfrak{A}\{\sim\}$, $h[2]$ forms a no-more-than-two-element subalgebra of $\mathfrak{B}\{\sim\}$, in which case $h[2] \in \{2, \{\frac{1}{2}\}\}$, and so $h[2] = 2$, for, otherwise, we would have both $(\text{img } h) = h[D] \supseteq h[2] = \{\frac{1}{2}\} \ni \frac{1}{2}$ and $\sim^{\mathfrak{B}} \frac{1}{2} = \frac{1}{2}$, in which case we would get $\sim^{\mathfrak{A}} \frac{1}{2} = \frac{1}{2}$ as well as, since $|\text{img } h| \neq 1$, both $\frac{1}{2} \in D = (\text{dom } h)$ and $h(\frac{1}{2}) \in 2$, and so would eventually get $2 \ni h(\frac{1}{2}) = \frac{1}{2}$.
- (ii) Assume h is injective, while $\{h \in \text{hom}(\mathcal{D}, \mathcal{B})$, in which case $\Delta_2^- \ni \langle 1, 0 \rangle \notin h$, for $\langle 1|0 \rangle \in \notin D^{\mathcal{A}\mathcal{B}}$, and so $\Delta_2^- \not\subseteq h$. Then, h is a bijection from D onto $\text{img } h$. Therefore, in case h is singular, we have $(\text{img } h) = \{\frac{1}{2}\} = D$, and so $h = \{\langle \frac{1}{2}, \frac{1}{2} \rangle\}$ is diagonal. Otherwise, by (i), $2 \subseteq D$, while $(h[2]) \subseteq h$ is diagonal. In particular, $h = (h[2])$ is diagonal, whenever $D = 2$. Otherwise, $D = A$, while $\frac{1}{2} \notin 2$, in which case, by the injectivity of h , we have $h(\frac{1}{2}) \notin h[2] = 2$, and so we get $h(\frac{1}{2}) = \frac{1}{2}$ (in particular, h is diagonal).

Then, (a/b) is by (ii) with $(\mathcal{B}/\mathcal{D}) = \mathcal{A}$ and and /bijective $h \in \text{hom}(\mathcal{D}, \mathcal{B})$ /“as well as $h^{-1} \in \text{hom}(\mathfrak{B}, \mathfrak{A})$ ”. \square

Corollary 5.15. *The following are equivalent:*

- (i) \mathcal{A} has no [unitary] equality determinant;
- (ii) \mathcal{A} is a strictly (surjectively) homomorphic counter-image of a \sim -classical Σ -matrix;
- (iii) \mathcal{A} is not {hereditarily} simple;
- (iv) $\theta^{\mathcal{A}} \in \text{Con}(\mathfrak{A})$ (in which case $\chi^{\mathcal{A}}$ is a strict surjective homomorphism from \mathcal{A} onto $\mathcal{C}_{\mathcal{A}} \triangleq \langle \chi^{\mathcal{A}}[\mathcal{A}], \{1\} \rangle$, being, in its turn, canonically \sim -classical).

Proof. First, (i) \Leftrightarrow (iii) is by Lemmas 3.1, 5.14(a) and Theorem 3.3.

Next, (ii) \Rightarrow (iii) is by Remark 2.5(i,ii), for $|\mathcal{A}| = 3 \not\leq 2$.

Further, (iii) \Rightarrow “ $\theta^{\mathcal{A}} \in \text{Con}(\mathfrak{A})$ ” is by the fact $\text{img}[\theta^{\mathcal{A}} \setminus \Delta_{\mathcal{A}}] = \{\{\frac{1}{2}, \mathbb{k}^{\mathcal{A}}\}\}$ is a singleton.

Finally, assume $\theta^{\mathcal{A}} \in \text{Con}(\mathfrak{A})$, in which case $h \triangleq \chi^{\mathcal{A}}$ is a strict surjective homomorphism from \mathcal{A} onto the classically-canonical (in particular, two-valued) Σ -matrix $\mathcal{C}_{\mathcal{A}}$, and so $h[2]$, being diagonal, is a strict surjective homomorphism from the \sim -negative Σ -matrix $(\mathcal{A}\{\sim\})[2]$ onto $\mathcal{C}_{\mathcal{A}}\{\sim\}$. Then, by Remark 2.7(ii)(a), $\mathcal{C}_{\mathcal{A}}\{\sim\}$ is \sim -negative, and so is $\mathcal{C}_{\mathcal{A}}$, in which case this is canonically \sim -classical. Thus, the optional part of (iv) holds, and so does (ii). \square

Lemma 5.16. *Let \mathcal{B} be a canonically \sim -[super-]classical Σ -matrix, I a finite set, $\bar{\mathcal{C}} \in \mathbf{S}_*(\mathcal{B})^I$ and \mathcal{D} a subdirect product of it. Then, the following hold:*

- (i) providing [\mathcal{B} is weakly conjunctive, whenever it is \sim -paraconsistent, and] \mathcal{D} is truth-non-empty [unless \mathcal{B} is \sim -paraconsistent], $(I \times \{j\}) \in \mathcal{D}$, for each $j \in 2$;
- (ii) providing $I \neq \emptyset$ (in particular, \mathcal{D} is consistent), while, for some $j \in 2$, $(I \times \{j\}) \in \mathcal{D}$, for each $\Sigma' \subseteq \Sigma$, $\{\langle b, I \times \{b \rangle \mid b = \varphi^{\mathfrak{B}}(0, 1), \varphi \in (\text{Var}_2[\cup \text{Fm}_{\Sigma'}^2])\}\}$ is an embedding of [the submatrix of] $\mathcal{B}\{\Sigma'\}$ [generated by 2] into $\mathcal{D}\{\Sigma'\}$.

Proof. (i) We use Remark 2.7(i)(c) tacitly. Consider the following cases:

- \mathcal{B} is \sim -paraconsistent, in which case it is both false-singular and weakly conjunctive, and so, by Lemma 3.12, $(I \times \{0\}) \in \mathcal{D}$.
- \mathcal{B} is not \sim -paraconsistent, in which case \mathcal{D} is truth-non-empty, and so there is some $d \in \mathcal{D}^{\mathcal{D}} \in \emptyset_{\infty \setminus 1}(\mathcal{D})$. Consider any $i \in I$ and the following complementary subcases:
 - \mathcal{B} is truth-singular, in which case $\pi_i(d) = 1$, and so $\pi_i(\sim^{\mathfrak{D}} d) = \sim^{\mathfrak{B}} \pi_i(d) = 0$.

- \mathcal{B} is not truth-singular, in which case it is false-singular, and so, as $\pi_i(\sim^{\mathfrak{D}}d) = \sim^{\mathfrak{B}}\pi_i(d) \notin D^{\mathfrak{B}}$, for, otherwise, (2.11) would not be true in \mathcal{B} under $[x_0/\pi_i(d), x_1/0]$, we have $\pi_i(\sim^{\mathfrak{D}}d) = 0$.

Thus, anyway, $\pi_i(\sim^{\mathfrak{D}}d) = 0$, so $D \ni \sim^{\mathfrak{D}}d = (I \times \{0\})$.

In this way, in any case, $(I \times \{0\}) \in D$, and so $D \ni \sim^{\mathfrak{D}}(I \times \{0\}) = (I \times \{1\})$.

- (ii) In that case, $D \ni \sim^{\mathfrak{D}}(I \times \{j\}) = (I \times \{1-j\})$, and so the fact that $2 = \{j, 1-j\}$ completes the argument. \square

Let $h_{+/2} : 2^2 \rightarrow A, \langle i, j \rangle \mapsto \frac{i+j}{2}$.

Theorem 5.17. *C is \sim -classical iff either of the following holds:*

- (i) $\theta^A \in \text{Con}(\mathfrak{A})$ (in which case $C_{\mathcal{A}} \triangleq \langle \chi^{\mathcal{A}}[\mathfrak{A}], \{1\} \rangle$ is a canonical \sim -classical Σ -matrix, being a strictly surjectively homomorphic image of \mathcal{A} , and so defines C);
- (ii) \mathcal{A} is truth-singular, while 2 forms a subalgebra of \mathfrak{A} , whereas $h_{+/2} \in \text{hom}((\mathfrak{A}|2)^2, \mathfrak{A})$ (in which case $h_{+/2} \in \text{hom}_{\mathfrak{S}}^{\mathfrak{S}}((\mathcal{A}|2)^2, \mathcal{A})$, and so $\mathcal{A}|2$ is a canonical \sim -classical Σ -matrix defining C).

Proof. Both the “if” part and the optional “in which case” one are immediate with using (2.15) and Corollary 5.15(iv). Conversely, assume C is \sim -classical, in which case, by (2.15), C is defined by a canonical \sim -classical (in particular, both simple and having no proper submatrix) Σ -matrix \mathcal{B} , while $\theta^A \notin \text{Con}(\mathfrak{A})$, in which case, by Corollary 5.15(iii) \Rightarrow (iv), \mathcal{A} is hereditarily simple, and so, by Lemma 3.7 with $M = \{\mathcal{B}|\mathcal{A}\}$, there is some finite set $I|J$, some $\overline{C|\mathcal{D}} \in \mathbf{S}_*(\mathcal{B}|\mathcal{A})^{I|J}$ some subirect product $\mathcal{E}|\mathcal{F}$ of it and some $(h|g) \in \text{hom}_{\mathfrak{S}}^{\mathfrak{S}}(\mathcal{E}|\mathcal{F}, \mathcal{A}|\mathcal{B})$ (in particular, $\mathfrak{A}|\mathfrak{B}$ belongs to the variety generated by $\mathfrak{B}|\mathfrak{A}$). Then, \mathcal{A} is truth-singular (in particular, non- \sim -paraconsistent), for \mathcal{B} is so, while truth-singularity is clearly preserved under \mathbf{P} as well as under both \mathbf{S} and \mathbf{H} , in view of Remark 2.7(ii)(c). And what is more, by Remark 2.7(ii)(b), $\mathcal{E}|\mathcal{F}$ is both truth-non-empty and consistent, for $\mathcal{A}|\mathcal{B}$ is so. Then, by Lemma 5.16(i) with $j = (0[+1])$, $(E|F) \ni (a|b)[\cdot] \triangleq ((I|J) \times \{j\})$. Let \mathcal{G} be the submatrix of \mathcal{A} generated by 2 , in which case it is simple, for \mathcal{A} is hereditarily so, and so, by Remark 2.5(ii) and Lemma 5.16(ii), $e \circ g$, where e is an embedding of \mathcal{G} into \mathcal{F} , is an embedding of \mathcal{G} into \mathcal{B} (in particular, is an isomorphism from \mathcal{G} onto \mathcal{B} , for this has no proper submatrix). Thus, $|G| = |B| = |2| = 2$, in which case $G \supseteq 2$ is equal to 2 , and so $2 = G$ forms a subalgebra of \mathfrak{A} , while $(\mathcal{A}|2) = \mathcal{G}$ is canonically \sim -classical and isomorphic (and so equal) to \mathcal{B} . And what is more, by the truth-singularity of \mathcal{A} , $h(a') = 1$, for $a' \in D^{\mathcal{E}}$, in which case $h(a) = h(\sim^{\mathcal{E}}a') = \sim^{\mathfrak{A}}1 = 0$, and so there is some $c \in (E \setminus \{a, a'\})$ such that $h(c) = \frac{1}{2}$. Then, $I \neq K \triangleq \{i \in I \mid \pi_i(c) = 1\} \neq \emptyset$, in which case $f \triangleq \{\langle \langle k, l \rangle, (K \times \{k\}) \cup ((I \setminus K) \times \{l\}) \rangle \mid k, l \in 2\}$ is an embedding of \mathcal{B}^2 into \mathcal{E} , and so $(f \circ h) \in \text{hom}(\mathfrak{B}^2, \mathfrak{A})$. Clearly, $f(\langle 1, 1 \rangle) = a'$, $f(\langle 0, 0 \rangle) = a$, $f(\langle 1, 0 \rangle) = c$, and so $f(\langle 0, 1 \rangle) = f(\sim^{\mathfrak{B}^2}\langle 1, 0 \rangle) = \sim^{\mathcal{E}}c$. Furthermore, the Σ -identity $\sim\sim x_0 \approx x_0$, being true in \mathfrak{B} , is so in \mathfrak{A} , for this belongs to the variety generated by \mathfrak{B} , in which case $\sim^{\mathfrak{A}}\sim^{\mathfrak{A}}\frac{1}{2} = \frac{1}{2} \notin 2$, and so $\sim^{\mathfrak{A}}\frac{1}{2} = \frac{1}{2}$. Thus, $(f \circ h) = h_{+/2}$, as required. \square

Corollary 5.18. *[Providing \mathcal{A} is either false-singular or \diamond -conjunctive/-disjunctive] C is \sim -classical iff $\theta^A \in \text{Con}(\mathfrak{A})$, (i.e., \mathcal{A} “has no {unitary} equality determinant” | “is not (hereditarily) simple” | “is not a strictly [surjectively] homomorphic counter-image of a \sim -classical Σ -matrix; cf. Corollary 5.15).*

Proof. By Remark 2.7(i)(a),(ii)(a), Lemma 5.10 and Theorem 5.17, because, for any \vee -disjunctive consistent truth-non-empty Σ -matrix \mathcal{B} , \mathcal{B}^2 is not \vee -disjunctive, since, for any $a \in D^{\mathcal{B}} \neq \emptyset$ and any $b \in (B \setminus D^{\mathcal{B}}) \neq \emptyset$, $(\langle a, b \rangle \vee^{\mathcal{B}^2} \langle b, a \rangle) \in D^{\mathcal{B}^2}$, while $\{\langle a, b \rangle, \langle b, a \rangle\} \subseteq (B^2 \setminus D^{\mathcal{B}^2})$. \square

In view of Example 1 of [14], this implies that U3VLSN are covered by the universal approach elaborated therein. On the other hand, the optional stipulation in the formulation of Corollary 5.18 can be neither omitted nor even weakened, because \mathcal{A} may be simple as well as weakly both conjunctive and disjunctive but \sim -classically-defining, in view of:

Example 5.19. Let $\Sigma \triangleq \Sigma_{\sim,01}$ and \mathcal{A} truth-singular with $\sim^{\mathfrak{A}} \frac{1}{2} \triangleq \frac{1}{2}$ and $(\perp/\top)^{\mathfrak{A}} \triangleq (0/1)$. Then, \mathcal{A} is weakly both \perp -conjunctive and \top -disjunctive as well as non- \sim -negative, in which case $\{x_0, \sim x_0\}$ is an equality determinant for \mathcal{A} , and so this simple (cf. Lemma 3.1). On the other hand, 2 forms a subalgebra of \mathfrak{A} , while $h_{+/\frac{1}{2}} \in \text{hom}((\mathfrak{A}|2)^2, \mathfrak{A})$, in which case, by Theorem 5.17, C is defined by the \sim -classical Σ -matrix $\mathcal{A}|2$, and so is \sim -classical. \square

Likewise, the item (ii) cannot be omitted in the formulation of Theorem 5.17.

Finally, Theorem 5.5(i) \Rightarrow (v) makes the next paragraph equally acute.

5.2.2.1. Classical models and extensions.

Lemma 5.20. *Let \mathcal{B} and \mathcal{D} be Σ -matrices. Suppose \mathcal{B} is weakly \sim -negative, while \mathcal{D} is consistent but not \sim -paraconsistent (in particular, \sim -negative; cf. Remark 2.7(i)(c)). Then, any $h \in \text{hom}(\mathcal{B}, \mathcal{D})$ is strict.*

Proof. Take any $d \in (D \setminus D^{\mathcal{D}}) \neq \emptyset$. If, for any $b \in (B \setminus D^{\mathcal{B}})$, $h(b)$ was in $D^{\mathcal{D}}$, then $h(\sim^{\mathfrak{B}} b) = \sim^{\mathcal{D}} h(b)$ would not belong to $D^{\mathcal{D}}$, for, otherwise, (2.11) would not be true in \mathcal{D} under $[x_0/h(b), x_1/d]$, in which case $\sim^{\mathfrak{B}} b$ would not belong to $D^{\mathcal{B}}$, for $h[D^{\mathcal{B}}] \subseteq D^{\mathcal{D}}$, and so \mathcal{B} would not be weakly \sim -negative. \square

Corollary 5.21. *Let \mathcal{B} be a [classically hereditary $\{\vee$ -disjunctive \sim -paraconsistent}] canonically \sim -[super-]classical Σ -matrix, C' the logic of \mathcal{B} and \mathcal{C} a (canonically) \sim -classical model of C' . Then, $\mathcal{B}[|2]$ is a canonical \sim -classical model of C' , isomorphic (and so equal) to \mathcal{C} , in which case it defines a unique \sim -classical extension of C' [relatively axiomatized by (3.2)].*

Proof. In that case, both $\mathcal{C} \in \text{Mod}(C')$ and $\mathcal{B}[|2]$ are finite, simple and \sim -negative as well as have no proper submatrices. Then, by Remark 2.5(ii) and Lemma 3.7, there are some finite set I , some $\bar{\mathcal{C}} \in \mathbf{S}_*(\mathcal{B})^I$, some subdirect product \mathcal{D} of it and some $g \in \text{hom}_{\mathbb{S}}^{\mathbb{S}}(\mathcal{D}, \mathcal{C})$, in which case \mathcal{D} is non-one-valued, for \mathcal{C} is so, and so there is some $d \in D$ distinct from $I \times \{0[+\frac{1}{2}]\}$. Then, there is some $i \in I$ such that $\pi_i(d) \neq (0[+\frac{1}{2}])$ [in particular, $\pi_i(d) \in 2$], in which case, by Remark 2.7(ii)(a), the submatrix \mathcal{E} of \mathcal{D} generated by $\{d\}$ is \sim -negative, for \mathcal{C} is so, while $f \triangleq (g|E) \in \text{hom}_{\mathbb{S}}^{\mathbb{S}}(\mathcal{E}, \mathcal{C})$, for \mathcal{C} has no proper submatrix, whereas [since $\pi[E] \subseteq 2$, for \mathcal{B} is classically hereditary] $e \triangleq (\pi_i|E) \in \text{hom}(\mathcal{E}, \mathcal{B}[|2])$ surjective, for $\mathcal{B}[|2]$ has no proper submatrix, as well as strict, in view of Lemma 5.20, and so, by Remark 2.5(ii) and Lemma 3.6 with $M = \{\mathcal{C}\}$, \mathcal{C} is isomorphic to $\mathcal{B}[|2]$. Then, (2.15) [as well as Remark 2.7(i)(c) and Theorem 3.11] complete the argument. \square

Theorem 5.22. *Suppose C is not \sim -classical. Then, the following hold:*

- (i) *for any \sim -classical $\mathcal{B} \in \text{Mod}(C)$, either of the following hold:*
 - (a) *if \mathcal{A} is classically hereditary, then $(\mathcal{A}|2) \in \text{Mod}(C)$ is both canonically \sim -classical and is isomorphic to \mathcal{B} , and so defines a unique \sim -classical extension of C ;*
 - (b) *if \mathcal{A} is not classically hereditary, then $L_4 \triangleq (A^2 \setminus (2^2 \cup \Delta_A))$ forms a subalgebra of \mathfrak{A}^2 , while $\theta^{A^2}|L_4 \in \text{Con}(\mathfrak{A}^2|L_4)$, in which case $\langle \chi^{A^2}|L_4[\mathfrak{A}^2|L_4], \{1\} \rangle \in \text{Mod}(C)$ is both canonically \sim -classical and isomorphic to \mathcal{B} , and so defines a unique \sim -classical extension of C , whereas \mathcal{B} is not disjunctive, as well as C is \sim -paraconsistent but is not weakly conjunctive;*

- (ii) [Providing C is not \sim -classical] C is [genuinely] \sim -subclassical iff] \mathcal{A} is [genuinely/conjunctively/disjunctively/implicatively] classically hereditary;
- (iii) [Providing C is not \sim -classical but is (either disjunctive|“weakly conjunctive/implicative” or) non- \sim -paraconsistent] C is \sim -subclassical iff] \mathcal{A} is classically hereditary.

Proof. In that case, by Corollary 5.18, \mathcal{A} is hereditarily simple. Consider any \sim -classical $\mathcal{B} \in \text{Mod}(C)$, in which case it is both finite and simple as well as has no proper submatrix. And what is more, taking (2.15) and Remark 2.7(ii)(a) into account, without loss of generality, one can suppose that \mathcal{B} is canonically \sim -classical, in which case $D^{\mathcal{B}} = \{1\}$. Assume \mathcal{A} is not classically hereditary, in which case it is generated by 2, and so there is some $\varphi \in \text{Fm}_{\Sigma}^2$ such that $\varphi^{\mathfrak{A}}(1, 0) = \frac{1}{2}$. Then, by Lemma 3.7, there are some finite set I , some $\bar{c} \in \mathbf{S}_*(\mathcal{A})^I$, some subdirect product \mathcal{D} of it and some $h \in \text{hom}_{\Sigma}^{\mathcal{S}}(\mathcal{D}, \mathcal{B})$, in which case, by Remark 2.7(ii)(a), \mathcal{D} is \sim -negative (in particular, both consistent and truth-non-empty), for \mathcal{B} is so, and so is any Σ -matrix embeddable into \mathcal{D} . Therefore,

$$(5.3) \quad (I \times \{j\}) \notin D,$$

for all $j \in 2$, because, otherwise, by Lemma 5.16(ii), there would be some embedding e of \mathcal{A} into \mathcal{D} , in which case, by Remark 2.5(ii), $e \circ h$ would be an embedding of \mathcal{A} into \mathcal{B} , and so we would get $3 = |A| \leq |B| = 2$. Hence, by (5.3) and Lemma 5.16(i), \mathcal{A} is not weakly conjunctive but is \sim -paraconsistent, in which case $\{\frac{1}{2}, \sim^{\mathfrak{A}}\frac{1}{2}\} \subseteq D^{\mathcal{A}}$ (in particular, $D^{\mathcal{A}} = \{\frac{1}{2}, 1\}$), and so

$$(5.4) \quad (I \times \{\frac{1}{2}\}) \notin D,$$

for \mathcal{D} is \sim -negative. Take any $a \in D^{\mathcal{D}} \neq \emptyset$, in which case $a \in \{\frac{1}{2}, 1\}^I$, and so, by (5.3) with $j = 1$ and (5.4), $I \neq J \triangleq \{i \in I \mid \pi_i(a) = 1\} \neq \emptyset$. Let \mathcal{E} be the submatrix of \mathcal{A}^2 generated by $\{\langle 1, \frac{1}{2} \rangle\}$. Given any $b, c \in A$, set $(b \wp c) \triangleq ((J \times \{b\}) \cup ((I \setminus J) \times \{c\})) \in A^I$, in which case $a = (1 \wp \frac{1}{2})$, and so $g \triangleq \{\langle \langle b, c \rangle, (b \wp c) \rangle \mid \langle b, c \rangle \in E\}$ is an embedding of \mathcal{E} into \mathcal{D} . Then, \mathcal{E} is \sim -negative, in which case E is disjoint with $2^2 \setminus \Delta_2$, and so, by (5.3) and (5.4), $E \subseteq L_4$. Therefore, $\sim^{\mathfrak{A}}\frac{1}{2} = \frac{1}{2}$, for, otherwise, $E \ni \langle 1, \frac{1}{2} \rangle$ would contain $\sim^{\mathfrak{A}^2}\langle 1, \frac{1}{2} \rangle = \langle 0, \sim^{\mathfrak{A}}\frac{1}{2} \rangle \in 2^2$. Hence, $E \ni \sim^{\mathfrak{A}^2}\langle 1, \frac{1}{2} \rangle = \langle 0, \frac{1}{2} \rangle$. Let $\psi \triangleq (\varphi[x_1/\sim x_0]) \in \text{Fm}_{\Sigma}^1$, in which case $\psi^{\mathfrak{A}}(1) = \varphi^{\mathfrak{A}}(1, 0) = \frac{1}{2}$, and so $\psi^{\mathfrak{A}}(\frac{1}{2}) \in 2$, for, otherwise, $E \ni \langle 1, \frac{1}{2} \rangle$ would contain $\psi^{\mathfrak{A}^2}(\langle 1, \frac{1}{2} \rangle) = \langle \frac{1}{2}, \frac{1}{2} \rangle \in \Delta_A$. Thus, $\{\langle \frac{1}{2}, 1 \rangle, \langle \frac{1}{2}, 0 \rangle\} = \{\psi^{\mathfrak{A}^2}(\langle 1, \frac{1}{2} \rangle), \sim^{\mathfrak{A}^2}\psi^{\mathfrak{A}^2}(\langle 1, \frac{1}{2} \rangle)\} \subseteq E$, for $E \ni \langle 1, \frac{1}{2} \rangle$ forms a subalgebra of \mathfrak{A}^2 , in which case $E = L_4$, and so L_4 forms a subalgebra of \mathfrak{A}^2 . Then, $f \triangleq (g \circ h) \in \text{hom}_{\Sigma}^{\mathcal{S}}(\mathcal{E}, \mathcal{B})$ (in particular, $\mathfrak{B} = f[\mathfrak{E}]$), for \mathcal{B} has no proper submatrix, in which case $f = \chi^{\mathcal{E}}$, for \mathcal{B} is classically-canonical, and so $\theta^{\mathcal{E}} = (\ker f) \in \text{Con}(\mathfrak{E})$. Finally, we prove that \mathcal{B} is not disjunctive, by contradiction. For suppose \mathcal{B} is \vee -disjunctive, in which case, by Remark 2.7(ii)(a), \mathcal{E} is so, and so, as $\langle 1, \frac{1}{2} \rangle \in D^{\mathcal{E}}$, while $\langle \frac{1}{2}, 0 \rangle \in L_4 = E$ we have $\{\langle 1, \frac{1}{2} \rangle \vee^{\mathfrak{E}} \langle \frac{1}{2}, 0 \rangle, \langle \frac{1}{2}, 0 \rangle \vee^{\mathfrak{E}} \langle 1, \frac{1}{2} \rangle\} \subseteq D^{\mathcal{E}}$. Then, $(\frac{1}{2} \vee^{\mathfrak{A}} 0) \in D^{\mathcal{A}} \ni (0 \vee^{\mathfrak{A}} \frac{1}{2})$, in which case $(\langle 0, \frac{1}{2} \rangle \vee^{\mathfrak{E}} \langle \frac{1}{2}, 0 \rangle) \in D^{\mathcal{E}}$, and so $(D^{\mathcal{E}} \cap \{\langle 0, \frac{1}{2} \rangle, \langle \frac{1}{2}, 0 \rangle\}) \neq \emptyset$. This contradicts to the fact that $0 \notin \{\frac{1}{2}, 1\} = D^{\mathcal{A}}$. Thus, \mathcal{B} is not disjunctive. In this way, (2.15), Remark 2.7(ii)(a), Corollaries 5.11, 5.21 and Lemma 5.10 complete the argument. \square

Theorem 5.22 provides an effective algebraic criterion of C 's being \sim -subclassical, according to which C , being \sim -subclassical but not \sim -classical, has a unique \sim -classical extension to be denoted by C^{PC} . The item (b) cannot be omitted in (i), while the optional stipulations in (ii,iii) cannot be omitted, whereas that of disjunctivity in (iii) cannot be even “weakened”, in view of:

Example 5.23. Let $\Sigma \triangleq \{\vee, \sim\}$, \mathcal{B} the canonically \sim -classical Σ -matrix with $(i \vee^{\mathfrak{B}} j) \triangleq 1$, for all $i, j \in 2$, and \mathcal{A} both false-singular and involutive (in particular, \sim -paraconsistent) with $(a \vee^{\mathfrak{A}} b) \triangleq (\min(a, 1 - a) + \frac{1}{2})$, for all $a, b \in A$, in which case, as $(\text{img } \vee^{\mathfrak{A}/\mathfrak{B}}) \subseteq D^{\mathfrak{A}/\mathfrak{B}}$, \mathcal{A}/\mathfrak{B} is weakly \vee -disjunctive, and so C/\mathfrak{B} is weakly \vee -disjunctive/“ \vee -conjunctive (cf. Remark 2.7(i)(a))”, respectively. Then, we have $(\langle \frac{1}{2}|a, a|\frac{1}{2} \rangle \vee^{\mathfrak{A}^2} b) = \langle 1|\frac{1}{2}, \frac{1}{2}|1 \rangle \in (L_4 \cap D^{\mathfrak{A}^2})$, for all $a \in 2$ and all $b \in A^2$. Hence, L_4 forms a subalgebra of \mathfrak{A}^2 , while $\chi^{\mathfrak{A}^2 \upharpoonright L_4} \in \text{hom}_{\mathfrak{S}}^{\mathfrak{S}}(\mathfrak{A}^2 \upharpoonright L_4, \mathfrak{B})$, in which case, by (2.15), $\mathfrak{B} \in \text{Mod}(C)$, and so C is \sim -subclassical, whereas $(0 \vee^{\mathfrak{A}} 1) = \frac{1}{2}$, in which case \mathcal{A} is not classically hereditary, and so, by Theorem 5.22(ii,iii), C is neither disjunctive nor weakly conjunctive nor genuinely \sim -subclassical (in particular, \mathfrak{B} is not genuinely \sim -classical, and so neither conjunctive nor disjunctive nor implicative). \square

5.2.2.2. Characteristic matrices. Perhaps, the principal peculiarity of non-classical U3VLSN is as follows:

Theorem 5.24. *Let \mathfrak{B} be a [canonical] \sim -super-classical Σ -matrix. Suppose C is non- \sim -classical and defined by \mathfrak{B} . Then, \mathfrak{B} is isomorphic [and so equal] to \mathcal{A} . In particular, any uniform three-valued expansion of C is defined by a unique expansion of \mathcal{A} , unless C is \sim -classical.*

Proof. Then, the canonization \mathcal{D} of \mathfrak{B} is isomorphic to \mathfrak{B} , in which case, by (2.15), C is defined by \mathcal{D} , and so, by Corollary 5.18, both \mathcal{A} and \mathcal{D} are simple. Hence, by Remark 2.5(ii) and Lemma 3.7, $(\mathcal{A}|\mathcal{D}) \in \mathbf{H}(\mathbf{P}^{\text{SD}}(\mathbf{S}(\mathcal{D}|\mathcal{A})))$ (in particular, \mathcal{A} is truth-singular iff \mathcal{D} is so, for truth-singularity is preserved under \mathbf{P} as well as both \mathbf{S} and \mathbf{H} ; cf. Remark 2.7(ii)(c)). Therefore, there are some finite set I , some $\bar{C} \in \mathbf{S}(\mathcal{A})^I$, some subdirect product \mathcal{E} of it and some $h \in \text{hom}_{\mathfrak{S}}^{\mathfrak{S}}(\mathcal{E}, \mathcal{D})$, in which case, by (2.15) and Remark 2.7(ii)(b), \mathcal{E} is a both consistent and truth-non-empty model of C , for \mathcal{D} is so, and so $I \neq \emptyset$. Consider the following complementary cases:

- $(I \times \{j\}) \in E$, for some $j \in 2$, in which case $E \ni \sim^{\mathfrak{E}}(I \times \{j\}) = (I \times \{1 - j\})$, and so, as $2 = \{j, 1 - j\}$, E contains both of $(a|b) \triangleq (I \times \{1|0\})$. Consider the following complementary subcases:
 - $(I \times \{\frac{1}{2}\}) \in E$, in which case, as $I \neq \emptyset$, $g \triangleq \{(a', I \times \{a'\}) \mid a' \in A\}$ is an embedding of \mathcal{A} into \mathcal{E} , and so, by Remark 2.5(ii), $g \circ h$ is an embedding of \mathcal{A} into \mathcal{D} (in particular, is an isomorphism from \mathcal{A} onto \mathcal{D} , because $|A| = 3 \leq l$, for no $l \in 3 = |D|$).
 - $(I \times \{\frac{1}{2}\}) \notin E$, in which case \mathcal{E} is non- \sim -paraconsistent, and so is \mathfrak{B} , in view of (2.15) (in particular, \mathcal{A} is so). Then, 2 forms a subalgebra of \mathfrak{A} , for, otherwise, there would be some $\phi \in \text{Fm}_{\Sigma}^2$ such that $\phi^{\mathfrak{A}}(1, 0) = \frac{1}{2}$, in which case E would contain $\phi^{\mathfrak{E}}(a, b) = (I \times \{\frac{1}{2}\})$, and so, by (2.15), $\mathcal{F} \triangleq (\mathcal{A} \upharpoonright 2)$ is a canonical \sim -classical model of C (in particular, the logic C' of \mathcal{F} is a \sim -classical extension of C). Moreover, as $a \in D^{\mathcal{E}} \not\cong b$, for $I \neq \emptyset$, $h(a) \in D^{\mathcal{D}} \not\cong h(b)$, in which case $h(b/a) = (0|1)$, whenever \mathcal{D} is false-/truth-singular, respectively, and so $(1|0) = \sim^{\mathfrak{D}}(0|1) = h(\sim^{\mathfrak{E}}(b/a)) = h(a/b)$ (in particular, $h[\{a, b\}] = 2$). And what is more, as $h[E] = D$, there is some $c \in E$ such that $h(c) = \frac{1}{2}$. Let \mathcal{G} be the submatrix of \mathcal{E} generated by $\{a, b, c\}$, in which case $h' \triangleq (h|G) \in \text{hom}_{\mathfrak{S}}^{\mathfrak{S}}(\mathcal{G}, \mathcal{D})$, for $h[\{a, b, c\}] = A$, and so, by (2.15), C , being defined by \mathcal{D} , is defined by \mathcal{G} . Hence, $J \triangleq \{i \in I \mid \pi_i(c) = \frac{1}{2}\} \neq \emptyset$, otherwise, $2^I \supseteq \{a, b\}$ would contain c , in which case it, forming a subalgebra of \mathfrak{A}^I , would include G , and so \mathcal{G} , being a submatrix of \mathcal{A}^I , would be a submatrix of $\mathcal{F}^I \in \text{Mod}(C')$ (in particular, by (2.15), C , being a sublogic of C' , would be equal to C' , and so would be \sim -classical, for C' is so). Take any $j \in J \neq \emptyset$, in which case $\pi_j(a|b|c) = (1|0|\frac{1}{2})$, and

so $g' \triangleq (\pi_j \upharpoonright G) \in \text{hom}(\mathcal{G}, \mathcal{A})$ is surjective, for $\{a, b, c\} \subseteq G$. We prove, by contradiction, that $g' \in \text{hom}_{\mathbb{S}}^{\mathbb{S}}(\mathcal{G}, \mathcal{A})$. For suppose $g' \notin \text{hom}_{\mathbb{S}}^{\mathbb{S}}(\mathcal{G}, \mathcal{A})$, in which case there is some $d \in (G \setminus D^{\mathcal{G}})$ such that $\pi_j(d) \in D^{\mathcal{A}}$, and so $\pi_j(\sim^{\mathfrak{E}} d) = \sim^{\mathfrak{A}} \pi_j(d) \notin D^{\mathcal{A}}$, for, otherwise, (2.11) would not be true in \mathcal{A} under $[x_0/\pi_j(d), x_1/0]$. Then, $\sim^{\mathfrak{D}} d \notin D^{\mathcal{G}}$, in which case $\sim^{\mathfrak{D}} h'(d) = h'(\sim^{\mathfrak{E}} d) \notin D^{\mathcal{D}} \not\equiv h'(d)$, and so $D^{\mathcal{D}} \not\equiv h'(d) = \frac{1}{2}$ (in particular, \mathcal{D} is truth-singular, that is, \mathcal{A} is so). Let \mathcal{H} be the submatrix of \mathcal{G} generated by $\{a, b, d\}$, in which case $h'' \triangleq (h' \upharpoonright \mathcal{H}) \in \text{hom}_{\mathbb{S}}^{\mathbb{S}}(\mathcal{H}, \mathcal{D})$, for $h'[\{a, b, d\}] = A$, since $h'(a|b|d) = (1|0|\frac{1}{2})$, respectively, and so, by (2.15), C , being defined by \mathcal{D} , is defined by \mathcal{H} . Then, as \mathcal{A} is truth-singular, $\pi_j(d) = 1$, in which case, for each $i \in J$, we get $\pi_i(d) = \pi_j(d) = 1$, because $\pi_i(a|b|c) = (1|0|\frac{1}{2}) = \pi_j(a|b|c)$, respectively, and so $d \in 2^I \supseteq \{a, b\}$. Therefore, 2^I , forming a subalgebra of \mathfrak{A}^I , includes H , in which case \mathcal{H} , being a submatrix of \mathcal{A}^I , is that of $\mathcal{F}^I \in \text{Mod}(C')$, and so, by (2.15), C , being a sublogic of C' , is equal to C' (in particular, C is \sim -classical, for C' is so). This contradiction shows that $g' \in \text{hom}_{\mathbb{S}}^{\mathbb{S}}(\mathcal{G}, \mathcal{A})$. In this way, since both \mathcal{A} and \mathcal{D} are simple, while $h' \in \text{hom}_{\mathbb{S}}^{\mathbb{S}}(\mathcal{G}, \mathcal{D})$, by Remark 2.5(ii) and Lemma 3.6 with $\mathbb{M} = \{\mathcal{A}\}$, we eventually conclude that \mathcal{A} is isomorphic to \mathcal{D} .

- $(I \times \{j\}) \notin E$, for each $j \in 2$, in which case, by Lemma 5.16(i), \mathcal{A} is \sim -paraconsistent (in particular, false-singular, i.e., non-truth-singular), that is, \mathcal{B} is so, and so \mathcal{E} is \sim -paraconsistent, in view of (2.15), as well as is not truth-singular, in view Remark 2.7(ii)(c). Then, first, there is some $e \in D^{\mathcal{E}}$ such that $\sim^{\mathfrak{E}} e \in D^{\mathcal{E}}$, in which case $E \ni e \triangleq (I \times \{\frac{1}{2}\})$, and so \mathcal{A} is extra-classically hereditary, for, otherwise, there would be some $\psi \in \text{Fm}_{\mathbb{S}}^1$ such that $j \triangleq \psi^{\mathfrak{A}}(\frac{1}{2}) \in 2$, in which case E would contain $\psi^{\mathfrak{E}}(e) = (I \times \{j\})$. Second, there is some $f \in D^{\mathcal{E}} \subseteq \{\frac{1}{2}, 1\}^I$ distinct from e , in which case $K \triangleq \{i \in I \mid \pi_i(f) = 1\} \neq \emptyset$, and so, since \mathcal{A} is extra-classically hereditary and generated by $A \setminus \{0\}$, $g'' \triangleq \{\langle b', (K \times \{b'\}) \cup ((I \setminus K) \times \{\frac{1}{2}\}) \rangle \mid b' \in A\}$ is an embedding of \mathcal{A} into \mathcal{E} . Hence, by Remark 2.5(ii), $g'' \circ h$ is an embedding of \mathcal{A} into \mathcal{D} , and so is an isomorphism from \mathcal{A} onto \mathcal{D} , for $|A| = 3 = |D|$.

Thus, anyway, \mathcal{A} is isomorphic to \mathcal{D} , and so to \mathcal{B} [in which case, by Lemma 5.14(b), $\mathcal{A} = \mathcal{B}$]. Then, as \sim is a subclassical negation for any expansion of C , (2.15) and Theorem 5.7 end the proof. \square

In view of Theorem 5.24, \mathcal{A} , being uniquely determined by C , unless this is \sim -classical, is said to be *characteristic for/of* C . In view of Example 5.13, the stipulation of C 's being non- \sim -classical cannot be omitted in the formulation of Theorem 5.24, even if C is both conjunctive and implicative (in particular, disjunctive).

5.2.2.3. Self-extensionality versus discriminating endomorphisms. A *(truth-)discriminating operator/endomorphism on/of* \mathcal{A} is any $h \in (A^A / \text{hom}(\mathfrak{A}, \mathfrak{A}))$ such that $\chi^{\mathcal{A}}(h(\frac{1}{2})) \neq \chi^{\mathcal{A}}(h(\mathbb{k}^A))$, in which case $h(\frac{1}{2}) \neq h(\mathbb{k}^A)$, and so h is neither diagonal nor singular, the set of all them being denoted by $(\partial/\partial)(\mathcal{A})$, respectively. Then, since $\text{img}[\theta^{\mathcal{A}} \setminus \Delta_{\mathcal{A}}] = \{\{\frac{1}{2}, \mathbb{k}^A\}\}$, by Example 4.2, Corollary 4.12 and Theorem 5.17(iii) \Rightarrow (i), we have:

Corollary 5.25. *[Providing \mathcal{A} is either implicative or both conjunctive and disjunctive] C is self-extensional iff either it is \sim -classical or $\partial(\mathcal{A}) \neq \emptyset$.*

Though there are $3^3 = 27$ unary operations on A , only few of them may be discriminating operators/endomorphisms on/of \mathcal{A} . More precisely, let $h_{+|- , a} \triangleq (\Delta_2^{+|-} \cup \{\langle \frac{1}{2}, a \rangle\}) \in A^A$, where $a \in A$, $\mathcal{H} \triangleq (\bigcup_{a \in A} \{h_{+, a}, h_{-, a}\})$ and $\mathcal{H}^{\mathcal{A}} \triangleq (\{h_{-, a} \mid$

$a \in A$, $\chi^{\mathcal{A}}(a) = \mathbb{k}^{\mathcal{A}} \cup \{h_{+,1-\mathbb{k}^{\mathcal{A}}}\}$). Clearly,

$$(5.5) \quad (\mathcal{H} \cap \partial(\mathcal{A})) = \mathcal{H}^{\mathcal{A}}.$$

Conversely, since $\bar{\partial}(\mathcal{A}) = (\partial(\mathcal{A}) \cap \text{hom}(\mathfrak{A}, \mathfrak{A}))$, by (5.5) and Lemma 5.14(i) with $\mathcal{D} = \mathcal{A} = \mathcal{B}$, we have:

Corollary 5.26. $\bar{\partial}(\mathcal{A}) \subseteq \mathcal{H}$. In particular, $\bar{\partial}(\mathcal{A}) = (\mathcal{H}^{\mathcal{A}} \cap \text{hom}(\mathfrak{A}, \mathfrak{A}))$.

Combining Corollaries 5.25 and 5.26, we eventually get:

Theorem 5.27. [Providing \mathcal{A} is either implicative or both conjunctive and disjunctive] C is self-extensional iff either it is \sim -classical or $(\mathcal{H}^{\mathcal{A}} \cap \text{hom}(\mathfrak{A}, \mathfrak{A})) \neq \emptyset$.

This yields a quite effective purely-algebraic criterion of the self-extensionality of C with either implicative or both conjunctive and disjunctive \mathcal{A} that can inevitably be enhanced a bit more under separate studying the alternatives involved excluding *a priori* some elements of $\mathcal{H}^{\mathcal{A}}$ from $\bar{\partial}(\mathcal{A})$ (i.e., from $\text{hom}(\mathfrak{A}, \mathfrak{A})$; cf. Corollary 5.26), because, under the stipulation of C 's being both self-extensional and non- \sim -classical, the alternatives under considerations are disjoint, as it is shown below.

5.2.2.3.1. Self-extensionality versus equational truth-definitions.

Lemma 5.28. Let $\bar{\cup}$ be an equational truth definition for \mathcal{A} . Suppose \mathcal{A} is either false-singular or \sqsupset -implicative, while C is not \sim -classical. Then, any non-singular endomorphism h of \mathfrak{A} is diagonal. In particular, providing \mathcal{A} is either implicative or both conjunctive and disjunctive, C is not self-extensional.

Proof. Then, for any $a \in A$, we have $(a \in D^{\mathcal{A}}) \Leftrightarrow (\mathfrak{A} \models (\bigwedge \bar{\cup})[x_0/a]) \Rightarrow (\mathfrak{A} \models (\bigwedge \bar{\cup})[x_0/h(a)]) \Leftrightarrow (h(a) \in D^{\mathcal{A}})$, in which case $h \in \text{hom}(\mathcal{A}, \mathcal{A})$ (in particular, $h(1) \neq 0$, for $1 \in D^{\mathcal{A}} \not\equiv 0$), and so, by Lemma 5.14(i) with $\mathcal{D} = \mathcal{A} = \mathcal{B}$, $h \upharpoonright 2$ is diagonal. Therefore, if $h(\frac{1}{2})$ was equal to $\mathbb{k}^{\mathcal{A}}$, then h would be equal to $\chi^{\mathcal{A}}$, in which case $\theta^{\mathcal{A}} = (\ker h)$ would be a congruence of \mathfrak{A} , and so, by Theorem 5.17, C would be \sim -classical. Hence, in case \mathcal{A} is false-singular, $h(\frac{1}{2}) = \frac{1}{2}$, for $\frac{1}{2} \in D^{\mathcal{A}} \not\equiv 0$. Otherwise, \mathcal{A} is \sqsupset -implicative, in which case $(\frac{1}{2} \sqsupset^{\mathfrak{A}} 0) = 1$ and $(1 \sqsupset^{\mathfrak{A}} 0) \neq 1$, and so $h(\frac{1}{2}) = \frac{1}{2}$, for otherwise, we would have $h(\frac{1}{2}) = 1$, in which case we would get $1 \neq 1$. Thus, in any case, $h(\frac{1}{2}) = \frac{1}{2}$, and so h is diagonal. In this way, Corollary 4.13 and Theorem 5.17(iii) \Rightarrow (i) complete the argument. \square

This ‘‘equational truth definition’’ analogue of Corollary 4.15 provides another and much more transparent insight into the non-self-extensionality of the instances discussed in Example 4.17 and summarized below. In this connection, we first have:

Corollary 5.29. Suppose \mathcal{A} is both \sqsupset -implicative and either weakly $\bar{\wedge}$ -conjunctive (in particular, \wr -negative with $\bar{\wedge} = \uplus \sqsupset$; cf. Remark 2.7(i)(a)) or truth-singular. Then, \mathcal{A} has a finitary equational truth-definition. In particular, C is not self-extensional, unless it is \sim -classical.

Proof. The case, when \mathcal{A} is truth-singular, is due to Remark 4.14(iv). Otherwise, \mathcal{A} is weakly $\bar{\wedge}$ -conjunctive, while $\{\frac{1}{2}\}$ does [not] form a subalgebra of \mathfrak{A} [that is, there is some $\varphi \in \text{Fm}_{\Sigma}^1$ such that $\varphi^{\mathfrak{A}}(a) \in 2$], so $\{(x_0 \sqsupset \phi) \approx \phi\}$ with $\phi \triangleq (\psi[\bar{\wedge}(\psi[x_0/\varphi])])$ and $\psi \triangleq (x_0 \bar{\wedge} \sim x_0)$ is a finitary equational truth definition for \mathcal{A} . In this way, Lemma 5.28 completes the argument. \square

This is why the contexts of the next two subparagraphs are disjoint, whenever C is self-extensional but not \sim -classical. Before coming to discussing them, we provide practically immediate applications of the above results of this subparagraph to some of the logics specified in Paragraph 5.2.1.1.

Remark 5.30. Suppose \mathcal{A} is both \sim -paraconsistent (and so false-singular), conjunctive and \vee -disjunctive as well as both classically- and extra-classically-hereditary. Then, $\{x_0 \approx (x_0 \vee \sim x_0)\}$ is an equational truth definition for \mathcal{A} , so, by Remark 2.7(i)(c) and Lemma 5.28, C is not self-extensional. \square

This subsumes disjunctive conjunctive \sim -paraconsistent LP and HZ , providing a more transparent insight into the non-self-extensionality of them than that given by Example 4.17. Likewise, $[I]P^1$ is subsumed by:

Remark 5.31. Suppose \mathcal{A} is both classically-valued and \diamond -conjunctive/ \vee -disjunctive (in particular, \sqsupset -implicative with $\diamond = \sqsupset\sqsupset$). Then, it is \wr -negative, where $\wr x_0 \triangleq \sim(x_0 \diamond x_0)$, in which case, by Remark 2.7(i)(a), \mathcal{A} is both $\bar{\wedge}$ -conjunctive and \vee -disjunctive, where $\bar{\wedge} \triangleq \diamond/\wr$ and $\vee \triangleq \diamond/\wr$, and so, by Remark 2.7(i)(b), \mathcal{A} is $\sqsupset\vee$ -implicative. On the other hand, as $\frac{1}{2} \notin 2$, any idempotent binary operation on A , being term-wise definable in \mathfrak{A} , is so by either x_0 or x_1 , in which case it is not symmetric, for A is not a singleton, and so \mathfrak{A} is not a semi-lattice (in particular, is not a [distributive] lattice). And what is more, $\{(x_0 \sqsupset\vee x_0) \sqsupset\vee x_0 \approx (x_0 \sqsupset\vee x_0)\}$ is a finitary equational truth definition for \mathcal{A} , so, providing \mathcal{A} is not \sim -negative (in which case it is \sim -paraconsistent/ (\vee, \sim) -paracomplete, whenever it is false- \wr -truth-singular), so, by Remark 2.7(i)(c) and Lemma 5.28, C is not self-extensional. \square

5.2.2.3.2. Conjunctive U3VLSN.

Lemma 5.32. *Let \mathcal{B} be a consistent/truth-non-empty weakly \diamond -conjunctive/ \vee -disjunctive Σ -matrix. Suppose \mathfrak{B} is a \diamond -semi-lattice with bound. Then, $\beta_{\diamond}^{\mathfrak{B}} \notin / \in D^{\mathfrak{B}}$.*

Proof. By the weak \diamond -conjunctivity/ \vee -disjunctivity of \mathcal{B} , we do have $\beta_{\diamond}^{\mathfrak{B}} = (\beta_{\diamond}^{\mathfrak{B}} \diamond^{\mathfrak{B}} a) \notin / \in D^{\mathfrak{B}}$, where $a \in ((B \setminus D^{\mathfrak{B}})/D^{\mathfrak{B}}) \neq \emptyset$. \square

Lemma 5.33. *Suppose C is weakly $\bar{\wedge}$ -conjunctive. Then, \mathfrak{A} is a $\bar{\wedge}$ -semi-lattice with bound such that the following hold:*

- (i) $(0 \bar{\wedge}^{\mathfrak{A}} 1) = \beta_{\bar{\wedge}}^{\mathfrak{A}}$;
- (ii) $\frac{1}{2} \leq_{\bar{\wedge}}^{\mathfrak{A}} 1$;
- (iii) [providing $\bar{\delta}(\mathcal{A}) \neq \emptyset$, $(g) \Rightarrow (a) \Rightarrow (b) \Rightarrow (c) \Leftrightarrow (d) \Leftrightarrow (e) \Leftrightarrow (f) \Rightarrow (g) \Rightarrow (h) [\Rightarrow (f)]$, where:
 - (a) $h_{+,1-\mathbb{k}^{\mathcal{A}}} \in \text{hom}(\mathfrak{A}, \mathfrak{A})$;
 - (b) \mathcal{A} is classically-hereditary;
 - (c) $\beta_{\bar{\wedge}}^{\mathfrak{A}} = 0$;
 - (d) $0 \leq_{\bar{\wedge}}^{\mathfrak{A}} \frac{1}{2}$;
 - (e) $0 \leq_{\bar{\wedge}}^{\mathfrak{A}} 1$;
 - (f) \mathcal{A} is not involutive;
 - (g) $h_{-,a} \in \text{hom}(\mathfrak{A}, \mathfrak{A})$, for no $a \in A$;
 - (h) $h_{-, \frac{1}{2}} \notin \text{hom}(\mathfrak{A}, \mathfrak{A})$;
- (iv) \mathcal{A} is not \sim -negative, unless $\bar{\delta}(\mathcal{A}) = \emptyset$.

Proof. In that case, by Theorem 4.6(i) \Rightarrow (iv), \mathfrak{A} , being finite, is a $\bar{\wedge}$ -semi-lattice with bound, so, by Lemma 5.32, $\beta_{\bar{\wedge}}^{\mathfrak{A}} \notin D^{\mathcal{A}}$. Let $\xi_{0[+1]} \triangleq [\sim]x_0$ as well as both $\phi_k \triangleq \xi_k(x_0 \bar{\wedge} \sim x_0)$ and $\psi_k \triangleq \phi_k(\sim x_0)$, where $k \in 2$.

- (i) In case $\beta_{\bar{\wedge}}^{\mathfrak{A}} = 0$, we have $0 = \beta_{\bar{\wedge}}^{\mathfrak{A}} \leq^{\mathfrak{A}} 1$, and so get $(0 \bar{\wedge}^{\mathfrak{A}} 1) = 0 = \beta_{\bar{\wedge}}^{\mathfrak{A}}$. Otherwise, as $1 \in D^{\mathcal{A}}$, we have $D^{\mathcal{A}} \not\cong \beta_{\bar{\wedge}}^{\mathfrak{A}} = \frac{1}{2}$, in which case \mathcal{A} is truth-singular, and so is non- \sim -paraconsistent, that is, C is so. Then, by (2.11) and the conjunctivity of C , we have $x_1 \in C(\phi_0)$, in which case, by Theorem 4.6(i) \Rightarrow (iv), we get $\beta_{\bar{\wedge}}^{\mathfrak{A}} \leq^{\mathfrak{A}} (0 \bar{\wedge}^{\mathfrak{A}} 1) = \phi_0^{\mathfrak{A}}(0) \leq_{\bar{\wedge}}^{\mathfrak{A}} \beta_{\bar{\wedge}}^{\mathfrak{A}}$, and so eventually get $(0 \bar{\wedge}^{\mathfrak{A}} 1) = \beta_{\bar{\wedge}}^{\mathfrak{A}}$.

- (ii) Consider the following complementary cases:
- \mathcal{A} is false-singular, in which case, by (i), for each $k \in 2$, $\phi_0^{\mathfrak{A}}(k) = \phi_0^{\mathfrak{A}}(0) = \beta_{\bar{\wedge}}^{\mathfrak{A}} = 0$, and so $(\phi|\psi)_1^{\mathfrak{A}}(k) = 1 \in D^{\mathcal{A}}$. Consider the following complementary subcases:
 - $\sim^{\mathfrak{A}}\frac{1}{2} = \frac{1}{2}$, in which case $\phi_1^{\mathfrak{A}}(\frac{1}{2}) = \frac{1}{2} \in D^{\mathcal{A}}$, for \mathcal{A} is false-singular, and so ϕ_1 is true in \mathcal{A} (in particular, $\phi_1 \in C(x_1)$). Then, by Theorem 4.6(i) \Rightarrow (iv), $\frac{1}{2} \leq_{\bar{\wedge}}^{\mathfrak{A}} \phi_1^{\mathfrak{A}}(0) = 1$.
 - $\sim^{\mathfrak{A}}\frac{1}{2} \neq \frac{1}{2}$, that is, $\sim^{\mathfrak{A}}\frac{1}{2} \in 2$, in which case $\psi_1^{\mathfrak{A}}(\frac{1}{2}) = \phi_1^{\mathfrak{A}}(\sim^{\mathfrak{A}}\frac{1}{2}) = 1 \in D^{\mathcal{A}}$, and so ψ_1 is true in \mathcal{A} (in particular, $\psi_1 \in C(x_1)$). Then, by Theorem 4.6(i) \Rightarrow (iv), $\frac{1}{2} \leq_{\bar{\wedge}}^{\mathfrak{A}} \psi_1^{\mathfrak{A}}(0) = 1$.
 - \mathcal{A} is truth-singular, in which case it is non- \sim -paraconsistent, that is, C is so, and so, by (2.11) and the $\bar{\wedge}$ -conjunctivity of C , $x_1 \in C(\phi_0)$. Consider the following complementary subcases:
 - $\frac{1}{2}$ is equal to either $\beta_{\bar{\wedge}}^{\mathfrak{A}}$ or $\sim^{\mathfrak{A}}\frac{1}{2}$, in which case we have $\frac{1}{2} = \phi_0^{\mathfrak{A}}(\frac{1}{2})$, and so, by Theorem 4.6(i) \Rightarrow (iv), get $\frac{1}{2} \leq_{\bar{\wedge}}^{\mathfrak{A}} 1$, for $x_1 \in C(\phi_0)$.
 - $\beta_{\bar{\wedge}}^{\mathfrak{A}} \neq \frac{1}{2} \neq \sim^{\mathfrak{A}}\frac{1}{2}$, in which case, as $1 \in D^{\mathcal{A}}$, by (i), for each $k \in 2$, $\phi_0^{\mathfrak{A}}(k) = (0 \bar{\wedge}^{\mathfrak{A}} 1) = \beta_{\bar{\wedge}}^{\mathfrak{A}} = 0$, and so $(\phi|\psi)_1^{\mathfrak{A}}(k) = 1 \in D^{\mathcal{A}}$ (in particular, $\psi_1^{\mathfrak{A}}(\frac{1}{2}) = \phi_1^{\mathfrak{A}}(\sim^{\mathfrak{A}}\frac{1}{2}) = 1 \in D^{\mathcal{A}}$). Then, ψ_1 is true in \mathcal{A} , in which case $\psi_1 \in C(x_1)$, and so, by Theorem 4.6(i) \Rightarrow (iv), $\frac{1}{2} \leq_{\bar{\wedge}}^{\mathfrak{A}} \psi_1^{\mathfrak{A}}(0) = 1$.
- (iii) First, (d/h) is a particular case of (c/g), while (d/e) \Rightarrow (e/c) is by (ii/i), whereas (b) \Rightarrow (e) is by the $\bar{\wedge}$ -conjunctivity of \mathcal{A} and the fact that $1 \in D^{\mathcal{A}} \not\equiv 0$. Next, (a) \Rightarrow (b) is by the fact that $\text{img}(h_{+,1-k^{\mathcal{A}}}) = 2$. Further, assume (f) holds, in which case $l \triangleq \sim^{\mathfrak{A}}\frac{1}{2} \in 2$, and so $\xi_{1-l}^{\mathfrak{A}}(\frac{1}{2}) = 1 \in D^{\mathcal{A}}$. We prove (e) by contradiction. For suppose (e) does not hold, in which case $\beta_{\bar{\wedge}}^{\mathfrak{A}} \neq 0$, and so, by Lemma 5.32, $\beta_{\bar{\wedge}}^{\mathfrak{A}} = \frac{1}{2}$, for $1 \in D^{\mathcal{A}}$ (in particular, $\phi_0^{\mathfrak{A}}(\frac{1}{2}) = \frac{1}{2}$). Likewise, by (i), for each $k \in 2$, $\phi_0^{\mathfrak{A}}(k) = (0 \bar{\wedge}^{\mathfrak{A}} 1) = \beta_{\bar{\wedge}}^{\mathfrak{A}} = \frac{1}{2}$, in which case ϕ_{1-l} is true in \mathcal{A} , and so $\phi_{1-l} \in C(x_1)$. Then, by Theorem 4.6(i) \Rightarrow (iv), $0 \leq_{\bar{\wedge}}^{\mathfrak{A}} \phi_{1-l}^{\mathfrak{A}}(0) = 1$. Thus, (e) holds. [Conversely, assume (f) does not hold, in which case $\sim^{\mathfrak{A}}a = (1-a)$, for all $a \in A$. Take any $h \in \bar{\partial}(\mathfrak{A}) \neq \emptyset$, in which case it is neither diagonal nor singular, and so, by Lemma 5.14, $(h|2) \in \{\Delta_2^+, \Delta_2^-\}$. Then, we have $h(\frac{1}{2}) = h(\sim^{\mathfrak{A}}\frac{1}{2}) = \sim^{\mathfrak{A}}h(\frac{1}{2}) = (1 - h(\frac{1}{2}))$, in which case we get $h(\frac{1}{2}) = \frac{1}{2}$, and so $h = h_{-, \frac{1}{2}}$, for, otherwise, h would be diagonal. Thus, (h) \Rightarrow (f) holds.] Now, assume (e) holds (that is, (c) does so), in which case, for each $k \in 2$, $\phi_0^{\mathfrak{A}}(k) = (0 \bar{\wedge}^{\mathfrak{A}} 1) = 0$, and so $\phi_1^{\mathfrak{A}}(k) = 1 \in D^{\mathcal{A}}$. We prove (f) by contradiction. For suppose $\sim^{\mathfrak{A}}\frac{1}{2} = \frac{1}{2}$, in which case $\phi_0^{\mathfrak{A}}(\frac{1}{2}) = \frac{1}{2}$, and so $\phi_1^{\mathfrak{A}}(\frac{1}{2}) = \frac{1}{2}$. Consider the following complementary cases:
 - \mathcal{A} is false-singular, in which case $\phi_1^{\mathfrak{A}}(\frac{1}{2}) = \frac{1}{2} \in D^{\mathcal{A}}$, and so ϕ_1 is true in \mathcal{A} (in particular, $\phi_1 \in C(x_1)$). Then, by Theorem 4.6(i) \Rightarrow (iv), $1 \leq_{\bar{\wedge}}^{\mathfrak{A}} \phi_1^{\mathfrak{A}}(\frac{1}{2}) = \frac{1}{2}$, in which case, by (ii), $\frac{1}{2} = 1$, and so $\frac{1}{2} \in 2$.
 - \mathcal{A} is truth-singular, in which case it is not \sim -paraconsistent, and so, by (2.11) and the $\bar{\wedge}$ -conjunctivity of C , $x_1 \in C(\phi_0)$. Then, by Theorem 4.6(i) \Rightarrow (iv), $\frac{1}{2} = \phi_0^{\mathfrak{A}}(\frac{1}{2}) \leq_{\bar{\wedge}}^{\mathfrak{A}} 0$, in which case, by (c), $\frac{1}{2} = 0$, and so $\frac{1}{2} \in 2$. Thus, as $\frac{1}{2} \notin 2$, (f) does hold. Furthermore, if any $h : A \rightarrow A$ with $(h|2) = \Delta_2^-$ was an endomorphism of \mathfrak{A} , then, by (e), we would have $1 = h(0) = h(0 \bar{\wedge}^{\mathfrak{A}} 1) = (h(0) \bar{\wedge}^{\mathfrak{A}} h(1)) = (1 \bar{\wedge}^{\mathfrak{A}} 0) = (0 \bar{\wedge}^{\mathfrak{A}} 1) = 0$, and so (g) holds. [Finally, (g) \Rightarrow (a) is by (5.5) and Lemma 5.14, for $\bar{\partial}(\mathcal{A}) = (\partial(\mathcal{A}) \cap \text{hom}(\mathfrak{A}, \mathfrak{A}))$.]

(iv) Assume $\bar{\partial}(\mathcal{A}) \neq \emptyset$. Then, \mathcal{A} is not \sim -negative, whenever it is involutive. Otherwise, by (iii)(f) \Rightarrow (a), $h \triangleq h_{+,1-k^{\mathcal{A}}} \in \text{hom}(\mathfrak{A}, \mathfrak{A})$, in which case, if \mathcal{A}

was \sim -negative, then we would have $\sim^{\mathfrak{A}} \frac{1}{2} = (1 - \mathbb{k}^{\mathcal{A}})$, and so would get $2 \ni \mathbb{k}^{\mathcal{A}} = \sim^{\mathfrak{A}}(1 - \mathbb{k}^{\mathcal{A}}) = \sim^{\mathfrak{A}} h(\frac{1}{2}) = h(\sim^{\mathfrak{A}} \frac{1}{2}) = h(1 - \mathbb{k}^{\mathcal{A}}) = (1 - \mathbb{k}^{\mathcal{A}})$. \square

Theorem 5.34. *Suppose C is $\bar{\wedge}$ -conjunctive, non- \sim -classical and self-extensional. Then, $\bar{\delta}(\mathcal{A}) \neq \emptyset$.*

Proof. Then, by Corollary 5.18, \mathcal{A} is hereditarily simple, while, by Theorem 4.6(i) \Rightarrow (iv) and Lemma 5.32, \mathfrak{A} , being finite, is a $\bar{\wedge}$ -semi-lattice with bound $\beta_{\bar{\wedge}}^{\mathfrak{A}} \notin D^{\mathcal{A}}$, in which case, as $\frac{1}{2} \notin 2 \ni \mathbb{k}^{\mathcal{A}}$ (in particular, $\frac{1}{2} \neq \mathbb{k}^{\mathcal{A}}$), by the commutativity identity for $\bar{\wedge}$, there are some $\bar{a} \in (\{\frac{1}{2}, \mathbb{k}^{\mathcal{A}}\}^2 \setminus \Delta_{\mathcal{A}})$ and some $i \in 2$ such that $a_{1-i} \neq (a_i \bar{\wedge}^{\mathfrak{A}} a_{1-i})$, and so $\mathcal{B} \triangleq \langle \mathfrak{A}, F \rangle$, where $a_i \in F \triangleq \{b' \in A \mid a_i \leq_{\bar{\wedge}}^{\mathfrak{A}} b'\} \not\ni a_{1-i}$, being both truth-non-empty and $\bar{\wedge}$ -conjunctive, is a finite consistent truth-non-empty model of C . Then, as 2 forms a subalgebra of $\mathfrak{A} \upharpoonright \Sigma_{\sim}$, by Remark 2.7(ii)(b), Lemmas 3.7, 5.16(i,ii) with $\Sigma' = \Sigma_{\sim}$ and the conjunctivity of \mathcal{A} , $((\mathcal{A} \upharpoonright \Sigma_{\sim}) \upharpoonright 2)$, being \sim -classical, belongs to $\mathbf{I}(\mathbf{S}(\mathbf{H}^{-1}(\mathbf{H}(\mathcal{B} \upharpoonright \Sigma_{\sim}))))$, in which case, by (2.15), \sim is a subclassical negation for the logic C' of \mathcal{B} , and so, by Theorem 5.7, \mathcal{B} , being three-valued, is \sim -super-classical. Let \mathcal{D} be the canonization of \mathcal{B} , in which case they are isomorphic, and so, by (2.15), C' is defined by \mathcal{D} . Consider the following complementary cases:

- C' is \sim -classical, in which case, as it is $\bar{\wedge}$ -conjunctive, for its sublogic C is so, by Corollary 5.18, \mathcal{D} is a strictly surjectively homomorphic counter-image of a \sim -classical Σ -matrix \mathcal{E} , and so is \mathcal{B} , being isomorphic to \mathcal{D} . Then, by (2.15), \mathcal{E} is a \sim -classical model of C , for $\mathcal{B} \in \text{Mod}(C)$, in which case C is both \sim -subclassical and $\bar{\wedge}$ -conjunctive, and so, by Corollary 5.21 and Theorem 5.22(iii), \mathcal{A} is classically hereditary, $\mathcal{A} \upharpoonright 2$ being isomorphic to \mathcal{E} (in particular, \mathcal{B} is a strictly [surjectively] homomorphic counter-image of $\mathcal{A} \upharpoonright 2$).
- C' is not \sim -classical, in which case, by Corollary 5.18, \mathcal{D} , being canonically \sim -super-classical and defining C' , is simple (in particular, \mathcal{B} , being isomorphic to \mathcal{D} , is so, in view of Remark 2.5(iii)). Hence, by Lemma 3.7, there are some finite set I , some $\bar{c} \in \mathbf{S}_*(\mathcal{A})^I$, some subdirect product \mathcal{G} of it and some $g \in \text{hom}_{\mathbf{S}}^{\mathbf{S}}(\mathcal{G}, \mathcal{B})$, in which case, by Remark 2.7(ii)(b), \mathcal{G} is both consistent (in particular, $I \neq \emptyset$) and truth-non-empty, for \mathcal{B} is so, and so, by Lemma 5.16(i), $a \triangleq (I \times \{1\}) \in G \ni b \triangleq (I \times \{0\})$. We prove, by contradiction, that $(I \times \{\frac{1}{2}\}) \in G$. For suppose $(I \times \{\frac{1}{2}\}) \notin G$, in which case \mathcal{A} is classically-hereditary, for, otherwise, there would be some $\varphi \in \text{Fm}_{\Sigma}^2$ such that $\varphi^{\mathfrak{A}}(0, 1) = \frac{1}{2}$, and so $G \supseteq \{a, b\}$ would contain $\varphi^{\mathfrak{B}}(b, a) = (I \times \{\frac{1}{2}\})$. Consider the following complementary subcases:

- \mathcal{A} is truth-singular, in which case \mathcal{B} is so, for truth-singularity is preserved under \mathbf{P} as well as, in view of Remark 2.7(ii)(c), under both \mathbf{S} and \mathbf{H} , and so $D^{\mathcal{B}} = \{a_i\}$ (in particular, by Lemma 5.33(ii), $a_i \neq \frac{1}{2}$, for $1 \neq \frac{1}{2}$). Then, $\beta_{\bar{\wedge}}^{\mathfrak{A}} \neq a_i = \mathbb{k}^{\mathcal{A}} = 0$, in which case, as $1 \in D^{\mathcal{A}}$, $\beta_{\bar{\wedge}}^{\mathfrak{A}} = \frac{1}{2} \neq 0$, and so, by Lemma 5.33(iii)(b) \Rightarrow (c), \mathcal{A} is not classically-hereditary.
- \mathcal{A} is false-singular, in which case, by Lemma 5.32, $\beta_{\bar{\wedge}}^{\mathfrak{A}} = 0$, and so, by Lemma 5.33(iii)(c) \Rightarrow (e/f), $(0 \leq_{\bar{\wedge}}^{\mathfrak{A}} 1) / (\sim^{\mathfrak{A}} \frac{1}{2} \in 2)$, respectively. And what is more, by Lemma 5.33(ii), $\frac{1}{2} \leq_{\bar{\wedge}}^{\mathfrak{A}} 1$, in which case $1 = \delta \beta_{\bar{\wedge}}^{\mathfrak{A}}$, while $a_i \neq \frac{1}{2}$, for, otherwise, we would have $\frac{1}{2} = a_i \not\leq_{\bar{\wedge}}^{\mathfrak{A}} a_{1-i} = \mathbb{k}^{\mathcal{A}} = 1$, and so $a_i = \mathbb{k}^{\mathcal{A}} = 1$ (in particular, $D^{\mathcal{B}} = \{1\}$, for $1 = \delta \beta_{\bar{\wedge}}^{\mathfrak{A}}$). Therefore, as $a \in D^{\mathcal{G}}$, for $1 \in D^{\mathcal{A}}$, $g(a) = 1$, in which case $g(b) = g(\sim^{\mathfrak{B}} a) = \sim^{\mathfrak{A}} g(a) = 0$, and so $g[\{a, b\}] = 2$. Furthermore, there is some $c \in G$ such that $g(c) = \frac{1}{2} \notin D^{\mathcal{B}}$, in which case $c \notin D^{\mathcal{G}}$, and so there is some

$j \in I$ such that $\pi_j(c) = 0$, for $\mathcal{C}_j \in \mathbf{S}_*(\mathcal{A})$, while 0 is the only non-distinguished value of \mathcal{A} . Let \mathcal{H} be the submatrix of \mathcal{G} generated by $\{a, b, c\}$, in which case $h \triangleq (g \upharpoonright H) \in \text{hom}_{\mathbb{S}}^{\mathbb{S}}(\mathcal{H}, \mathcal{B})$, for $g[\{a, b, c\}] = A$, and so, by (2.15), C' , being defined by \mathcal{B} , is defined by \mathcal{H} , while, since $\pi_j[\{a, b, c\}] = 2$ forms a subalgebra of \mathfrak{A} , $f \triangleq (\pi_j \upharpoonright H) \in \text{hom}(\mathcal{H}, \mathcal{A} \upharpoonright 2)$ is surjective. Consider the following complementary (for $\sim^{\mathfrak{A}} \frac{1}{2} \in 2$) subcases:

- $\ast \sim^{\mathfrak{A}} \frac{1}{2} = 1$, in which case \mathcal{B} is weakly \sim -negative, for $\sim^{\mathfrak{A}} 0 = 1 \in D^{\mathcal{B}}$, and so is \mathcal{H} , in view of Remark 2.7(ii)(a). Then, by Lemma 5.20, $f \in \text{hom}_{\mathbb{S}}^{\mathbb{S}}(\mathcal{H}, \mathcal{A} \upharpoonright 2)$, for $\mathcal{A} \upharpoonright 2$ is \sim -negative, in which case, by (2.15), C' , being defined by \mathcal{H} , is defined by the \sim -classical Σ -matrix $\mathcal{A} \upharpoonright 2$, and so is \sim -classical.
- $\ast \sim^{\mathfrak{A}} \frac{1}{2} = 0$, in which case \mathcal{A} is \sim -negative, and so, by Remarks 2.5(ii), 2.7(i)(a,b) and Theorem 3.2, h is injective, for \mathcal{A} is conjunctive and hereditarily simple. Then, $(h^{-1} \circ f) \in \text{hom}_{\mathbb{S}}^{\mathbb{S}}(\mathcal{B}, \mathcal{A} \upharpoonright 2)$, for $(h \upharpoonright f)(a/b) = (1/0) \in / \notin D^{\mathcal{B} \upharpoonright \mathcal{A}}$ and $(h \upharpoonright f)(c) = (\frac{1}{2} | 0) \notin D^{\mathcal{B} \upharpoonright \mathcal{A}}$, in which case, by (2.15), C' , being defined by \mathcal{B} , is defined by the \sim -classical Σ -matrix $\mathcal{A} \upharpoonright 2$, and so is \sim -classical.

Thus, anyway, we come to a contradiction, in which case $(I \times \{\frac{1}{2}\}) \in G$, and so, as $I \neq \emptyset$, while $a \in G \ni b$, $e \triangleq \{\langle a', I \times \{a'\} \mid a' \in A \rangle\}$ is an embedding of \mathcal{A} into \mathcal{G} . Therefore, by Remark 2.5(ii), $e' \triangleq (e \circ g)$ is an embedding of \mathcal{A} into \mathcal{B} , for \mathcal{A} is simple, in which case it is an isomorphism from \mathcal{A} onto \mathcal{B} , as $|A| = 3 \not\leq k$, for no $k \in 3 = |B|$, and so $e'^{-1} \in \text{hom}(\mathcal{B}, \mathcal{A})$ is strict.

In this way, in any case, there is some strict $h' \in \text{hom}(\mathcal{B}, \mathcal{A}) \subseteq \text{hom}(\mathfrak{A}, \mathfrak{A})$, in which case $h'(a_i) \in D^{\mathcal{A}} \not\equiv h'(a_{1-i})$, for $a_i \in D^{\mathcal{B}} \not\equiv a_{1-i}$, and so $h' \in \mathfrak{D}(\mathcal{A})$, as required. \square

Then, combining Theorems 5.22(iii), 5.34 and Corollary 5.25 with Lemmas 5.32 and 5.33(ii, iii,iv), we immediately get the following two corollaries:

Corollary 5.35. *Suppose C is both $\bar{\wedge}$ -conjunctive and non- \sim -classical, while \mathcal{A} is false-/truth-singular. Then, C is self-extensional iff /either $h_{+,1-\mathbb{k}\mathcal{A}}$ /“or $h_{-, \frac{1}{2}}$ ” is an endomorphism of \mathfrak{A} [while \mathfrak{A} is a $\bar{\wedge}$ -semi-lattice with $\frac{1}{2} \leq \frac{\mathfrak{A}}{\bar{\wedge}} 1$, whereas it is that with bound 0 and/iff it is that with dual bound 1 and/iff \mathcal{A} is non-involutive and/iff \mathcal{A} is classically-hereditary (i.e., C is \sim -subclassical), as well as \mathcal{A} is not \sim -negative].*

Corollary 5.36. *Suppose \mathcal{A} is both $\bar{\wedge}$ -conjunctive and \vee -disjunctive, while C is not \sim -classical. Then, C is self-extensional iff $h_{+,1-\mathbb{k}\mathcal{A}} \in \text{hom}(\mathfrak{A}, \mathfrak{A})$, in which case \mathfrak{A} is a distributive $(\bar{\wedge}, \vee)$ -lattice with zero 0 and unit 1, while \mathcal{A} is neither involutive nor \sim -negative as well as classically-hereditary, and so C is \sim -subclassical.*

These immediately yield the self-extensionality of $[P]G_3^{(*)}$, for $h_{+,1-\mathbb{k}\mathcal{A}}$ is an endomorphism of the underlying algebra of its conjunctive (disjunctive) characteristic matrix. And what is more, they immediately imply the non-self-extensionality of $[I]P^1$, for the underlying algebra of its conjunctive (disjunctive) characteristic matrix is not a semi-lattice at all {cf. Remark 5.31}. Likewise, the non-self-extensionality of the conjunctive (disjunctive) HZ {cf. Subparagraph 5.2.1.1.3} ensues from either the involutivity of its conjunctive (disjunctive) classically-hereditary characteristic matrix or the fact that the underlying algebra of this matrix, though being a distributive lattice, is not that with both zero 0 and unit 1. Finally, the above corollaries imply *immediately* the non-self-extensionality of $LP_{[01]}/K_{3[01]}$, in view of the involutivity of their conjunctive (disjunctive) classically-hereditary characteristic matrices, providing, as opposed to Example 4.17, a more [perhaps, the most]

transparent and immediate *generic* insight into the non-self-extensionality of the latter independent from that of the former, and so into that of Łukasiewicz' finitely-valued logics [6] {cf. Example 4.16}, for these are expansions of K_3 . On the other hand, Corollary/Theorem 5.36/4.7 does not subsume Corollary/Theorem 5.35/5.34, due to existence of self-extensional conjunctive but non-disjunctive non- \sim -classical uniform three-valued Σ -logics with subclassical negation \sim , in view of:

Example 5.37. Let $\Sigma \triangleq \{\wedge, \sim\}$ and \mathcal{A} the Σ -reduct of the [non-]truth-singular $\Sigma_{\sim,+,01}^{\supset}$ -matrix specified in Subparagraph 5.2.1.1.2, in which case the former is both \wedge -conjunctive and non- \sim -negative, for the latter is so, and so $[P]G_3^\wedge \triangleq C$, being the Σ -fragment of the self-extensional [paraconsistent counterpart of] Gödel's three-valued logic $[P]G_3$ [2], is both \wedge -conjunctive and self-extensional as well as, by Remark 5.12 and Corollary 5.18, not \sim -classical. On the other hand, by induction on construction of any $\varphi \in \text{Fm}_\Sigma^2$, we prove that either $\varphi^{\mathfrak{A}}(\frac{1}{2}, \frac{1}{2}) \neq \frac{1}{2}$ or there are some $a, b \in A$ such that $\max(a, b) \not\leq \varphi^{\mathfrak{A}}(a, b)$. In case $\varphi = x_{0|1}$, taking $a \triangleq (0|1)$ and $b \triangleq (1|0)$, we get $\max(a, b) = 1 \not\leq 0 = \varphi^{\mathfrak{A}}(a, b)$. Likewise, in case $\varphi = \sim\xi$, where $\xi \in \text{Fm}_\Sigma^2$, as $(\text{img } \sim^{\mathfrak{A}}) \subseteq 2 \not\supseteq \frac{1}{2}$, we have $\varphi^{\mathfrak{A}}(\frac{1}{2}, \frac{1}{2}) \neq \frac{1}{2}$. Finally, in case $\varphi = (\phi \wedge \psi)$, where $\phi, \psi \in \text{Fm}_\Sigma^2$, if $\varphi^{\mathfrak{A}}(\frac{1}{2}, \frac{1}{2})$ is equal to $\frac{1}{2}$, then so is either $\phi^{\mathfrak{A}}(\frac{1}{2}, \frac{1}{2})$ or $\psi^{\mathfrak{A}}(\frac{1}{2}, \frac{1}{2})$, for \mathcal{A} is classically-hereditary, while, if, for any $a, b \in A$, it holds that $\max(a, b) \leq \varphi^{\mathfrak{A}}(a, b) = \min(\phi^{\mathfrak{A}}(a, b), \psi^{\mathfrak{A}}(a, b))$, then both $\max(a, b) \leq \phi^{\mathfrak{A}}(a, b)$ and $\max(a, b) \leq \psi^{\mathfrak{A}}(a, b)$ hold, and so the induction hypothesis completes the argument. In particular, $\max \cap A^2$ is not term-wise definable in \mathfrak{A} . Therefore, by Lemma 5.10 and Corollary 5.36, $[P]G_3^\wedge$ is not disjunctive. \square

Example 5.38. Let $\Sigma \triangleq \{\wedge, \sim\}$ and \mathcal{A} both truth-singular and involutive (in particular, non- \sim -negative) with $(a \wedge^{\mathfrak{A}} a) \triangleq a$, for all $a \in A$, as well as $(a \wedge^{\mathfrak{A}} b) \triangleq \frac{1}{2}$, for all $b \in (A \setminus \{a\})$. Then, \mathfrak{A} is a \wedge -semi-lattice with bound $\frac{1}{2}$ and maximal elements in 2, in which case \mathcal{A} is \wedge -conjunctive and, being involutive, is not \sim -negative, and so C is $\bar{\wedge}$ -conjunctive and, by Remark 5.12 and Corollary 5.18, not \sim -classical. Moreover, $h_{-\frac{1}{2}}$ is an endomorphism of \mathfrak{A} , so, by Corollary 5.35, C is self-extensional, while, by Corollary 5.36, C is not disjunctive. \square

The latter example shows that the “involutive” alternative cannot be disregarded in Corollary 5.35, by which, among other things, any conjunctive self-extensional uniform three-valued non- \sim -classical logic with subclassical negation \sim is a \sim -conservative term-wise definitional expansion of either of the three instances discussed above, and so is \sim -paraconsistent, unless its characteristic matrix is truth-singular. Likewise, by Corollary 5.36, any conjunctive \vee -disjunctive self-extensional uniform three-valued non- \sim -classical logic with subclassical negation \sim and [non-]truth-singular characteristic matrix is a \sim -conservative term-wise definitional expansion of $[P]G_3^*$, and so is [not] non- \sim -paraconsistent as well as [non-] (\vee, \sim) -paracomplete.

5.2.2.3.3. Implicative U3VLSN. We start from marking the framework of the self-extensionality of C under its being both non- \sim -classical and implicative:

Corollary 5.39. *Suppose \mathcal{A} is \sqsupset -implicative. Then, C is not self-extensional, unless it is either \sim -paraconsistent or \sim -classical. In particular, C is not self-extensional, whenever \mathcal{A} is truth-singular (in particular, both (\vee, \sim) -paracomplete and weakly \vee -disjunctive).*

Proof. If \mathcal{A} is both false-singular and non- \sim -paraconsistent, then it is \sim -negative. So, Remarks 2.7(i)(c), 4.14(iv), Lemma 5.28 and Corollary 5.29 end the proof. \square

Theorem 5.40. *Suppose \mathcal{A} is \sqsupset -implicative, while C is not \sim -classical. Then, C is self-extensional iff $h_{-, \frac{1}{2}} \in \text{hom}(\mathfrak{A}, \mathfrak{A})$, in which case \mathfrak{A} is an \sqsupset -implicative intrinsic semi-lattice with bound $\frac{1}{2}$, whereas \mathcal{A} is involutive as well as not classically-hereditary, and so C is not \sim -subclassical.*

Proof. Assume C is self-extensional. Then, by Theorem 4.9, \mathfrak{A} is an \sqsupset -implicative intrinsic semi-lattice with bound $a \triangleq (\frac{1}{2} \sqsupset^{\mathfrak{A}} \frac{1}{2}) = (b \sqsupset^{\mathfrak{A}} b)$, for any $b \in A$, while, by Corollary 5.39, \mathcal{A} is \sim -paraconsistent (in particular, false-singular), in which case $a \in D^{\mathcal{A}} = \{\frac{1}{2}, 1\}$, and so $a = \frac{1}{2}$ [in particular, $\sim^{\mathfrak{A}} a \in D^{\mathcal{A}}$, and so $\sim^{\mathfrak{A}} a = \frac{1}{2}$], for, otherwise, we would have $[\sim^{\mathfrak{A}}]a = 1$, in which case we would get $\sim^{\mathfrak{A}}[\sim^{\mathfrak{A}}]a = \sim^{\mathfrak{A}}1 = 0 \notin D^{\mathcal{A}}$, and so \mathcal{A} would be \wr -negative, where $\wr x_0 \triangleq (x_0 \sqsupset \sim[\sim](x_0 \sqsupset x_0))$ (in particular, by Corollary 5.29, C would not be self-extensional). In that case, \mathcal{A} is involutive as well as not classically-hereditary, for $(0 \sqsupset^{\mathfrak{A}} 0) = a = \frac{1}{2} \notin 2 \ni 0$, while, for any $h \in \text{hom}(\mathfrak{A}, \mathfrak{A})$, we have $h(\frac{1}{2}) = (h(\frac{1}{2}) \sqsupset^{\mathfrak{A}} h(\frac{1}{2})) = \frac{1}{2}$, and so Theorems 5.22(iii) and 5.27 complete the argument. \square

Corollaries 5.36/5.35 and 5.40, in particular, “provide one more insight into their context’s being disjoint, in view of opposite requirements on the involutivity of characteristic matrices”/“taking Example 4.2 into account, immediately yield the following essential (mainly, due to elimination of the disjunctivity stipulation) enhancement of Theorem 5.27”:

Corollary 5.41. *Suppose \mathcal{A} is either implicative or conjunctive. Then, C is self-extensional iff either it is \sim -classical or $(\{h_{+, 1-\mathbb{k}\mathcal{A}}, h_{-, \frac{1}{2}}\} \cap \text{hom}(\mathfrak{A}, \mathfrak{A})) \neq \emptyset$.*

Finally, we present a term-wise definitionally minimal instance of a self-extensional paraconsistent implicative U3VLSN:

Example 5.42. Let $\Sigma \triangleq \Sigma \sqsupset$ and \mathcal{A} both false-singular and involutive with $(a \sqsupset^{\mathfrak{A}} a) \triangleq \frac{1}{2}$ and $(a \sqsupset^{\mathfrak{A}} b) \triangleq b$, for all $a \in A$ and all $b \in (A \setminus \{a\})$. Then, \mathcal{A} is both \sim -paraconsistent and \sqsupset -implicative. And what is more, $h_{-, \frac{1}{2}} \in \text{hom}(\mathfrak{A}, \mathfrak{A})$. Hence, by Theorem 5.40, C is self-extensional. Now, let $\Sigma' \ni \sim$ be a signature with (possibly, secondary) binary connective \sqsupset , \mathcal{A}' an \sqsupset -implicative canonical \sim -superclassical Σ' -matrix and C' the logic of \mathcal{A}' . Assume C' is self-extensional. Then, by Corollary 5.39 and Theorem 5.40, \mathcal{A}' is false-singular, in which case $D^{\mathcal{A}'} = D^{\mathcal{A}}$, as well as involutive, in which case $\sim^{\mathfrak{A}'} = \sim^{\mathfrak{A}}$, while \mathfrak{A}' is an \sqsupset -implicative intrinsic semi-lattice with bound $\frac{1}{2} = (a \sqsupset^{\mathfrak{A}'} a)$, for any $a \in A' = A$, whereas $h \triangleq h_{-, \frac{1}{2}} \in \text{hom}(\mathfrak{A}', \mathfrak{A}')$. Therefore, by (4.2), for all $a \in A$, $(\frac{1}{2} \sqsupset^{\mathfrak{A}'} a) = ((a \sqsupset^{\mathfrak{A}'} a) \sqsupset^{\mathfrak{A}'} a) = a$. Furthermore, by the \sqsupset -implicativity and false-singularity of \mathcal{A}' , for each $b \in D^{\mathcal{A}'}$, $(b \sqsupset^{\mathfrak{A}'} 0) = 0$, and so $(h(b) \sqsupset^{\mathfrak{A}'} 1) = h(0) = 1$. Likewise, $(0 \sqsupset^{\mathfrak{A}'} b) \in D^{\mathcal{A}'}$, in which case $(0 \sqsupset^{\mathfrak{A}'} \frac{1}{2}) = \frac{1}{2}$, for, otherwise, $D^{\mathcal{A}'} \ni (1 \sqsupset^{\mathfrak{A}'} \frac{1}{2}) = h(1) = 0 \notin D^{\mathcal{A}'}$, while $(0 \sqsupset^{\mathfrak{A}'} 1) = 1$, for, otherwise, $D^{\mathcal{A}'} \ni (1 \sqsupset^{\mathfrak{A}'} 0) = h(\frac{1}{2}) = \frac{1}{2} \in D^{\mathcal{A}'}$, and so $(1 \sqsupset^{\mathfrak{A}'} \frac{1}{2}) = h(\frac{1}{2}) = \frac{1}{2}$. In this way, $\sqsupset^{\mathfrak{A}'} = \sqsupset^{\mathfrak{A}}$. Thus, C' is a \sim -conservative term-wise definitional expansion of C . \square

6. CONCLUSIONS

Aside from quite useful general results and their equally illustrative generic applications (sometimes, even multiple ones providing different insights, and so demonstrating the whole power of universal tools elaborated here) to infinite classes of particular logics, the incompatibility of the self-extensionality of either implicative or both conjunctive and disjunctive finitely-valued logics with unitary equality determinant and the algebraizability (in the sense of [13, 12]) of two-side sequent calculi (associated with such logics according to [14]), discovered here, looks quite remarkable, especially due to its providing a new insight into the non-“self-extensionality

of”/“algebraizability of sequent calculi associated with” certain logics of such a kind proved originally *ad hoc*, and so justifying the thesis of the first paragraph of Section 1. Finally, Subsection 5.2 constitutes foundations of an algebraic theory of U3VLSN. In this connection, taking Theorem 5.34 into account, the most acute problem remaining still open is marking the framework of elimination of disjunctivity stipulation in the formulation of Theorem 4.7.

REFERENCES

1. N. D. Belnap, Jr, *A useful four-valued logic*, Modern uses of multiple-valued logic (J. M. Dunn and G. Epstein, eds.), D. Reidel Publishing Company, Dordrecht, 1977, pp. 8–37.
2. K. Gödel, *Zum intuitionistischen Aussagenkalkül*, Anzeiger der Akademie der Wissenschaften im Wien **69** (1932), 65–66.
3. K. Hałkowska and A. Zajac, *O pewnym, trójwartościowym systemie rachunku zdań*, Acta Universitatis Wratislaviensis. Prace Filozoficzne **57** (1988), 41–49.
4. S. C. Kleene, *Introduction to metamathematics*, D. Van Nostrand Company, New York, 1952.
5. J. Łoś and R. Suszko, *Remarks on sentential logics*, Indagationes Mathematicae **20** (1958), 177–183.
6. J. Łukasiewicz, *O logice trójwartościowej*, Ruch Filozoficzny **5** (1920), 170–171.
7. A. I. Mal’cev, *To a general theory of algebraic systems*, Mathematical Collection (New Series) **35 (77)** (1954), 3–20, In Russian.
8. ———, *Algebraic systems*, Springer Verlag, New York, 1965.
9. G. Priest, *The logic of paradox*, Journal of Philosophical Logic **8** (1979), 219–241.
10. A. P. Pynko, *Algebraic study of Sette’s maximal paraconsistent logic*, Studia Logica **54** (1995), no. 1, 89–128.
11. ———, *On Priest’s logic of paradox*, Journal of Applied Non-Classical Logics **5** (1995), no. 2, 219–225.
12. ———, *Definitional equivalence and algebraizability of generalized logical systems*, Annals of Pure and Applied Logic **98** (1999), 1–68.
13. ———, *Functional completeness and axiomatizability within Belnap’s four-valued logic and its expansions*, Journal of Applied Non-Classical Logics **9** (1999), no. 1/2, 61–105, Special Issue on Multi-Valued Logics.
14. ———, *Sequential calculi for many-valued logics with equality determinant*, Bulletin of the Section of Logic **33** (2004), no. 1, 23–32.
15. ———, *A relative interpolation theorem for infinitary universal Horn logic and its applications*, Archive for Mathematical Logic **45** (2006), 267–305.
16. ———, *Subquasivarieties of implicative locally-finite quasivarieties*, Mathematical Logic Quarterly **56** (2010), no. 6, 643–658.
17. ———, *Many-sorted General Algebraic Logic: Subprevarieties versus Extensions*, Kindle Direct Publishing, 2019.
18. ———, *Four-valued expansions of Dunn-Belnap’s logic (I): Basic characterizations*, Bulletin of the Section of Logic **49** (2020), no. 4, 401–437.
19. ———, *Four-valued expansions of Belnap’s logic: Inheriting basic peculiarities*, EasyChair Preprint no. 3118, EasyChair, 2020.
20. J.A. Robinson, *A machine-oriented logic based on the resolution principle*, Journal of Association of Computing Machinery **12** (1965), 23–41.
21. A. M. Sette, *On the propositional calculus P^1* , Mathematica Japonica **18** (1973), 173–180.

DEPARTMENT OF DIGITAL AUTOMATA THEORY (100), V.M. GLUSHKOV INSTITUTE OF CYBERNETICS, GLUSHKOV PROSP. 40, KIEV, 03680, UKRAINE

Email address: pynko@i.ua