



## 2 x 2 Integer Matrices: Composition of Binary Quadratic Forms

---

Rama Garimella

EasyChair preprints are intended for rapid dissemination of research results and are integrated with the rest of EasyChair.

February 14, 2024

## 2 x 2 INTEGER MATRICES: COMPOSITION OF BINARY QUADRATIC FORMS

Garimella Rama Murthy,  
Professor, Mahindra University,  
Hyderabad, INDIA

### ABSTRACT

In this research paper, we consider  $2 \times 2$  integer matrices and identify interesting binary quadratic forms which naturally arise. Specifically, we consider such symmetric integer matrices and derive compositions of pure binary quadratic forms naturally arising in association with determinant of such matrices. We also, discover number-theoretic results associated with ternary quadratic forms naturally arising in connection with  $2 \times 2$  symmetric integer matrices. We formulate a “generalized Waring problem” using real quadratic algebraic numbers. We also discuss composition of binary quadratic forms naturally arising in other interesting structured  $2 \times 2$  integer matrices. We explore representation of integers using ternary as well as binary quadratic forms. Also, an arbitrary ternary quadratic form is expressed as  $Trace(\bar{X}^2)$  for a  $2 \times 2$  matrix  $\bar{X}$ . Results related to Ramanujan’s ternary quadratic form are explored.

### 1. INTRODUCTION:

Ever since the dawn of civilization, integers stimulated the curiosity of several mathematicians. Algebraic symbolism helped defining negative numbers based on the concept of “ZERO”. Also, linear algebraic equation in one variable with integer coefficients enabled the introduction of rational numbers. Similarly, quadratic equations naturally led to the proposal of novel class of numbers, called “complex numbers”.

Diophantus considered linear algebraic equations in two variables that are constrained to be integers. The so called simplest linear Diophantine equation is of the form

$$ax + by = c$$

was solved using the Greatest Common Divisor (GCD of the integers  $\{a, b\}$ ) algorithm. The book “Arithmetica” by Diophantus and some volumes of Euclid’s elements contained interesting results related to integers and specifically prime numbers, perfect numbers (among other number-theoretic results). Fermat acquired a copy of Arithmetica and contributed several interesting number-theoretic theorems that survived passage of time. For instance, Fermat proved that a prime number of the form  $\{4l+1, l=1,2,\dots\}$  (i.e.  $p \equiv 1 \pmod{4}$ ) can be expressed uniquely as the sum of squares of two integers. Further, a prime of the form  $\{4l+3, l=1,2,\dots\}$  (i.e.  $p \equiv 3 \pmod{4}$ ) can never be expressed as the sum of squares of two integers. This result was combined with the following algebraic identity

$$\begin{aligned}(x_1^2 + x_2^2)(y_1^2 + y_2^2) &= (x_1y_1 - x_2y_2)^2 + (x_1y_2 + x_2y_1)^2 \\ &= (x_1y_1 + x_2y_2)^2 + (x_1y_2 - x_2y_1)^2\end{aligned}$$

resulting in the so called GENUS theorem in algebraic number theory.

The author in his research efforts became interested in  $2 \times 2$  integer matrices. Several interesting results were documented in the technical report [3]. The results reported

in this research paper deal with number-theoretic concepts/ideas applied to  $2 \times 2$  integer matrices.

This research paper is organized as follows. In Section 2, composition of binary quadratic forms arising in the case of  $2 \times 2$  symmetric integer matrices are discussed. In Section 3, ternary quadratic forms naturally arising in association with symmetric  $2 \times 2$  integer matrices are identified and interesting results are derived. In Section 4, arbitrary ternary quadratic form is expressed as  $Trace(\bar{X}^2)$  for a  $2 \times 2$  matrix  $\bar{X}$  whose elements are quadratic surds. In Section 5, composition of binary quadratic forms naturally arising in association with certain structured quadratic forms arising in structured  $2 \times 2$  matrices is discussed. The research paper concludes in Section 6.

## 2. $2 \times 2$ Symmetric Integer Matrices: Sums of Squares of Two Integers: Compositions:

Consider a symmetric  $2 \times 2$  integer matrix of the form  $X = \begin{bmatrix} a & b \\ b & c \end{bmatrix}$

where  $\{a, b, c\}$  are integers. It readily follows that

$$X^2 = \begin{bmatrix} a^2 + b^2 & b(a+c) \\ b(a+c) & b^2 + c^2 \end{bmatrix}. \text{ Thus, we have}$$

$$Det(X^2) = (Det(X))^2 = (a^2 + b^2)(b^2 + c^2) - b^2(a+c)^2.$$

Using the fact that  $Trace(X) = a + c$ , we have that

$$(Det(X))^2 = (a^2 + b^2)(b^2 + c^2) - b^2(Trace(X))^2.$$

Hence,

$$(Det(X))^2 + (b(Trace(X)))^2 = (a^2 + b^2)(b^2 + c^2).$$

Using the standard identity on product of sum of squares of two integers, we have that

$$(a^2 + b^2)(b^2 + c^2) = (ab + bc)^2 + (ac - b^2)^2 = (ab - bc)^2 + (ac + b^2)^2.$$

Thus, genus theorem from algebraic number theory readily applies. Let  $X_1, X_2$  be

two symmetric integer matrices with elements  $\{a_1, b_1, c_1\}; \{a_2, b_2, c_2\}$

respectively. Using the above discussion, we have that

$$(Det(X_1))^2 + (b_1(Trace(X_1)))^2 = (a_1^2 + b_1^2)(b_1^2 + c_1^2)$$

$$(Det(X_2))^2 + (b_2(Trace(X_2)))^2 = (a_2^2 + b_2^2)(b_2^2 + c_2^2)$$

Since the LHS as well as RHS of the above two expressions are binary quadratic Forms, genus theorem can be readily invoked. The binary quadratic form associated with the symmetric matrix,  $X$  is based on the quantity

$$(Det(X))^2 + (b(Trace(X)))^2 = (\mu_1\mu_2)^2 + b^2(\mu_1 + \mu_2)^2, \text{ where}$$

$\mu_1, \mu_2$  are eigenvalues of  $X$ .

**3. 2 x 2 Symmetric Integer Matrices: Sum of Squares of 3 integers ( Trinary Quadratic Form ): Equivalence of Trinary and Binary Quadratic Forms:**

We now consider another interesting quantity associated with a 2 x 2 symmetric integer matrix:  $X = \begin{bmatrix} a & b \\ b & c \end{bmatrix}$ . We readily have that

$$X^2 = \begin{bmatrix} a^2 + b^2 & b(a + c) \\ b(a + c) & b^2 + c^2 \end{bmatrix}.$$

Thus,  $Trace(X^2) = a^2 + 2b^2 + c^2 = \mu_1^2 + \mu_2^2$ , where  $\mu_1, \mu_2$  are eigenvalues of  $X$ , which are in general quadratic algebraic numbers.

- It readily follows that all the 8 possible combinations of components of  $X$  i.e.  $\{\mp a, \mp b, \mp c\}$  will all lead to the same value of  $Trace(X^2)$ .
- The eigenvalues are “complimentary” quadratic surds in the sense that if  $\mu_1 = \delta + \sqrt{\theta}$ , then  $\mu_2 = \delta - \sqrt{\theta}$  with  $\mu_1 + \mu_2 = Trace(X)$ .

Note: We thus have a trinary quadratic form in integers equal to the binary quadratic form in eigenvalues of symmetric 2 x 2 integer matrix. Since  $X$  is symmetric, its eigenvalues are real numbers. We now determine the nature of eigenvalues under some conditions on the integers  $a, b, c$ .

CASE I: Eigenvalues are rational numbers/integers:

$$\mu_1 + \mu_2 = a + c \quad \text{and} \quad \mu_1 \mu_2 = ac - b^2.$$

$$\text{Also, } (\mu_1 - \mu_2)^2 = (a - c)^2 + (2b)^2 = \Delta^2$$

Hence, we have that  $\mu_1, \mu_2 = \frac{(a+c) \mp \sqrt{(a-c)^2 + (2b)^2}}{2}$ . It readily follows that the eigenvalues depend on  $\{a, c\}$  through  $\{a+c, a-c\}$  only.

Thus, if  $\{(a-c), (2b), \Delta\}$  form a Pythagorean triple, the eigenvalues of  $X$  are rational Numbers i.e.  $\mu_1, \mu_2 = \frac{(a+c) \mp \Delta}{2}$ . If  $\{(a+c), \Delta\}$  are even integers, then the both the eigenvalues are integers.

Note:  $\{(a+c), (a-c)\}$  are both even/odd integers. Hence, if  $\{a, c\}$  are both even/odd, the eigenvalues will be integers.

CASE (ii):  $(a - c)^2 + (2b)^2 = p$ , a prime number.

By Fermat’s Theorem,  $p \equiv 1(mod) 4$ .

For any given  $p$ ,  $(a-c)=k$  is unique by Fermat’s Theorem.

In such case,  $a+c = k + 2c$ . Thus,  $\mu_1, \mu_2 = \frac{(k+2c) \mp \sqrt{p}}{2}$ .

Hence, in this case,  $\mu_1, \mu_2$  are real algebraic numbers.

Note: A related case is the one, where  $(a-c)^2 + (2b)^2 = q$ , an integer which is not a perfect square (even or odd number). Even in this case the eigenvalues are algebraic numbers.

- INTERESTING TRINARY QUADRATIC FORM:

We now focus on the following equation from the above discussion:

$$, \quad \text{Trace}(X^2) = a^2 + 2b^2 + c^2 = \mu_1^2 + \mu_2^2$$

We reduce the above equality of trinary and binary quadratic forms ( with  $\mu_1, \mu_2$  being real algebraic numbers ) to the case of equality between two binary quadratic forms under some conditions:

(I)  $\{a, c, d\}$  form a Pythagorean triple

$$\text{Trace}(X^2) = a^2 + 2b^2 + c^2 = d^2 + 2b^2 = \mu_1^2 + \mu_2^2.$$

If the eigenvalues  $\{\mu_1, \mu_2\}$  are integers, then the genus Theorem associated

With binary quadratic forms can be invoked. Also, letting  $X_1, X_2$  be two such  $2 \times 2$  Symmetric integer matrices, we have

$$\text{Tr}(X_1^2) = d_1^2 + 2b_1^2 ; \quad \text{Tr}(X_2^2) = d_2^2 + 2b_2^2.$$

We can readily invoke Brahmagupta's identity for the composition of such quadratic forms

- Bramhagupta's Identity:

For a given  $n$ , the product of two numbers of the form  $a^2 + nb^2$  is itself a number of that form i.e.

$$\begin{aligned} (a^2 + nb^2)(c^2 + nd^2) &= (ac - nbd)^2 + n(ad + bc)^2 \\ &= (ac + nbd)^2 + n(ad - bc)^2 \end{aligned}$$

The identity holds in any commutative ring.

We now invoke the Brahmagupta's identity:

$$\begin{aligned} \text{Tr}(X_1^2) \text{Tr}(X_2^2) &= (d_1^2 + 2b_1^2)(d_2^2 + 2b_2^2) = (d_1d_2 - 2b_1b_2)^2 + 2(d_1b_2 + d_2b_1)^2 \\ &= (d_1d_2 + 2b_1b_2)^2 + 2(d_1b_2 - d_2b_1)^2 \end{aligned}$$

Note: The other interesting cases which lead to composition of binary quadratic forms are

$$a^2 + 2b^2 = e^2 \text{ or } c^2 + 2b^2 = f^2, \text{ for suitable integers } \{e, f\}.$$

- From linear algebra, we readily have that

$$\text{Tr}(X^{2m}) = \mu_1^{2m} + \mu_2^{2m} \text{ for } m \geq 1.$$

As in the case of  $m=1$ , the above expression reduces to interesting polynomial in  $\{a, b, c\}$

- **Fermat's Theorem on Sum of Squares of Two Integers: Pell's Equation: Connections:**

The following Fermat's theorem is well known.

**Fermat's Theorem:** Given a prime,  $p$  of the form,  $p \equiv 1 \pmod{4}$ , it can be uniquely expressed as sum of squares of two positive integers i.e  $p = a^2 + c^2$  for unique positive integers  $\{a, c\}$ . Furthermore, a prime  $q \equiv 3 \pmod{4}$  can never be expressed as sum of squares of two positive integers.

The author observed that for some such primes,  $p$  i.e  $p = a^2 + c^2$ ,  $a + c$  is also a prime,  $q$ . But ' $q$ ' can be such that  $q \equiv 1 \pmod{4}$  or  $q \equiv 3 \pmod{4}$ . Some examples are

$$1^2 + 2^2 = 5 \text{ with } 1 + 2 = 3 ; 2^2 + 5^2 = 29 \text{ with } 2 + 5 = 7 \equiv 3 \pmod{4}$$

Also,  $4^2 + 5^2 = 41$  with  $4 + 5 = 9$ . It can be confirmed by numerical evidence.

We interpret the observation from the viewpoint of  $2 \times 2$  symmetric integer matrices. Let  $\mu_1, \mu_2$  be the eigenvalues of symmetric  $2 \times 2$  integer matrix,  $\bar{X}$  such that

$$\mu_1 + \mu_2 = \text{Trace}(\bar{X}) = q, \quad a \text{ prime number and}$$

$$\mu_1^2 + \mu_2^2 = \text{Trace}(\bar{X}^2) = p, \quad a \text{ prime number. Also let}$$

$$\mu_1 \mu_2 = \text{Determinant}(\bar{X}) = r.$$

It readily follows that  $q^2 = p + 2r$ . Suppose ' $r$ ' is a perfect square of an integer i.e.

$r = b^2$ . We readily have a Diophantine equation  $q^2 - 2b^2 = p$  which is a Pell - type equation.

In view of the above discussion, we want to synthesize a  $2 \times 2$  symmetric integer matrix,

$$X = \begin{bmatrix} a & b \\ b & c \end{bmatrix} \text{ with } a + c = q \text{ and } a^2 + c^2 = p. \text{ If } \bar{X} \text{ is a diagonal matrix, then } \{a, c\} \text{ are}$$

the eigenvalues and Fermat's theorem with the above observation applies.

Now, suppose  $\bar{X}$  is not a diagonal matrix, but it is a singular matrix i.e.  $ac = b^2$ . We readily have that  $(a + c)^2 = q^2 = a^2 + c^2 + 2b^2 = p + 2b^2$ . Thus, in this case Pell-type Diophantine equation naturally follows. We now provide some examples:

$$X = \begin{bmatrix} a & b \\ b & c \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \text{ with } a + c = 5, a^2 + c^2 = 17 \text{ and } b = 2. \text{ In the same spirit, we have}$$

$$\begin{bmatrix} 1 & 6 \\ 6 & 36 \end{bmatrix} \text{ and } \begin{bmatrix} 9 & 6 \\ 6 & 4 \end{bmatrix}. \text{ In all the three examples, we have that}$$

$$a = (\tilde{a})^2, \quad c = (\tilde{c})^2 \text{ and } b = \tilde{a} \tilde{c}. \text{ Hence } L^1 - \text{norm}(\bar{X}) = a + c + 2b = (\tilde{a} + \tilde{c})^2.$$

Hence, in these three examples

$$a + c = (\tilde{a})^2 + (\tilde{c})^2 \text{ is a prime and } a^2 + c^2 = (\tilde{a})^4 + (\tilde{c})^4 \text{ is also a prime.}$$

The above examples motivate the theme of investigating the nature of primes,  $p$  expressible as  $(\tilde{a})^4 + (\tilde{c})^4 = p$ .

In view of the connection between Fermat's Theorem and Pell's equation, we introduce an interesting class of Diophantine equations:

Primal Diophantine Equation: An algebraic equation in which the variables are constrained to be primes is called a PRIMAL DIOPHANTINE EQUATION i.e. integer solutions in a Diophantine equation are constrained to be PRIMES.

We now reason that the Pell-type Diophantine equation that we derived above i.e.

$q^2 - 2s = p = q^2 - 2ac = p$  is never a PRIMAL Diophantine equation (i.e.  $p$  and  $q$  and  $s$  are all never primes) if the eigenvalues of the associated symmetric matrix  $X = \begin{bmatrix} a & b \\ b & c \end{bmatrix}$ , are integers. Trivially  $a = 1$  and  $c = r$ , a prime or  $a = r$  and  $c = 1$ . Since  $a+c$  is constrained to be prime,  $r = 2$ . It readily follows that in such case  $\begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}$  or  $\begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}$  will have eigenvalues that are complimentary quadratic surds.

Thus, such a matrix is the only  $2 \times 2$  symmetric integer matrix with the associated Pell-type Diophantine equation being a PRIMAL Diophantine equation.

- Now, we consider representation of a prime number,  $p$  using the specific trinary quadratic form considered above.

$$\text{Trace}(X^2) = a^2 + 2b^2 + c^2 = p = \mu_1^2 + \mu_2^2.$$

We provide some examples which illustrate the fact that  $p \equiv 1 \pmod{4}$  or  $p \equiv 3 \pmod{4}$ . It readily follows (from Fermat's Theorem) that, if  $\text{Trace}(X^2) = q$ , a prime with  $q \equiv 3 \pmod{4}$ , then,  $\{\mu_1, \mu_2\}$  are real quadratic algebraic numbers (quadratic surds) and not integers.

Example 1:  $\bar{X} = \begin{bmatrix} 1 & 2 \\ 2 & 2 \end{bmatrix}$ . We have that  $\text{Trace}(X^2) = 13 \equiv 1 \pmod{4}$ . Also

$$\text{Trace}(X) = \mu_1 + \mu_2 = 3 \text{ and determinant}(X) = \mu_1 \mu_2 = -2.$$

$$\mu_1 = \frac{3 + \sqrt{17}}{2}, \quad \mu_2 = \frac{3 - \sqrt{17}}{2}$$

i.e. eigenvalues are quadratic surds

Example 2:  $\bar{X} = \begin{bmatrix} 1 & 3 \\ 3 & 2 \end{bmatrix}$ . We have that  $\text{Trace}(X^2) = 23 \equiv 3 \pmod{4}$ . Also

$$\text{Trace}(X) = \mu_1 + \mu_2 = 3 \text{ and determinant}(X) = \mu_1 \mu_2 = -7.$$

$$\mu_1 = \frac{3 + \sqrt{37}}{2}, \quad \mu_2 = \frac{3 - \sqrt{37}}{2}$$

Note: In both the examples, the eigenvalues are NOT integers.

- Suppose the  $2 \times 2$  symmetric integer matrix,  $\bar{X}$  is singular (i.e.  $b^2 = ac$ ). In this case, we have that  $\text{Det}(\bar{X}) = \mu_1 \mu_2 = 0$ . Furthermore,  
 $\text{Trace}(X^2) = a^2 + 2b^2 + c^2 = \mu^2$ .  
*Note:* We can call such 4 integers  $\{a, b, c, \mu\}$  as Pythagorean Quadruples.

Note: If  $\text{Trace}(\bar{X}^2) = p$ , a prime number, then  $\bar{X}$  cannot be singular.

In the above two examples, we realized that if  $\text{Trace}(\bar{X}^2)$  is a prime,  $p$ , it can be such that  $p \equiv 1 \pmod{4}$  or  $p \equiv 3 \pmod{4}$ , when the eigenvalues are real quadratic algebraic numbers. The following Lemma is in the spirit of Fermat's theorem which states that if

$$J^2 + K^2 = q,$$

then  $q \equiv 1 \pmod{4}$  and given 'q';  $\{J, K\}$  are unique.

Lemma: Given  $\bar{X}$  is a  $2 \times 2$  symmetric integer matrix with given trace value and

$\text{Trace}(X^2) = a^2 + 2b^2 + c^2 = \mu_1^2 + \mu_2^2 = g$ , with  $g$  being a prime and  $\mu_1, \mu_2$  being the eigenvalues of  $\bar{X}$  that are "complimentary" quadratic surds (but not integers),  $\{\mu_1, \mu_2\}$  are unique.

**Proof:** The Lemma follows from the fact that  $\text{Trace}(\bar{X})$  and  $\text{Trace}(X^2)$  uniquely determine the pair of eigenvalues  $\{\mu_1, \mu_2\}$ . Q. E. D.

Note: The  $2 \times 2$  matrices  $\begin{bmatrix} a & b \\ b & c \end{bmatrix}$  and  $\begin{bmatrix} c & b \\ b & a \end{bmatrix}$  have the same set of eigenvalues. Hence, given prime  $g$ , the associated  $2 \times 2$  matrix is not unique. In fact, based on the sign, there are 8 associated  $2 \times 2$  symmetric matrices (i.e. with  $\pm a, \pm b, \pm c$ ) which lead to the same value of  $g$ , a prime number. It is possible, to determine the conditions under which two  $2 \times 2$  symmetric integer matrices have the same value of  $\text{Trace}(\bar{X}^2)$ .

The following Theorem deals with  $\{\text{Trace}(\bar{X}^s) \text{ for } s \geq 2\}$ .

**THEOREM:** Let  $\bar{X} = \begin{bmatrix} a & b \\ b & c \end{bmatrix}$  be a  $2 \times 2$  symmetric integer non-singular, non-diagonal matrix (i.e.  $b^2 \neq ac$  and  $b \neq 0$ ) and let  $\{a, c\}$  are both even or both are odd. Also, let  $\{(a-c), 2b, K\}$  form a Pythagorean triple i.e.  $(a-c)^2 + (2b)^2 = K^2$ . Let  $\text{Trace}(\bar{X}^s) = f_s(a, b, c)$  be a trivariate polynomial in  $\{a, b, c\}$ .

Under these conditions,

- (i) we have that  $\text{Trace}(\bar{X}^s) \neq \mu^s$  for any integer  $s \geq 2$ .
- (ii) Also,  $\text{Det}(\bar{X}^s) = \beta^s = (\text{Det}(\bar{X}))^s = (ac - b^2)$  for all  $s$ .



(iii) If  $\bar{X}$  is singular, then  $Trace(\bar{X}^s) = \delta^s$  for all  $s$   
 (where  $\delta$  is the non-zero eigenvalue of  $\bar{X}$ .)

**PROOF:** From the conditions in the statement of theorem (based on earlier discussion) that the eigenvalues of  $\bar{X}$  are integers and  $Trace(\bar{X}^s) = \mu_1^s + \mu_2^s$  for all  $s \geq 1$ .

From Fermat's Last Number Theorem, we have that

$$Trace(\bar{X}^s) \neq \mu^s \text{ for any integer } s \geq 3.$$

Now, let us consider the case of  $s=2$ . It readily follows that (for  $\bar{X}$  non-singular)

$$\mu_1^2 + \mu_2^2 = \frac{(a+c)^2 + k^2}{2}.$$

The RHS in the above equation cannot be an even or odd integer (based on properties of even/odd integers). Thus, the result in (i) follows.

The claims in (ii), (iii) follow from basic linear algebra results Q. E. D.

Note: From the above Lemma it follows that on the manifold/hypersurface  $f_p(a, b, c) - \mu^p$ , there are no integral/rational points (i.e.  $a, b, c, \mu$ 's) for  $p \geq 2$ . But for  $p=1$ , there are infinitely many rational/integral points on the associated plane ( $a+c=\mu$ )

Note:  $f_2(a, b, c) = a^2 + 2b^2 + c^2$ ,  $f_3(a, b, c) = a^3 + 3ab^2 + 3b^2c + c^3$  and

$f_4(a, b, c) = (a^2 + b^2)^2 + 2(b(a+c))^2 + (c^2 + b^2)^2$  i.e.  $f_4(a, b, c)$  is of the same form as  $f_2(a, b, c)$  i.e. weighted sum of three squares of integers. Further, it can be readily reasoned that  $f_m(a, b, c)$  is a sum of squares of 3 integers for any even 'm'.

Note: By the conditions in the above theorem, we have that the associated 4-manifolds  $f_m(a, b, c) - \mu^m$  have no rational/integral points on them for  $m \geq 2$ .

Note: In the spirit of known theme in number theory, we are interested in  $2 \times 2$  symmetric integer matrices for which  $f_2(a, b, c) = a^2 + 2b^2 + c^2$  is a prime number. Also, we are interested in the case where:  $f_m(a, b, c) = \alpha^2 + 2\beta^2 + \gamma^2$  is a prime number (with 'm' being an even number). Thus the functional form of  $Trace(\bar{X}^{2m})$  for m, an integer is the above sum of squares of three integers.

- Suppose the conditions of the above Theorem are violated i.e. the eigenvalue are not integers but are "complimentary quadratic surds". Then, we provide an example where  $\mu_1^p + \mu_2^p = \mu^p$  can happen for  $p = 2$ .

Example:  $\bar{X} = \begin{bmatrix} 3 & 10 \\ 10 & 4 \end{bmatrix}$ . The eigenvalues of  $\bar{X}$  are  $\mu_1 = \frac{7+\sqrt{401}}{2}$ ,  $\mu_2 = \frac{7-\sqrt{401}}{2}$

which are complimentary quadratic surds. It follows that  $\mu_1^2 + \mu_2^2 = 15^2$ .

$$\text{Thus } f_2(a, b, c) = a^2 + 2b^2 + c^2 = f_2(a, b, c) = 3^2 + 2(10^2) + 4^2.$$

But  $f_3(a, b, c) = 2191$  which is not the perfect cube of any integer. We are led to the following conjecture in the spirit of Fermat's Last Number Theorem based on quadratic surds.

**CONJECTURE:** Given that  $\mu_1, \mu_2$  are “complimentary” quadratic surds which are the eigenvalues of a non-singular,  $2 \times 2$  symmetric integer matrix. For any  $p \geq 3$ , we have that

$$\mu_1^p + \mu_2^p \neq \mu^p \text{ for an integer } \mu.$$

We provide the following facts related to the above conjecture. Let the eigenvalues of  $\bar{X}$  be the “complimentary” quadratic surds i.e.  $\mu_1 = \frac{\delta + \sqrt{\gamma}}{2}$ ,  $\mu_2 = \frac{\delta - \sqrt{\gamma}}{2}$ . It readily follows that

$$\mu_1^p + \mu_2^p = \left(\frac{\delta + \sqrt{\gamma}}{2}\right)^p + \left(\frac{\delta - \sqrt{\gamma}}{2}\right)^p = \mu^p = \frac{1}{2^p} \left[ \sum_{\substack{j=0 \\ j \text{ even}}}^p \binom{p}{j} \delta^{p-j} \sqrt{\gamma}^j \right] = \mu^p \text{ for } p \geq 2.$$

In the case of  $p=3, p=4$  we are led to the following Diophantine equations in  $\delta, \gamma$ .

$$p=3 \dots \dots \delta^3 - 4\mu^3 + 3\delta\gamma = 0.$$

$$p=4 \dots \dots \delta^4 + 6\delta^2\gamma + \gamma^2 - 8\mu^4\gamma = 0.$$

The conjecture boils down to proving that the above Diophantine equations have no solutions when the  $2 \times 2$  symmetric integer matrix is non-singular.

4-Manifolds: Reduction to Lower Dimensional Surface: Parametrization using Eigenvalues:

Let  $K_p(\mu_1, \mu_2) = \mu_1^p + \mu_2^p$ . We readily have that  $K_p(\mu_1, \mu_2) - \theta^p = f_p(a, b, c) - \theta^p$ . Thus, using the eigenvalues of  $\bar{X}$  as the variables ( parametrization ), a 4-manifold/hyper surface is reduced to a lower dimensional surface

Note: We generalize the essential idea of above Theorem to arbitrary finite dimensional symmetric square integer matrices. Let B be a symmetric  $N \times N$  dimensional integer matrix. Let the real eigenvalues of such a matrix be  $\mu_i$  for  $i = 1$  to  $N$ . It readily follows that

$$Tr(\bar{B}^p) = \sum_{i=1}^N \mu_i^p. \text{ Also } Tr(\bar{B}^p) \text{ is a function of } M = \frac{N(N+1)}{2} \text{ variables. Let}$$

$$h_p(a_1, a_2, \dots, a_M) = \mu^p \text{ is a manifold/hypersurface in the variables } a_1, a_2, \dots, a_M.$$

Hence, based on the known results in number theory, the number of rational/integral points on such a manifolds/hypersurface can be studied. It also readily follows that

$$h_p(a_1, a_2, \dots, a_M) = \sum_{i=1}^N \mu_i^p = \mu^p,$$

leads to an associated manifold in the variables  $\mu_i$  for  $i = 1$  to  $N$ . We expect interesting connections between number theory, group theory and topology of manifolds.

Note: As specified in the case of  $2 \times 2$  matrices, using eigenvalues of  $\bar{X}$  as the variables ( by parametrization ), an  $M+1$  dimensional hypersurface/manifold is reduced to an  $N+1$  ( lower dimensional ) dimensional hypersurface.

Note: As stated earlier, it can readily be reasoned that ( in the case of  $2 \times 2$  symmetric integer matrices )  $f_p(a, b, c)$  is always a sum of squares of three integers if 'p' is an even number. This fact will compliment the claims of the above Thoerem.

In view of the above results, we formulate an interesting generalization of Waring problem.

**GENERALIZED WARING PROBLEM:**

- In the simplest case, determine the number of ways in which an integer,  $f$  ( from natural numbers ) can be expressed as sum of squares of two real quadratic algebraic numbers,  $\mu_1, \mu_2$  ( quadratic surds ) i.e

$$\mu_1^2 + \mu_2^2 = f .$$

More generally, we are interested in number of representations of the following form;

$$\mu_1^2 + \mu_2^2 + \dots + \mu_M^2 = f .$$

with  $\mu_i$ 's being quadratic algebraic numbers.

In the spirit of above generalization, we can consider weighted sum of squares. Most generally, we consider number of representations of an integer as a general multi-variate polynomial ( could be homogeneous ) in quadratic algebraic numbers. Generalizations in the spirit of Hilbert's 10<sup>th</sup> problem are possible ( with the variables being quadratic surds instead of integers from the natural numbers ).

**4. Quadratic Surd Element based 2 x 2 Symmetric Matrices: Ternary Quadratic Forms: Ramanujan's Ternary Quadratic Form:**

We now reason that a most general arbitrary ternary quadratic form can be expressed in the form  $Trace (\bar{X}^2)$ , where  $X$  is a 2 x 2 symmetric matrix whose elements are quadratic surds.

Consider a 2 x 2 symmetric integer matrices of the form:

$$\bar{X} = \begin{bmatrix} \sqrt{\alpha} a & \sqrt{\frac{\beta}{2}} b \\ \sqrt{\frac{\beta}{2}} b & \sqrt{\gamma} c \end{bmatrix} \text{ and } \bar{X} = \begin{bmatrix} \sqrt{\alpha} a & \sqrt{\frac{\gamma}{2}} c \\ \sqrt{\frac{\gamma}{2}} c & \sqrt{\beta} b \end{bmatrix} . \text{ It readily follows that}$$

$$Trace (\bar{X}^2) = \alpha a^2 + \beta b^2 + \gamma c^2 .$$

Thus, all the results of section 3, can easily be generalized to this most general ternary quadratic form. For instance, the conditions for the eigenvalues to be integers can be readily derived. Details are avoided for brevity. In view of the results of section 3 and the above expression for an arbitrary ternary quadratic form, the themes investigated in number theory ( such as expression of an integer by ternary quadratic form ) can be addressed based on the associated linear algebraic results.

Ramanujan's ternary quadratic form is of the following form

$$a^2 + 10 c^2 + b^2$$

It can be readily expressed as  $Trace (\bar{X}^2)$  of the following matrices i.e.  $\bar{X}'$ s

$$\bar{X} = \begin{bmatrix} a & \sqrt{5}c \\ \sqrt{5}c & b \end{bmatrix} \text{ and } \bar{X} = \begin{bmatrix} a & \frac{b}{\sqrt{2}} \\ \frac{b}{\sqrt{2}} & \sqrt{10}c \end{bmatrix}$$

From the above 2 x 2 symmetric matrices ( not integer matrices), the condition for their determinant to be zero can be readily derived. The results based on the Theorem in section 3, are readily generalizable to the above 2 x 2 matrices. Specifically, the condition for

$Trace(\bar{X}^2) \neq \delta^2$  for any integer  $\delta$  is readily derived. Further, expression for the eigenvalues of the above 2 x 2 matrices can be readily derived in terms of the elements of the matrices. Detailed expressions are avoided for brevity.

### 5. Structured 2 x 2 Integer Matrices: Composition of Binary Quadratic Forms:

We now consider structured 2 x 2 integer matrices. First we consider a 2 x 2 integer matrix, X of the following form:

$$\bar{X} = \begin{bmatrix} a & -\alpha b \\ b & a \end{bmatrix}, \text{ where } \alpha \text{ is an integer.}$$

Such class of matrices reduce to the 2 x 2 matrices representing complex numbers when  $\alpha = 1$ .

It readily follows that the determinant of such a structured matrix is an interesting binary quadratic form:

$$Det(\bar{X}) = a + \alpha b^2.$$

Given two such integer matrices  $\{\bar{X}_1, \bar{X}_2\}$ , it readily follows that

$$Det(\bar{X}_1 \bar{X}_2) = Det(\bar{X}_1) Det(\bar{X}_2).$$

Thus, the RHS in the above equation can be expressed as the interesting binary quadratic form using the Brahmagupta's identity. Details are avoided for brevity.

Now, we consider a structured 2 x 2 integer matrix of the following form:

$$\bar{X} = \begin{bmatrix} a & b \\ b & -a \end{bmatrix}. \text{ It readily follows that } \bar{X}^2 = \begin{bmatrix} a^2 + b^2 & 0 \\ 0 & a^2 + b^2 \end{bmatrix} = (a^2 + b^2) I.$$

It readily follows that compositions of binary quadratic forms can be readily invoked in association with two diagonal matrices  $\bar{X}^2, \bar{Y}^2$  ( associated with structured matrices  $\bar{X}, \bar{Y}$  ).

Infact several number-theoretic results ( such as MATRIX PYTHAGOREAN THEOREM, MATRIX GENUS THEOREM ) can be readily invoked in association with such structured 2 x 2 integer matrices.

### 5. CONCLUSIONS:

In this research paper, several interesting results related to composition of binary quadratic forms arising in connection with 2 x 2 integer matrices are explored. It is expected that these results have interesting implications to algebraic number theory based on quadratic surds. Theorem in Section 3 could also lead to new insights into topology of manifolds.

REFERENCES:

- [1] G. Andrews," Number Theory," Dover Publications, New York
- [2] Harvey Cohn,"Advanced Number Theory," Dover Publications, New York
- [3] G. Rama Murthy,"NP Hard Problems: Many Linear Objective Function Optimization Problem: Integer/Rational Matrices," IIIT-Hyderabad Technical report: IIIT/TR/2016/23, MAY 2016